

RESEARCH ARTICLE

Walking economy at simulated high altitude in human healthy young male lowlanders

Masahiro Horiuchi^{1,*}, Yoko Handa¹, Daijiro Abe² and Yoshiyuki Fukuoka³

ABSTRACT

We measured oxygen consumption during walking per unit distance (C_w) values for 12 human healthy young males at six speeds from 0.667 to 1.639 m s⁻¹ (four min per stage) on a level gradient under normobaric normoxia, moderate hypoxia (15% O₂), and severe hypoxia (11% O₂). Muscle deoxygenation (HHb) was measured at the vastus lateralis muscle using near-infrared spectroscopy. Economical speed which can minimize the C_w in each individual was calculated from a U-shaped relationship. We found a significantly slower economical speed (ES) under severe hypoxia [1.237 (0.056) m s⁻¹; mean (s.d.)] compared to normoxia [1.334 (0.070) m s⁻¹] and moderate hypoxia [1.314 (0.070) m s⁻¹, $P < 0.05$ respectively] with no differences between normoxia and moderate hypoxia ($P > 0.05$). HHb gradually increased with increasing speed under severe hypoxia, while it did not increase under normoxia and moderate hypoxia. Changes in HHb between standing baseline and the final minute at faster gait speeds were significantly related to individual ES ($r = 0.393$ at 1.250 m s⁻¹, $r = 0.376$ at 1.444 m s⁻¹, and $r = 0.409$ at 1.639 m s⁻¹, $P < 0.05$, respectively). These results suggested that acute severe hypoxia slowed ES by ~8%, but moderate hypoxia left ES unchanged.

KEY WORDS: Bipedal locomotion, Optimal speed, Muscle O₂ extraction, Peripheral circulation, Energy expenditure

INTRODUCTION

As humans moved into high-altitude regions over the past 20,000 years, these populations adapted culturally and physiologically to the reduced availability of oxygen in the atmosphere (Beall, 2007). Cardiovascular adaptation to altitude has been primarily studied at rest (Beall, 2007), and comparatively little is known about metabolic responses during walking under hypoxia. Energy costs in Himalayan porters and Tibetan migrants were significantly lower during walking compared to lowlanders (Marconi et al., 2005; Minetti et al., 2006). Similarly, East African women (Maloiy et al., 1986) and Himalayan porters (Bastien et al., 2005) who live at high altitudes can walk and carry heavy baggage as part of their daily lives with unchanged energy cost. Although the underlying mechanisms of the lower and effective energy cost during walking in these populations remain unclear, these results suggest that chronic

exposure-induced hypoxia due to sustained and specific training at high altitudes may cause specialized adaptation in these ethnic groups (Marconi et al., 2005; Minetti et al., 2006). It has also been reported that the arterial oxygen content of Tibetans is markedly lower than that of Andeans, whereas exhaled nitric oxide (NO) concentration, which is a potent vasodilator, is higher compared to Andeans and lowlanders (Beall, 2007). Additionally, native highlanders were found to have higher forearm blood flow and circulating concentrations of bioactive NO products than low-altitude residents (Erzurum et al., 2007). These results may imply that adaptation to high altitude (e.g. oxygen delivery to tissues) may be affected by vasodilator function linked to enhanced blood flow, although this interesting hypothesis has never been applied to lowlanders.

It is well known that there is a U-shaped relationship between oxygen consumption during walking per unit distance (C_w ; ml kg⁻¹ m⁻¹) and gait speeds (v ; m s⁻¹) (Saibene and Minetti, 2003). This indicates that every individual has a particular gait speed that minimizes C_w , which is called the economical speed (ES) (Abe et al., 2008a,b; Horiuchi et al., 2014c, 2015a; Saibene, 1990; Wezenberg et al., 2013); however, to the best of our knowledge, no study has been conducted on the ES of individuals at simulated high-altitude.

At high altitude, it is known that peak aerobic capacity is reduced with a reduction of inspired oxygen pressure (Calbet et al., 2009). Under these conditions, it is possible that ES would be slow, because diminished aerobic capacity causes relatively higher exercise intensity. Traditionally, an individual's ES may be estimated by physical characteristics (Donelan and Kram, 1997), including height (Abe et al., 2008b) and leg length (Horiuchi et al., 2015a). Additionally, preferred walking speed, which was found to be almost consistent with ES in previous studies (Malatesta et al., 2003; Wezenberg et al., 2013), has been suggested to be related to peak aerobic capacity and muscle mitochondrial capacity (Coen et al., 2013).

There are, however, technical limitations to continuously measuring muscle mitochondrial capacity in active skeletal muscles during walking. According to the Fick equation (McArdle et al., 1996), oxygen uptake can be determined by a function of O₂ delivery and O₂ extraction. O₂ extraction occurs at active skeletal muscle, which is defined as an arterial-venous O₂ difference ($a-v$ O₂ difference). As muscle O₂ extraction and skeletal muscle mitochondrial capacity have been related to exercise performance (Jacobs et al., 2013), the continuous measurement of muscle O₂ extraction can provide new information toward a better understanding of the factors that explain individual ES. An alternative approach would be to measure muscle deoxygenation (HHb) derived from near infrared spectroscopy (NIRS), because changes in HHb have been considered a surrogate of microvascular O₂ extraction (DeLorey et al., 2003; Grassi et al., 2003).

In the present study, we sought to investigate the potential impact of hypoxic conditions on an individual's ES during walking. We

¹Division of Human Environmental Science, Mt. Fuji Research Institute, Kamiyoshida 5597-1, Fuji-yoshida-city, Yamanashi 4030005, Japan. ²Center for Health and Sports Science, Kyushu Sangyo University, Matsukadai 2-3-1, Higashi-ku, Fukuoka-city, Fukuoka 8138503, Japan. ³Faculty of Health and Sports Science, Doshisha University, Tatara 1-3, Kyotanabe, Kyoto 6100394, Japan.

*Author for correspondence (mhoriuchi@mfr.i.pref.yamanashi.jp)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Table 1. Cardiorespiratory variables and energy expenditure at resting baseline under each oxygen concentration

	Normoxia	Moderate hypoxia	Severe hypoxia
$\dot{V}O_2$, ml min ⁻¹	296±52	288±28	310±34
$\dot{V}CO_2$, ml min ⁻¹	265±47	260±24	309±38 [†]
\dot{V}_E , l min ⁻¹	11.1±1.2	11.0±1.4	12.4±1.3 [†]
RER	0.89±0.03	0.91±0.06	0.99±0.04 [†]
HR, bpm	82±10	84±9	94±12 [†]
SpO ₂ , %	97.0±0.6	90.9±1.6 [‡]	77.4±5.7 [†]
EE, J s ⁻¹	102±18	100±9	109±12

Values are mean±s.d. $\dot{V}O_2$, pulmonary oxygen uptake; $\dot{V}CO_2$, carbon dioxide output; \dot{V}_E , pulmonary ventilation; RER, respiratory gas exchange ratio; HR, heart rate; SpO₂, arterial saturation; EE, energy expenditure. * $P<0.05$ between normoxia and severe hypoxia, [†] $P<0.05$ between moderate and severe hypoxia, [‡] $P<0.05$ between normoxia and moderate hypoxia.

hypothesized that ES would be slower with a decrease in fractional inspiratory oxygen concentration (FiO₂), and that possible alterations in ES would account for changes in muscle O₂ extraction. To test this hypothesis, in addition to oxygen consumption measurement, muscle deoxygenation profiles (the balance between $\dot{V}O_2$ and \dot{Q} in exercising muscle) at the vastus lateralis muscle were simultaneously measured by NIRS.

RESULTS

Table 1 shows cardiorespiratory variables at resting baseline with different inspired-oxygen concentrations. There were no differences in pulmonary oxygen uptake ($\dot{V}O_2$), and energy expenditure (EE) between the different oxygen concentrations. Meanwhile, carbon dioxide output ($\dot{V}CO_2$), pulmonary ventilation (\dot{V}_E), respiratory gas exchange ratio (RER) and heart rate (HR) linearly increased as concentration of FiO₂ fell ($P<0.001$). These parameters under severe hypoxia were significantly higher compared to normoxia and moderate hypoxia ($P<0.05$, respectively). In contrast, arterial O₂ saturation (SpO₂) linearly decreased with decreasing inspired O₂ concentrations ($P<0.001$), and the differences between conditions were statistically significant ($P<0.05$). During walking, no differences in $\dot{V}O_2$ were observed among conditions at any gait speed, whereas $\dot{V}CO_2$, \dot{V}_E , and energy expenditure (EE) at faster gait speeds under severe hypoxia were significantly greater than normoxia and moderate hypoxia ($P<0.05$, respectively, Fig. 1).

Mean values of the C_w at each oxygen concentration are shown in Fig. 2A. The figure shows that the averaged correlation coefficient values of the C_w - v relationship in the 36 trials of the present study (12 participants×three oxygen levels) were 0.982 (0.937~0.999). Two-way repeated measures analysis of variance (ANOVA) revealed no significant main effect of oxygen concentrations




Fig. 1. Changes in cardiorespiratory variables and energy expenditure (EE) at all gait speeds under normoxia (○), moderate hypoxia (△), and severe hypoxia (●). Values are mean±s.d. * $P<0.05$ between normoxia and severe hypoxia, [†] $P<0.05$ between moderate and severe hypoxia within a same speed, respectively (Tukey *post hoc* test was used). $\dot{V}O_2$, oxygen uptake; $\dot{V}CO_2$, carbon dioxide output; \dot{V}_E , pulmonary ventilation. Error bars indicate s.d.




Fig. 2. Energy cost of walking and economical speed. Energy cost of walking (C_w) at all gait speeds (A); averaged C_w during slower and faster three stages at each oxygen concentration (B); and comparisons in the economical speed (ES) at each oxygen concentration (C). White circles and bars indicate normoxia, gray triangles and bars indicate moderate hypoxia, and black circles and bars indicate severe hypoxia. In A, values are mean \pm s.d. Error bars indicate s.d. * $P < 0.05$ between normoxia and moderate hypoxia, † $P < 0.05$ between moderate and severe hypoxia (Tukey *post hoc* test was used).

($P > 0.05$), while significant main effects were observed for gait speed and interaction ($P < 0.05$, Fig. 2A). Fig. 2B shows the averaged C_w during the slower three gait speeds (between 0.667 and 1.056 m s^{-1}), and faster three gait speeds (between 1.250 and 1.639 m s^{-1}). From ANOVA with linear trend analysis, the averaged C_w at faster three gait speeds linearly increased ($P < 0.05$) among three FiO_2 with no statistical differences in pairwise comparisons. The averaged C_w at the faster gait speeds under severe hypoxia ($3.51 \pm 0.44 \text{ J kg}^{-1} \text{ m}^{-1}$) was $\sim 6.4\%$ greater than normoxia ($3.30 \pm 0.24 \text{ J kg}^{-1} \text{ m}^{-1}$); meanwhile, almost equivalent values were found for C_w during the slower gait speeds in all conditions ($P > 0.05$). Fig. 2C illustrates the comparison of ES under the three oxygen concentrations. ES linearly decreased as concentration of FiO_2 fell ($P < 0.001$). Moreover, ES under severe hypoxia was significantly slower than under normoxia and moderate hypoxia ($P < 0.05$, respectively), while no differences were observed in ES between normoxia and moderate hypoxia ($1.334 \pm 0.070 \text{ m s}^{-1}$ in normoxia, $1.314 \pm 0.070 \text{ m s}^{-1}$ in moderate hypoxia, and $1.237 \pm 0.056 \text{ m s}^{-1}$ in severe hypoxia). Yet there were no statistically significant differences in the coefficients a and b of the quadratic equation among conditions ($P > 0.05$, Table 2).

Fig. 3 shows the relative changes from the standing baseline values in HHb at each gait speed. Although HHb remained unchanged at the slower speeds, it gradually increased in accordance with increasing speed under severe hypoxia; however, it stabilized under normoxia and moderate hypoxia during walking. There were significant differences in relative changes of HHb between severe hypoxia and normoxia above 1.056 m s^{-1} of gait speed ($P < 0.05$). Additionally, significant differences in HHb were observed between moderate and severe hypoxia above 1.250 m s^{-1} of gait speed ($P < 0.05$).

Fig. 4 shows the relationships between ES at each oxygen level, and changes in HHb signals from the baseline values to the last 1 min at each gait speed. Increases in HHb were significantly related to ES for the three faster gait speeds (i.e. above 1.250 m s^{-1}) when the data were pooled.

Table 2. The coefficient a and b derived by the quadratic equation during walking among each oxygen concentration

	Normoxia	Moderate hypoxia	Severe hypoxia
a	$8.83 \times 10^{-4} \pm 1.99 \times 10^{-4}$	$7.92 \times 10^{-4} \pm 2.27 \times 10^{-4}$	$9.00 \times 10^{-4} \pm 2.86 \times 10^{-4}$
b	-0.141 ± 0.028	-0.125 ± 0.036	-0.133 ± 0.038

Values are mean \pm s.d.

DISCUSSION

To the best of our knowledge, this is the first study to examine whether hypoxic conditions alter C_w and ES during walking. The major findings of the present study were threefold. First, averaged C_w at faster gait speeds linearly increased as concentration of FiO_2 fell, although overall C_w under moderate and severe hypoxia was not significantly different from normoxia. Second, ES linearly decreased as concentration of FiO_2 fell and a significantly slower ES was observed only under severe hypoxia; there were no differences in ES between normoxia and moderate hypoxia. Finally, relative changes in HHb at the vastus lateralis muscle from standing to faster gait speeds were inversely correlated with decelerated ES. Collectively, these results suggest that the U-shaped relationship between gait speeds and C_w showed only a leftward shift under severe hypoxia, resulting in a slower ES. In addition, these changes in ES might be partly accountable for local muscle O_2 extraction.

Although no studies have investigated alterations of C_w and ES under hypoxia, hypoxia by itself may reflect relatively higher exercise intensity. However, it is supposed that mechanical energy demands of walking are the same regardless of oxygen concentration, although we did not measure it. Thus, unchanged overall C_w , even under severe hypoxia, may not be a surprising finding. Previous studies reported that $\dot{V}\text{O}_2$ kinetics slowed at the onset of moderate exercise while breathing hypoxic gas mixture (12~15% O_2); these studies, however, also demonstrated that $\dot{V}\text{O}_2$ showed a stable phase 1~2 min after the onset of exercise with no differences between normoxia and moderate hypoxia (DeLorey et al., 2004; Engelen et al., 1996). These results suggested that hypoxia by itself did not affect steady-state whole-body $\dot{V}\text{O}_2$, resulting in unchanged C_w values (Fig. 2A).

Nevertheless, it should be noted that effects of hyperventilation-induced higher $\dot{V}\text{CO}_2$ under severe hypoxia caused higher EE at faster speeds under severe hypoxia than normoxia and moderate hypoxia (Fig. 1). Therefore, it is still possible that C_w may be influenced by higher $\dot{V}\text{CO}_2$. We found significant linear increases in averaged C_w during the faster gait speeds (1.250~1.639 m s^{-1}) as concentration of FiO_2 fell. Specifically, the averaged C_w above 1.250 m s^{-1} under severe hypoxia was 6.4% higher than normoxia despite no significant differences in pairwise comparisons (Fig. 2B). This higher C_w during faster walking would lead to a steeper quadratic curve, resulting in a higher coefficient a . It was notable that the difference in the averaged ES between normoxia and severe hypoxia was about 7.8% (1.334 vs 1.237 m s^{-1}),




Fig. 3. Relative changes from standing baseline values in deoxygenated hemoglobin (HHb) corresponding to gait speed at each oxygen concentration. White circles indicate normoxia, gray circles indicate moderate hypoxia, and black circles indicate severe hypoxia. Values are mean±s.d. * $P<0.05$ between normoxia and moderate hypoxia; † $P<0.05$ between moderate and severe hypoxia; (Tukey *post hoc* test was used).

indicating that a considerable 6.4% higher C_w during the faster gait speeds under severe hypoxia could explain the different ES. Otherwise, as the ES was determined by the coefficients a and b [see Eqn (5) in the method section], greater coefficient a and/or lesser coefficient b resulted in slower ES (Abe et al., 2015). We found no significant differences in the coefficients a and b among conditions, because the coefficients a and b are determined from C_w values observed at six gait speeds. Indeed, the coefficient a under severe hypoxia was higher than normoxia by ~2%, and the coefficient b under severe hypoxia was lower than normoxia by ~6% (Table 2). Taken together, the fact that the slower ES was unchanged in the C_w

under severe hypoxia indicates that the U-shaped relationship shifted leftward only.

Another concern is that statistical power might be lower than we expected. The linear trend analysis was further applied for the data set. We addressed the concern of low statistical power by focusing on the overall slope and fit of the response in ES across the levels of hypoxia, thus reducing the number of comparisons made. This approach was particularly advantageous because changes in ES were subtle among different conditions. However, we must acknowledge that other potential factors in addition to gas exchange variables should be considered to explain the significant




Fig. 4. Relationships between an individual's ES and delta changes in HHb between standing baseline and last 1-min data at each gait speed. Data are pooled in all oxygen concentrations ($n=36$). Data points indicate normoxia (○), moderate hypoxia (△), and severe hypoxia (●). Gait speed of (A) 0.667 m s⁻¹; (B) 0.861 m s⁻¹; (C) 1.056 m s⁻¹; (D) 1.250 m s⁻¹; (E) 1.444 m s⁻¹; and (F) 1.639 m s⁻¹.

linear trend for the ES. For example, higher HR during exercise under hypoxia compared to normoxia, which can cause greater blood flow (BF) to exercising muscles, has been suggested to be a contributor for maintaining similar $\dot{V}O_2$ at each speed (Engelen et al., 1996).

We measured muscle deoxygenation profiles using NIRS to assess muscle O_2 extraction indirectly. It has been suggested that HHb as an indicator of muscle O_2 extraction (DeLorey et al., 2003; Grassi et al., 2003) could reflect the balance between muscle O_2 utilization ($\dot{V}O_{2m}$) and O_2 delivery ($\dot{Q}O_2$, the product of arterial O_2 content and muscle BF). Under severe hypoxia, HHb gradually increased with increasing gait speed, showing significantly higher values above 1.250 m s^{-1} than normoxia and moderate hypoxia. We speculate that an increased leg muscle BF could compensate for a reduction in the arterial O_2 content (CaO_2) under moderate hypoxia, causing similar results in HHb between normoxia and moderate hypoxia. Conversely, under severe hypoxia, greater reduction in CaO_2 could not be compensated by enhanced muscle BF, resulting in greater HHb increase compared with normoxia and moderate hypoxia.

It is well known that reduction of alveolar partial pressure of oxygen limits pulmonary O_2 diffusion capability at high-altitude, which induces a decrease in SpO_2 (Calbet and Lundby, 2009; Schoene, 2001). In the face of impaired pulmonary O_2 diffusion, the rate of peripheral O_2 delivery may play an important role in circulating arterial O_2 . It is thus possible that these differences in peripheral circulation derived by NIRS signals may affect an individual's ES in order to maintain similar $\dot{V}O_2$ at each speed. In the present study, slower ES was related to higher HHb during faster gait speeds. As shown in Fig. 3, HHb increased during faster gait speeds, in particular under severe hypoxia; therefore, the fact that muscle O_2 extraction compensated from reduced $\dot{Q}O_2$ might be partly accounted for by an individual ES.

Methodological considerations

There are several limitations to interpret our results. First, we set 11% FiO_2 as severe hypoxia conditions. We could not completely rule out the effect of this lower oxygen concentration on energy cost. Indeed, $\dot{V}CO_2$ was significantly higher during faster gait speeds under severe hypoxia. These results indicated that a subset of subjects were walking with a potential increase in anaerobic glycolysis to energy turnover; however, a previous study reported that $\dot{V}O_2$ manifested a delayed quasi-steady state even more than lactate threshold (Poole et al., 1988), which was a higher exercise intensity compared to our study. Thus, we believe that our main conclusion may not be strongly affected. In addition, we recruited only healthy young active subjects and performed the experiment only at level gradient. From the viewpoint of clinical implication, e.g. mountain climbing in middle-aged- and aged-populations, such information has not been available at this stage. Future studies should be warranted with various populations as well as with a larger field study.

In conclusion, moderate hypoxia at ~15% O_2 did not affect C_w and ES during level walking in healthy young males. On the other hand, severe hypoxia at 11% O_2 slowed the ES without changing the greater C_w at faster gait speeds compared with normoxia and moderate hypoxia. From observing HHb dynamic profile under severe hypoxia, HHb responses may indicate greater O_2 extraction rather than enhanced hypoxic-induced $\dot{Q}O_2$. Thus, a significantly slower ES might be associated with hypoxic-induced higher $\dot{V}CO_2$ and greater O_2 extraction only at severe hypoxia.

MATERIALS AND METHODS

Participants

Subjects were twelve fit and healthy male athletes (sprinters, middle-distance runners, soccer and baseball players), who engaged in strenuous daily training (2 h per day, 5-6 days per week). Their mean age, height, and body mass were 24 ± 8 years, 1.74 ± 0.06 m, and 70 ± 10 kg, respectively (values are mean \pm s.d.). Researchers explained all procedures, possible risks, and benefits of participation, and obtained written informed consent from each participant. They were asked to refrain from intense physical activity on the 2 days before and from drinking any alcohol and caffeinated beverages the day before testing. This study conformed to the Declaration of Helsinki, and the Mount Fuji Research Institute ethical committee approved all study procedures (No: ECMFRI-03-2014).

Exercise protocols

All experiments were carried out on a motor-driven treadmill, 2.21 m long and 0.88 m wide (T7000, Johnson Health Tech. Co., Ltd, Taichung Hsein, Taiwan). Under all experimental conditions, participants walked on the same treadmill, and they were free to choose their step frequency at each speed. All participants wore underwear, shirts, socks, shorts and lightweight training shoes (Abe et al., 2004). They were allowed to familiarize themselves with treadmill walking while wearing a gas collection mask during at least three preliminary practices on the same treadmill at several gait speeds and gradients (Abe et al., 2004). Inspired oxygen concentrations were set at normobaric normoxia (21%, room air), moderate hypoxia (FiO_2 ; 15%, equivalent to a simulated altitude of 2700 m, at which there is increased risk of acute mountain sickness), and severe hypoxia (FiO_2 ; 11%, equivalent to a simulated altitude of 5000 m, that of the highest permanent human residences on earth). Each oxygen concentration was supplied by 200-L Douglas bag with hypoxic gas generator system (see below 'Measurements') and performed on different days in random order, and a single blind method was used. Participants began by sitting in a chair for 10 min, and then standing for 5 min on the treadmill while baseline values were measured. They then began to walk on the treadmill. Six gait speeds were set incrementally at 0.667, 0.861, 1.056, 1.250, 1.444, and 1.639 m s^{-1} . In keeping with our recent work (Abe et al., 2015), each gait speed was maintained for 4 min.

Measurements

\dot{V}_E and gas-exchange variables were measured by an online computerized breath-by-breath method (AE-310S, Minato Medical Science, Osaka, Japan). Inspired and expired gas volumes were measured using a hot wire respiratory flow system. Flow signals were electrically integrated for the duration of each breath to calculate minute ventilation. The expired fractions of O_2 and CO_2 were analyzed using a zirconium solid electrolyte oxygen analyzer and an infrared carbon dioxide analyzer, respectively. The standard known gases (O_2 15.23%, CO_2 4.999%, and N_2 balance) and room air were used for the calibration of the gas analyzer. Each gas was supplied via a 200-L Douglas bag with a hypoxic gas generator system (Everest summit II, Will Co. Ltd., Tokyo, Japan). Throughout the study, participants' HR was recorded with a wireless HR monitor (POLAR RC800X, POLAR electro, Tokyo, Japan).

Local tissue oxygenation profiles of the vastus lateralis muscle were measured using NIRS (BOM-L1TRW, Omega Wave, Tokyo, Japan), as previously described (Horiuchi et al., 2015b, 2014b). This instrument uses three laser diodes (780, 810, and 830 nm), and calculates relative tissue levels of oxygenated hemoglobin (HbO_2) and HHb according to the modified Beer-Lambert law (Kashima, 2003). NIRS optodes were placed on the lower third of the vastus lateralis muscle (10~12 cm above the knee joint) (Koga et al., 2007). The probe holder contained one light source probe, and two detectors were placed 2 cm (detector 1) and 4 cm (detector 2) away from the source. Hb concentrations received by detector 1 were subtracted from those received by detector 2. This procedure allowed us to minimize the influence of skin blood flow (Ando et al., 2013; Horiuchi et al., 2014a), and to provide a NIRS signal traversing approximately 20 mm, because it has been reported that NIRS signals can reach a half of the depth of the distance between the probe and detector (Patterson et al., 1989).

The thigh muscle, with attached optodes and covering, was wrapped with an elastic bandage to minimize movement of the optodes while permitting

freedom of movement for treadmill walking. Pen marks were made on the skin to indicate the margins of the holder so that optodes could be positioned in exactly the same place for each test. NIRS signals were measured at 1-s intervals throughout the experiment. SpO₂ was monitored by pulse oximeter on the left middle finger every 1 min throughout the study (TM-2564G, A&D, Tokyo, Japan).

Data analysis

Baseline values for all physiological responses (i.e. gas exchange variables, HR, SpO₂, and NIRS signals) were averages of the last 2 min of standing prior to starting walking. A single sample with an average final 1-min pulmonary $\dot{V}O_2$ value (ml min⁻¹) and $\dot{V}CO_2$ (ml min⁻¹) at each gait speed were used to obtain the energy expenditure (EE) during walking, based on the following equation (Brouwer, 1957; Masschelein et al., 2012):

$$EE(J \text{ sec}^{-1}) = [(3.869 \times (\dot{V}O_2/1000)) + (1.195 \times (\dot{V}CO_2/1000))] \times (4.186/60) \times 1000. \quad (1)$$

To calculate each particular C_w , this equation can be transported as follows:

$$C_w(J \text{ kg}^{-1} \text{ m}^{-1}) = EE(J \text{ sec}^{-1})/\text{Body mass}(\text{kg})/\text{speed}(\text{m s}^{-1}). \quad (2)$$

The C_w - v relationship can be mathematically described by the following equation (Abe et al., 2015; Wall-Scheffler and Myers, 2013):

$$C_w(v) = av^2 + bv + c, \quad (3)$$

where the constants a, b, and c are determined by the least squares regressions with the actually observed C_w values at each gait speed. A differential function of the original quadratic Eqn (2) of each individual can be described as follows:

$$C_w'(v) = 2av + b. \quad (4)$$

Then, the individual ES was determined at the gait speed when $C_w'(v)$ equaled zero, that is, the individual ES could be observed using the following equation:

$$ES = -b/2a. \quad (5)$$

We recently reported that standing $\dot{V}O_2$ amounted approximately 50% of the absolute $\dot{V}O_2$ at the level gradient at 0.667 m s⁻¹ under normoxia, indicating that a careful consideration should be necessary to calculate C_w and ES by subtracting the standing metabolic rate (Abe et al., 2015). Indeed, the ES calculated including the standing metabolic rate matches preferred walking speed in many previous studies (Browning et al., 2006; Martin et al., 1992; Malatesta et al., 2003; Peyrot et al., 2012; Wall-Scheffler and Myers, 2013; Wezenberg et al., 2013). With this background, we included the standing metabolic rate to calculate C_w and ES.

A single sample of the mean NIRS signals during the final 1 min at each speed was also analyzed. To compare HHb between participants, the changes in this metric were quantified as percentage changes from the baseline values. Briefly, each resting baseline value, which was represented as an arbitrary unit, was defined as 100%. Thus, changes in NIRS signals at each gait speed were represented as relative changes from baseline values. Similarly, a single sample with an average final 1 min of \dot{V}_E , RER, and SpO₂ was calculated.

Statistics

All data are presented as means±s.d. One-way repeated measures ANOVA with linear trend analysis and pairwise (Tukey) *post hoc* tests were used to evaluate the changes in cardiorespiratory variables at rest, the averaged C_w during the slower and faster three stages, and the ES among different oxygen concentrations. Moreover, two-way repeated measure ANOVA was used to compare changes in the C_w and the NIRS signals between different oxygen concentrations and gait speed within participants. Tukey's *post hoc* test for two-way ANOVA was employed when interactions were significant. To estimate the relationship between changes in ES and NIRS signals, a Pearson correlation coefficient was conducted. Statistical analysis was performed by commercial software packages: Sigma Stat ver 3.5 (Hulinks,

IL, USA) and GraphPad Prism 7 (GraphPad Software, Inc, La, Jolla, CA, USA). A *P*-value of <0.05 was considered statistically significant.

Acknowledgements

We thank all participants for their time and effort. We also offer thanks to Dr Katsuhiko for providing study equipment and to Mr. Shohei Doashi for recruiting participants.

Competing interests

The authors declare no competing or financial interests.

Author contributions

M.H., D.A., and Y.F. conceived and designed the study. M.H., and Y.H. performed experiments. M.H., D.A., Y.H., and Y.F. analyzed data. M.H. D.A., and Y.F. interpreted results. Y.H. prepared tables and figures. M.H. drafted the first manuscript. M.H., D.A., Y.H., and Y.F. edited/revised the manuscript and approved the final version for publication.

Funding

This study was financially supported in part by Japan Society for the Promotion of Science [26440268JP to M.H., 26440266JP to D.A., and 26650175JP to Y.F.].

References

- Abe, D., Yanagawa, K. and Niihata, S. (2004). Effects of load carriage, load position, and walking speed on energy cost of walking. *Appl. Ergon.* **35**, 329-335.
- Abe, D., Muraki, S. and Yasukouchi, A. (2008a). Ergonomic effects of load carriage on energy cost of gradient walking. *Appl. Ergon.* **39**, 144-149.
- Abe, D., Muraki, S. and Yasukouchi, A. (2008b). Ergonomic effects of load carriage on the upper and lower back on metabolic energy cost of walking. *Appl. Ergon.* **39**, 392-398.
- Abe, D., Fukuoka, Y. and Horiuchi, M. (2015). Economical speed and energetically optimal transition speed evaluated by gross and net oxygen cost of transport at different gradients. *PLoS ONE* **10**, e0138154.
- Ando, S., Hatamoto, Y., Sudo, M., Kiyonaga, A., Tanaka, H. and Higaki, Y. (2013). The effects of exercise under hypoxia on cognitive function. *PLoS ONE* **8**, e63630.
- Bastien, G. J., Schepens, B., Willems, P. A. and Heglund, N. C. (2005). Energetics of load carrying in Nepalese porters. *Science* **308**, 1755.
- Beall, C. M. (2007). Two routes to functional adaptation: tibetan and Andean high-altitude natives. *Proc. Natl. Acad. Sci. USA* **104** Suppl. 1, 8655-8660.
- Brouwer, E. (1957). On simple formulae for calculating the heat expenditure and the quantities of carbohydrate and fat oxidized in metabolism of men and animals, from gaseous exchange (Oxygen intake and carbonic acid output) and urine-N. *Acta Physiol. Pharmacol. Neerl.* **6**, 795-802.
- Browning, R. C., Baker, E. A., Herron, J. A. and Kram, R. (2006). Effects of obesity and sex on the energetic cost and preferred speed of walking. *J. Appl. Physiol.* **100**, 390-398.
- Calbet, J. A. L. and Lundby, C. (2009). Air to muscle O₂ delivery during exercise at altitude. *High Alt. Med. Biol.* **10**, 123-134.
- Calbet, J. A. L., Rådegran, G., Boushel, R. and Saltin, B. (2009). On the mechanisms that limit oxygen uptake during exercise in acute and chronic hypoxia: role of muscle mass. *J. Physiol.* **587**, 477-490.
- Coen, P. M., Jubrias, S. A., Distefano, G., Amati, F., Mackey, D. C., Glynn, N. W., Manini, T. M., Wohlgenuth, S. E., Leeuwenburgh, C., Cummings, S. R. et al. (2013). Skeletal muscle mitochondrial energetics are associated with maximal aerobic capacity and walking speed in older adults. *J. Gerontol. A Biol. Sci. Med. Sci.* **68**, 447-455.
- DeLorey, D. S., Kowalchuk, J. M. and Paterson, D. H. (2003). Relationship between pulmonary O₂ uptake kinetics and muscle deoxygenation during moderate-intensity exercise. *J. Appl. Physiol.* (1985) **95**, 113-120.
- DeLorey, D. S., Shaw, C. N., Shoemaker, J. K., Kowalchuk, J. M. and Paterson, D. H. (2004). The effect of hypoxia on pulmonary O₂ uptake, leg blood flow and muscle deoxygenation during single-leg knee-extension exercise. *Exp. Physiol.* **89**, 293-302.
- Donelan, J. M. and Kram, R. (1997). The effect of reduced gravity on the kinematics of human walking: a test of the dynamic similarity hypothesis for locomotion. *J. Exp. Biol.* **200**, 3193-3201.
- Engelen, M., Porszasz, J., Riley, M., Wasserman, K., Maehara, K. and Barstow, T. J. (1996). Effects of hypoxic hypoxia on O₂ uptake and heart rate kinetics during heavy exercise. *J. Appl. Physiol.* (1985) **81**, 2500-2508.
- Erzurum, S. C., Ghosh, S., Janocha, A. J., Xu, W., Bauer, S., Bryan, N. S., Tejero, J., Hemann, C., Hille, R., Stuehr, D. J. et al. (2007). Higher blood flow and circulating NO products offset high-altitude hypoxia among Tibetans. *Proc. Natl. Acad. Sci. USA* **104**, 17593-17598.
- Grassi, B., Pogliaghi, S., Rampichini, S., Quaresima, V., Ferrari, M., Marconi, C. and Cerretelli, P. (2003). Muscle oxygenation and pulmonary gas exchange kinetics during cycling exercise on-transitions in humans. *J. Appl. Physiol.* (1985) **95**, 149-158.

- Horiuchi, M., Endo, J., Takayama, N., Murase, K., Nishiyama, N., Saito, H. and Fujiwara, A. (2014a). Impact of viewing vs. not viewing a real forest on physiological and psychological responses in the same setting. *Int. J. Environ. Res. Public Health* **11**, 10883-10901.
- Horiuchi, M., Fadel, P. J. and Ogoh, S. (2014b). Differential effect of sympathetic activation on tissue oxygenation in gastrocnemius and soleus muscles during exercise in humans. *Exp. Physiol.* **99**, 348-358.
- Horiuchi, M., Muraki, S., Horiuchi, Y., Inada, N. and Abe, D. (2014c). Energy cost of pushing a wheelchair on various gradients in young men. *Int. J. Indus. Ergon.* **44**, 442-447.
- Horiuchi, M., Endo, J., Horiuchi, Y. and Abe, D. (2015a). Comparisons of energy cost and economical walking speed at various gradients in healthy active younger and older adults. *J. Exerc. Sci. Fit.* **13**, 79-85.
- Horiuchi, M., Endo, J. and Thijssen, D. H. (2015b). Impact of ischemic preconditioning on functional sympatholysis during handgrip exercise in humans. *Physiol. Rep.* **3**, e12304.
- Jacobs, R. A., Fluck, D., Bonne, T. C., Burgi, S., Christensen, P. M., Toigo, M. and Lundby, C. (2013). Improvements in exercise performance with high-intensity interval training coincide with an increase in skeletal muscle mitochondrial content and function. *J. Appl. Physiol.* (1985) **115**, 785-793.
- Kashima, S. (2003). Spectroscopic measurement of blood volume and its oxygenation in a small volume of tissue using red laser lights and differential calculation between two point detections. *Optics Laser Technol.* **35**, 485-489.
- Koga, S., Poole, D. C., Ferreira, L. F., Whipp, B. J., Kondo, N., Saitoh, T., Ohmae, E. and Barstow, T. J. (2007). Spatial heterogeneity of quadriceps muscle deoxygenation kinetics during cycle exercise. *J. Appl. Physiol.* (1985) **103**, 2049-2056.
- Malatesta, D., Simar, D., Dauvilliers, Y., Candau, R., Borrani, F., Prefaut, C. and Caillaud, C. (2003). Energy cost of walking and gait instability in healthy 65- and 80-yr-olds. *J. Appl. Physiol.* (1985) **95**, 2248-2256.
- Maloiy, G. M. O., Heglund, N. C., Prager, L. M., Cavagna, G. A. and Taylor, C. R. (1986). Energetic cost of carrying loads: have African women discovered an economic way? *Nature* **319**, 668-669.
- Marconi, C., Marzorati, M., Sciuto, D., Ferri, A. and Cerretelli, P. (2005). Economy of locomotion in high-altitude Tibetan migrants exposed to normoxia. *J. Physiol.* **569**, 667-675.
- Martin, P. E., Rothstein, D. E. and Larish, D. D. (1992). Effects of age and physical activity status on the speed-aerobic demand relationship of walking. *J. Appl. Physiol.* **73**, 200-206.
- Masschelein, E., Van Thienen, R., Wang, X., Van Schepdael, A., Thomis, M. and Hespel, P. (2012). Dietary nitrate improves muscle but not cerebral oxygenation status during exercise in hypoxia. *J. Appl. Physiol.* (1985) **113**, 736-745.
- McArdle, W. D., Katch, F. I. and Katch, V. L. (1996). *Exercise Physiology: Nutrition, Energy, and Human Performance 4th edition*, Alphen aan den Rijn. The Netherlands: Wolters Kluwer.
- Minetti, A. E., Formenti, F. and Ardigo, L. P. (2006). Himalayan porter's specialization: metabolic power, economy, efficiency and skill. *Proc. Biol. Sci.* **273**, 2791-2797.
- Patterson, M. S., Chance, B. and Wilson, B. C. (1989). Time resolved reflectance and transmittance for the noninvasive measurement of tissue optical properties. *Appl. Optics* **28**, 2331-2336.
- Peyrot, N., Thivel, D., Isacco, L., Morin, J.-B., Belli, A. and Duche, P. (2012). Why does walking economy improve after weight loss in obese adolescents? *Med. Sci. Sports Exerc.* **44**, 659-665.
- Poole, D. C., Ward, S. A., Gardner, G. W. and Whipp, B. J. (1988). Metabolic and respiratory profile of the upper limit for prolonged exercise in man. *Ergonomics* **31**, 1265-1279.
- Saibene, F. (1990). The mechanisms for minimizing energy expenditure in human locomotion. *Eur. J. Clin. Nutr.* **44** Suppl. 1, 65-71.
- Saibene, F. and Minetti, A. E. (2003). Biomechanical and physiological aspects of legged locomotion in humans. *Eur. J. Appl. Physiol.* **88**, 297-316.
- Schoene, R. B. (2001). Limits of human lung function at high altitude. *J. Exp. Biol.* **204**, 3121-3127.
- Wall-Scheffler, C. M. and Myers, M. J. (2013). Reproductive costs for everyone: how female loads impact human mobility strategies. *J. Hum. Evol.* **64**, 448-456.
- Wezenberg, D., van der Woude, L. H., Faber, W. X., de Haan, A. and Houdijk, H. (2013). Relation between aerobic capacity and walking ability in older adults with a lower-limb amputation. *Arch. Phys. Med. Rehabil.* **94**, 1714-1720.