

Connectin, a target of homeotic gene control in *Drosophila*

ALEX P. GOULD* and ROBERT A. H. WHITE

Department of Anatomy, University of Cambridge, Downing Street, Cambridge CB2 3DY, UK

*Present address: National Institute for Medical Research, The Ridgeway, Mill Hill, London NW7 1AA, UK

Summary

The homeotic genes of *Drosophila* encode transcription factors that specify morphological differences between segments. To identify the genes that they control, we developed a chromatin immunopurification approach designed to isolate *in vivo* binding sites for the products of the homeotic gene *Ultrabithorax*. Here, we report the analysis of one immunopurified binding site. This 110 bp fragment maps within a regulatory region of a gene under homeotic control, *connectin*. A 4 kb DNA fragment, including the immunopurified binding site, is sufficient to reproduce the appropriate homeotic control within a subset of the full tissue distribution of *connectin*. Analysis of the role of the 110 bp binding site

indicates that it mediates transcriptional controls by *Ultrabithorax* and other homeotic genes. This is the first report of a functional *in vivo* binding site isolated using the chromatin immunopurification method. We also show that the protein product of the *connectin* gene is predicted to be a cell-surface molecule containing leucine-rich repeats. The protein, connectin, can mediate cell-cell adhesion thus suggesting a direct link between homeotic gene function and processes of cell-cell recognition.

Key words: homeotic gene, *Ultrabithorax*, leucine-rich repeat, *Drosophila* development, homeobox.

Introduction

Differences between metameres in *Drosophila* are controlled by a small group of homeotic selector genes, expressed and required in specific anatomical domains (Lewis, 1963; Garcia-Bellido, 1975; Lewis, 1978; Akam, 1987). For example, the *Ultrabithorax* gene (*Ubx*) is expressed predominantly in a domain extending from parasegment (PS)5 to PS13 and is required primarily in the third thoracic and first abdominal segments (Lewis, 1978; Akam and Martinez-Arias, 1985; White and Wilcox, 1985). As long ago as 1975, it was proposed that the homeotic loci select particular developmental pathways by regulating a battery of subordinate target genes and that these target genes are required to coordinate the building of structures characteristic of each segment (Garcia-Bellido, 1975). More recently, the wealth of molecular data on homeotic genes have demonstrated that they encode sequence-specific DNA-binding proteins (Beachy et al., 1988; Muller et al., 1988) capable of activating or repressing transcription from a variety of reporter gene constructs *in vitro* (Thali et al., 1988; Krasnow et al., 1989; Winslow et al., 1989). However, the *in vivo* target genes have been difficult to identify (Andrew and Scott, 1992).

The classical genetic approaches, which have been so successful in identifying the early pattern-forming genes in *Drosophila* development, have so far failed to identify genes that act immediately downstream of the homeotics in the developmental genetic hierarchy, although Kennison

and Tamkun (1988) have isolated a number of loci interacting with dominant homeotic mutations. Recently, Wagner-Bernholtz and colleagues have used P-element-mediated enhancer detection to screen for regulatory elements that respond, either directly or indirectly, to ectopic expression of *Antennapedia* (*Antp*). Using this approach, a previously unidentified gene, *spalt major*, has been identified as a target of negative regulation by *Antp* (Wagner-Bernholtz et al., 1991). In addition, a few promising candidate targets have been identified from amongst the many genes that have previously been cloned. Perhaps the best characterised so far is the *decapentaplegic* (*dpp*) gene, which plays a role in both dorsoventral and proximodistal pattern formation (St. Johnston and Gelbart, 1987). The protein encoded by *dpp* is related to a vertebrate growth factor and functions in the visceral mesoderm in *Drosophila* to induce the expression of another homeotic gene, *labial*, in the midgut endoderm (Immergluck et al., 1990; Panganiiban et al., 1990). Genetic experiments show that the *dpp* gene is positively regulated by *Ubx* and negatively regulated by another homeotic gene, *abdominal-A* (Immergluck et al., 1990; Reuter et al., 1990). However, none of the approaches described so far have been able to demonstrate a direct interaction between homeotic proteins and the regulatory DNA of a candidate target gene.

In an effort to identify direct targets of homeotic regulation, we have developed an immunopurification approach that enriches for short chromatin fragments through their association with endogenous *Ubx* protein (Gould et al.,

1990). In brief, embryonic nuclei are digested with a restriction enzyme and then lysed; the soluble chromatin is affinity-purified against a matrix containing antibodies to *Ubx* proteins; the DNA fragments from the immunopurified chromatin are then cloned. Using this method, at least two candidate *Ubx* in vivo binding sites were isolated. Both target sites (clones 35 and 48) are capable of highly specific binding to *Ubx* proteins in vitro and are located in genomic DNA close to transcription units regulated by *Ubx* in vivo. These two transcripts were therefore presented as good candidates for genes directly regulated by *Ubx* (Gould et al., 1990). We have continued the analysis of these candidates in order to ask, firstly, whether the immunopurified target sites identify regulatory elements mediating homeotic control and, secondly, what can be learnt of the developmental function of these putative target genes.

Here we report the molecular characterisation of the gene, *connectin*, found adjacent to clone 35 and we demonstrate that the short (110 bp) immunopurified fragment is an important component of a *connectin* regulatory element under homeotic gene control. Nose et al. (1992) have independently isolated this gene from a screen of enhancer trap insertions that are expressed in muscle cells. The protein produced by this locus has been named connectin as it may have a role in the formation of connections between certain motor neurons and their target muscles (Nose et al., 1992).

Materials and methods

Fly strains

Flies were maintained at 25°C on cornmeal-agar medium and embryos were collected onto yeasted apple-juice agar plates. Standard genetic markers are as described by Lindsley and Zimm (1985, 1990). Wild-type embryos were of the Oregon R strain. Homeotic null mutations used are as follows: *Df(3R)bx^{d100}*; a deficiency of the entire *Ubx* gene (Lewis, 1978), *abd A^{M1}* and *Ubx^{MX6} abd-A^{M1}* (Sanchez-Herrero et al., 1985) and *Antp^{W10}* (Wakimoto and Kaufman, 1981). Recipients for P-element transformation were homozygous for *cn* and *ry*.

General DNA techniques and sequencing

Standard techniques used are as described by Sambrook et al. (1989). An EMBL3 genomic library made from the iso-1 strain (gift of J. Tamkun) was used to accomplish a genomic walk of about 65 kb at the cytological region 64C. The initial isolation of clones from this region has been described previously (Gould et al., 1990).

A 10 kb *SalI* genomic fragment, detecting the embryonic transcript distribution, was used to screen a *gt10* cDNA library constructed from 3-12 hour embryonic poly(A)⁺ RNA. Subsequently the longest hybridising clone, *gt35.8* (2.6 kb), was used to screen two independent size-selected plasmid cDNA libraries, one from 8-12 hour embryos and the other from 12-24 hour embryos (Brown and Kafatos, 1988); 40 primary isolates were characterised and all were found to be of one class but with differing 5' and 3' extents. Sequence analysis shows that the 5' end of pNB35.8 is identical with that of two other independent cDNA clones, one isolated from a separate library. However, the 3' extents of several clones sequenced were found to be unique and in no case was a poly(A) tract found. It therefore seems likely that additional (but short) 3' message sequences are not included within pNB35.8.

cDNA and genomic clones were sequenced by the chain termination method (Sanger et al., 1977) using double-stranded templates and Sequenase (USB). Large regions were covered using nested deletions together with primer synthesis. Nested deletions of *gt35.8* were created using the exonuclease III/mung bean nuclease system (Stratagene) and sequenced as single-stranded M13 templates using Klenow polymerase.

Sequences were analysed using the Staden (Staden, 1982) and GCG (Devereux et al., 1984) software.

Whole-mount in situ hybridisations using digoxigenin-labelled DNA probes were carried out according to Tautz and Pfeifle (1989) with some modifications (Gould et al., 1990). For hybridizations using RNA probes, the digoxigenin-labelled RNA probes were made according to manufacturers instructions (Boehringer-Mannheim, RNA labelling kit) and the in situ hybridization carried out as for the DNA probes except that the prehybridizations, hybridizations and post-hybridization washes were at 70°C.

Transformation constructs

P-element mediated germ-line transformation of *cn ry* embryos was according to Spradling (1986) but using pUCHs 2-3 (gift of Ernst Hafen) as a constitutive source of transposase. Regulatory regions of the *connectin* gene were cloned into the blunted *XbaI* site of pHZ50PL (Hiromi and Gehring, 1987).

The 10 kb *SalI* genomic fragment was inserted into pBluescript (Stratagene). From this 10 kb subclone, a 4 kb *EcoRI* fragment (representing the 4 kb *EcoRI-SalI* genomic interval) was excised, blunted with Klenow polymerase, and then inserted into pHZ50PL to give the 4 kb construct. The 1 kb *Pst-SalI* genomic region was subcloned into pBluescript and excised as a blunt *SmaI-HincII* fragment for insertion into pHZ50PL yielding the 1 kb construct.

Precise deletion of clone 35 sequences (a 110 bp *HaeIII* fragment; (Gould et al., 1990)) was accomplished from a 0.7 kb *PstI-SacI* genomic subclone using the Inverse Polymerase Chain Reaction with Pfu DNA polymerase (Stratagene) at high stringency. Primers were chosen to generate an additional GC dinucleotide at the deletion junction, thus converting the two *HaeIII* half-sites into a diagnostic *ApaI* site. Sequencing confirmed that no inadvertent mutations had been introduced. A 0.4 kb *SacI-SpeI* fragment, containing the deletion of clone 35, was exchanged with the wild-type 0.5 kb *SacI-SpeI* fragment in both the 4 kb and 1 kb constructs to give the 4 kb-35 and 1 kb-35 constructs, respectively.

X-gal staining

X-gal staining was performed as described (Protocol 76; Ashburner, 1989b), except that 5% glutaraldehyde in PBS was used as the fixative. After the staining reaction, embryos were mounted in PBS, 10% glycerol and 0.1% Triton X-100 and photographed using differential-interference or bright-field optics.

Results

Cloning and sequencing of the connectin gene

The immunopurified 110 bp clone 35 was used as a probe to initiate a genomic walk at the cytological location 64C. Screening by in situ hybridization with genomic probes from this region revealed a transcription unit under homeotic gene control (Gould et al., 1990; Fig. 1). To characterize this 35 transcription unit, we isolated many cDNAs covering the cloned interval, the longest being approximately 4.5 kb in length (pNB35.8). This is close to the size of the major polyadenylated RNA species (approximately




Fig. 1. Organisation of the 35 transcription unit. The positions of restriction sites for *Hind*III (H), *Eco*RI (R), and *Sal*I (S) are indicated above a line representing the genomic walk (scale above line in kb). Restriction fragments hybridising to cDNA clone pNB35.8 (see text and Materials and methods) are shown as solid bars beneath the walk with the 5' to 3' direction indicated. The location of the 110 bp clone 35 used to initiate this genomic walk is indicated, as are the extents of the lambda phage covering the cloned region. The position of the *Eco*RI-*Sal*I fragment used in the 4 kb transformation construct is also shown.

5 kb) detected on blots of 0-24 hour embryonic RNA using probes from this region (data not shown). Using this cDNA as a probe in embryo whole-mount in situ hybridisation, the same transcript distribution was obtained as had been seen previously with genomic probes.

To define the 5' and 3' limits of any open reading frame (ORF) present, several different cDNA clones were transcribed and translated in vitro. This indicated that both gt35.8 (2.6 kb) and pNB35.8 (4.5 kb) cDNAs encode a similarly sized polypeptide of about $72 \times 10^3 M_r$ (data not shown). Sequencing of the gt35.8 cDNA revealed an ORF of 2046 nucleotides (ORF:35), encoding a polypeptide of 682 amino acids ($76 \times 10^3 M_r$; Fig. 2). The codon usage found within this ORF:35 agrees well with that predicted for a *Drosophila* gene (Ashburner, 1989a). The second methionine codon of this reading frame is predicted to be slightly more favourable as a translation start than the first methionine codon, lying 27 nucleotides upstream (Cavener, 1987). Hydropathy analysis (Fig. 3A) indicates that both the amino and carboxy termini are predicted to be highly hydrophobic. The amino-terminal hydrophobic region of 26 amino acids is characteristic of a signal sequence (von Heijne, 1986). There is no likely transmembrane domain but the carboxy-terminal hydrophobic region of 25 residues is consistent with the presence of a glycosyl-phosphatidyl inositol linkage to the cell membrane (GPI-anchor; Ferguson and Williams, 1988). The existence of a GPI anchor has been confirmed by the phosphatidylinositol specific-phospholipase C induced release of this protein from transformed *Drosophila* S2 cells expressing this ORF:35 (L. Meadows, A. P. G., R. A. H. W., unpublished results).

The most striking feature of the predicted polypeptide is the presence of 10 copies in tandem of a 24 amino acid leucine-rich repeat (LRR; Fig. 3B,C). This motif has a characteristic spacing of hydrophobic residues, in particular leucine and is found in many evolutionarily and functionally diverse proteins (Takahashi et al., 1985; Kataoka et al., 1985; Titani et al., 1987; Fisher et al., 1989; Reinke et al., 1988; Hashimoto et al., 1988; Rothberg et al., 1990). The consensus motif for the LRRs of ORF:35 is shown in Fig. 3B. In several molecules containing blocks of LRRs, there

are also conserved flanking regions containing cysteine clusters (Lopez et al., 1987; Hickey et al., 1989; Schneider and Schweiger, 1991). Mutations affecting cysteine residues in a carboxy-terminal flanking region of the *Drosophila* LRR-protein Toll give rise to dominant *Toll* mutations (Schneider et al., 1991). ORF:35 shows only very limited homology to these conserved flanking sequences; however, there are cysteines at positions 100, 110, 112 and 126 followed by the sequence DTT occurring just on the amino-terminal side (137-139) of the LRR block and ICDCRL on the carboxy-terminal side (409-413). The other notable feature of the sequence of ORF:35 is that the amino-terminal region preceding the LRR block is serine-rich.

Comparison of the sequence of ORF:35 with that from a gene isolated by Nose et al. (1992) revealed identity and we refer to this gene as *connectin*.

The expression of the connectin gene

Before considering the regulation of the *connectin* gene, we first discuss its expression pattern in more detail than has been presented previously (Gould et al., 1990). Transcripts are first detected by in situ hybridization in the extended germ band (stage 10; stages according to Campos-Ortega and Hartenstein, 1985) in a subset of the splanchnopleura, the cell layer that will subsequently give rise to the visceral mesoderm (Fig. 4A). There is expression in the presumptive foregut and hindgut mesoderm and in a segmentally repeated subset of the presumptive midgut visceral mesoderm. The labelling in the mesoderm around the foregut and hindgut persists but the expression in the midgut visceral mesoderm subsets narrows to small segmentally repeated patches by the end of stage 11 (Fig. 4B) and disappears after germband retraction. In late stage 11, *connectin* RNA appears in the ventral nervous system and in the somatic mesoderm (Fig. 4B). The labelling in the ventral nervous system is initially in two longitudinal bands of cells with little segment-to-segment modulation in intensity. In contrast, the initial expression in the somatic mesoderm is highly modulated, being strong in the three thoracic segments and relatively weak in abdominal segments. There

1	GATTAGTCTCGTTAGGCGGTCTTTGTGTCAACAACGTGTGTGCTCGCTCCCGGTGAAAAATCGAAAAACCCAGCAATCGCCCTCGCGATTTTCGGCTCGGTTTTTCCACAGGACTAACG	120
121	ACCAAAGTGCTTGTGTGTTTTTTCGGCCCCATAAGCAAGTGTGCGTGCCAAAGTGCAACAAGTCTTAAGCGTAATTTAATGCAACAATTTCCGCCGTTTCCACATTTCCCGAATTTGA	240
241	ACAAGTGAAAAACGACTCGATCGTATCGCCAGGAAGATATATCAGTGTGCGAGTGTTCGTGCAAAAGTGGAGCATTATCTCCAAAGAAAACAGAAAAAAGAAAAACAGTCAAATAC	360
361	ATTGCATTAGTACAGCAGATCGAGAACTCTGGTGTACGTGACTTGGCCTTTTCCAGACTCAATCGAAAAGTCCGCTCTTCCGCCGAGCGATAGGAATAAGCTCGCCATGTTGCCGCAT	480
481	CATCAGCGCCATAAAATGGCGACTTTGGCGGACAGCGCATTGTTTCTGCTCCTTTCTGCTCTGCTAATTTGGCGCTGCGCTGGTTCACGCCACCGAAGGACGCGCAAAAGATGACCGG	600
1	M A T L A D S A I C F L L L S L L L I G A C L V T P T E G R A K D D R	35
601	CGCACCAGGGGGCGTGGCAGCAGCAGTGGCGTCTCTCGTCTCCTCCAGCAGCAGCAACAACATGAACAATGGCTACTATAGTGGCTCCTCGTCGACAGCGGGCTCCTCCTCGGCTAT	720
36	R T R G R G S S S G V L S S S S S S N N M N N G Y Y S G S S S T A G S S S G Y	75
721	GTGTTCCACGACAGTAGTGGCGTGAACCTCGGAAGCACTGGCTACAGCGGACCGATGGACAGCAGTGGATTCTGCACCCGGCGGCTGACATGAAGCTAATGTGCTACTGTACCCCGGAT	840
76	V F T S S S A V N S G S T G Y S G P M D S T G F C T R R R D M K L M C Y C T P D	115
841	GAGAACCACGTGCGAGTGCAGAAGCCGAGTGTGGGTGTTCTCCGAGGATTGCATCAGAATGACACCACCTGGACCGGATTCTATCAGCAGAAGCGTTTGGCGGAAGTGAAGTTTGTG	960
116	E N H V P V Q K A E C W V F S E G G L H Q N D T T W T R F Y Q Q K R L R E L K F V	155
961	ATCCAAAACAAATGCCGATTTGGACTATATACCACCATGATAATGAACCGCTGAAGAACCTGAGCAGCATTGTGATGAATACTCGCAGGTGGAGATAGTCAAGAGCTATGCCCTTCGCA	1080
156	I Q N N A R L D Y I P T M I I E P L K N L S S I V I E Y S Q V E I V K S Y A F A	195
1081	AATCTGCCCTTCTTGGAAAGATTATCTTGAATAACAATCACATAATGGCCCTCGACCAGGATGCCCTCGCCAATCATATCAGGCTGAGGAAATGAACCTGGAGCACAACCAGATCTTC	1200
196	N L P F L E R I I L N N N H I M A L D Q D A F A N H I R L R E L N L E H N Q I F	235
1201	GAGATGGATCGCTATGCATTCGGTAATCTGCCACTTTGCGAACGGCTCTCCTGAATAACAACAATATCAGCAGCTTTCAGCGGGTCTATTCGCTGATATGGCCAGGCTAACCTTCCTG	1320
236	E M D R Y A F R N L P L C E R L F L N N N N I S T L H E G L F A D M A R L T F L	275
1321	AACTGGCGCACAACCAGATCAATGTGCTGACCGCAGAGATTTTCCGAGGACTGGGCAATCTAAATGTGTGAAGCTGACCCGGAATAATTTGAACCTTATCGCGGACTGTCTTTGGC	1440
276	N L A H N P V Q K A E C W V F S E G G L H Q N D T T W T R F Y Q Q K R L R E L K F V	315
1441	GAATTTGGAGCCTGAGTGAAGCTGAGCAGATAATCGCATTTGAGCGCAITTCGAGCGCTCTGGATGGCTGAATACACTGAAAACACTGAATCTGCGAAAACACTGTGTAAG	1560
316	E L W S L S E L E L D D N R I E R I S E R A L D G L N T L K T L N L R N N L L K	355
1561	AAGATCGCAATGGCTGCTCGCGGGAACCCCGCCCTGTGTCCATCAACGTGACAGGCAAAACAAATTTGAAAACGCTGACGTTCTACACATTCCAGCCAATTTGGACAATTTGGTCAAC	1680
356	K I D N G L L R G T P A L L S I N V Q A N K L E T L T F Y T F Q P I M D N L V N	395
1681	AGCACCAGCGAGCTGCTGTGTGACAGCAACAATTCATTGCGACTGTGCTGTGAGTGGATCTCGAATGAAAAATCGCACAGTCACTGCAGCTGCGTGAATCGCTGGAGGACCTG	1800
396	S T S E L L V S D N K F I C D C R L Q W I F E L K N R T R H L Q L R D S L E D L	435
1801	CACTGCACGCTGACAGGACGCGAAACTGTGCGCACTTTGTTGGACCCCGTCCCGCCGACCATCTGGAGTGTCTCAATATCGGGGATTCACGGCGATTGGCAGCAATAGCCGACGATGGGC	1920
436	H C T L Q E P K L S H F V D P V P P T I L D V L N I G S N L N F I G S D T V F A	475
1921	GGTGTGGCAACACGCTGGTGGCAGCAGCTACAGCGCTTACTACTGGACAGCAGGAAAGCATTGGGCGAGCAGATCCCGACAGCAGTCCGGGACAGCGGAGTTCGCTCGAGT	2040
476	G V G N S V V G S S Y S G L T M D D S R K H L G S R S R Q A L R G Q R Q F A S S	515
2041	CGCGAGAACGTGGTGGAGTCCAAGATCGCGAGGAGGAGGAGCGCAGGAGGAGTCAAGGAGAAGGACCTGGCGGAGTGGCTCCGGCTCACAAGCGATATGACTACTACGATGACAAC	2160
516	A E N V V E S K M R R R R K R Q E E V K E K D L A A V A P A H K R Y D Y Y D D N	555
2161	AACGGAGCAGTCCCTGGGTATGCGCTAGACCTGGATGACAATCTTAGCCTGCACAAGCAGGAGTCTATGGTCCGGTTCTCCGCTAGTGGGTGACAAACGATGGATGCTCGATG	2280
556	N G G M S L G H G L D L D D N L S L H K Q G F Y G A G S P V V G H N D V D V L M	595
2281	ACATCGCACTCCGCGTCCGGAGACCCCTCGACITCTGACCACCAAGAAGCAATCTACATCAAGTTGTTCTGCTGAAGCCGGAAATGCTGCCCTGCCACGATGGACTCAGCGATCCC	2400
596	T S H S A S G D A L I L T K N A I Y I K L N L K P E M L P C H D G L S D P	635
2401	ACCGAGCTGCCCTCTCCCGCATCTGATGGATGTGCGCAGCAATGTGGCCAGGACATGTGACGGCAGGGCAAATTTCTTGGCCAGGGCATGACATCATAGTGTAGTCTAGTCGCA	2520
636	T E L P L S R D L M D V R S N V G Q D M S T A G A N S L A O G M T I I V S L V A	675
2521	CTGATGATGATCAGCGGGGTTGAAGACTCAACTCCGAGAATGGTGTGATGGTACCCCGACGTTTGGTTCTACATTGTACATACTGTCTTGGAGATGGTGCCTTGGGAAACGA	2640
676	L M M I S R G	682
2641	GCTTCAACACGTTAATACGAAAGATACCAGCTACAGTCTGTAATATGAAAAAGGGTAACAAAATTCGATAAAATTTAATACTATAAACCTAGCTGATAGCTGTAATGATAGACTCG	2760
2761	ACTTTTTCAACAACCTTCAGCAGCAGATTGAGATTGAGATATAGAAACCGATACAAAACCATTTACCGAACTATAAGCCAAAACGAAACCAAAGAGAACCTCCGAGCAAACTGGCGTT	2880
2881	TCATTTAAAAAGAAAGATACTCAACTTATCGCAGGATGATGACTACCACTACTACTACATAGAATTAGTTTAAACGTTAGCCATAGT	2973

Fig. 2. Nucleotide and predicted amino acid sequences of the 35 cDNAs. The sequence of nucleotides 1 to 343 is from cDNA pNB35.8, whilst nucleotides 344 (arrowhead) to 2973 are from the overlapping clone gt35.8 (see Materials and methods). Conceptual translation is initiated at position 496 but the upstream ATG at position 469 is indicated by double underlining. Amino and carboxy terminal hydrophobic stretches of amino acids are underlined and potential N-linked glycosylation sites are shown in bold. This sequence has been submitted to the EMBL sequence data base (accession number X68701).

are differences, too, between the thoracic segments with the labelling in T1 being slightly weaker and having a different arrangement of labelled cells (Fig. 4C,D). By stage 14 (Fig. 4E,F), the intensity differences between segments in the somatic mesoderm are less pronounced but there are clear differences in the arrangement of mesodermal cell groups between thoracic and abdominal segments. From studies with antibodies against the *connectin* protein, the thoracic ventrolateral clusters correspond to the precursors of the pleural external transverse muscles, whilst in the abdomen the ventral clusters correspond to the ventral superficial oblique muscles and the lateral clusters to the

pleural external transverse group (L. Meadows, A. P. G., R. A. H. W., unpublished observations; see also Nose et al., 1992). The labelling in the mesoderm declines shortly after stage 14. In the ventral nervous system, the initially uniform labelling is succeeded by a more modulated pattern. Differences between the thoracic and abdominal neuromeres appear at stage 14 and, by stage 16, there is strong labelling in the anterior central nervous system, including the brain lobes and the thoracic neuromeres but little labelling in most of the abdominal neuromeres. Labelling is also seen associated with the gnathal sense organs from stage 11 onwards.




Fig. 3. Analysis of the *connectin* protein precursor. (A) Hydropathy plot using the method of Kyte and Doolittle (1982). Significant regions of hydrophobicity (scoring positively) are found at both the amino and carboxy termini. (B) Alignment of individual LRRs showing highly conserved amino acids boxed. Bracketed numbers to the right indicate the position of each 24 amino acid LRR within the protein precursor. A consensus sequence for the 10 LRRs is shown below. (C) Schematic representation of the amino acid sequence of the protein precursor. Amino (N) and carboxy (C) terminal hydrophobic regions are indicated by black boxes. Shaded boxes show the position of individual LRRs and asterisks mark potential N-linked glycosylation sites.

Transcription driven by a 4 kb region encompassing clone 35

The *connectin* gene was isolated as a transcription unit neighbouring the immunopurified clone 35, a candidate *in vivo* binding site for *Ubx* protein. We have previously shown that *connectin* expression is under the control of *Ubx* and other homeotic genes (Gould et al., 1990). If the immunopurification has identified a functional *Ubx* protein binding site, it should be possible to demonstrate that the 110 bp clone 35 sequence mediates *Ubx*-dependent regulation of *connectin*. The approach used was first to ask whether the immunopurified fragment had led us to a *connectin* regulatory element under homeotic gene control. We considered it unlikely that the 110 bp sequence would function efficiently by itself as an enhancer element and so larger genomic restriction fragments spanning the clone 35

region were tested for their ability to drive expression of a *lacZ* reporter gene from an *hsp70* minimal promoter in transformed embryos (Hiromi and Gehring, 1987; Fig. 5). In Fig. 6, we show that the 4 kb *SalI-EcoRI* fragment encompassing clone 35 is sufficient to give a consistent pattern of *lacZ* expression corresponding to a subset of the total *connectin* expression pattern (compare Fig. 4D-F and Fig. 6A-C). (We note that this element failed to give the same staining pattern when placed upstream of the *hsp70* promoter in the opposite orientation and therefore does not behave like a true enhancer.) High levels of reporter gene expression were detected, from the extended germ band stage onwards, in small groups of cells in the gnathal and thoracic segments and in a posterior segment, probably A9 (Fig. 6A-C). The pattern of gnathal staining correlates with the developing gnathal sense organs, which express *connectin* RNA, and the thoracic labelling correlates with the ventrolateral *connectin* expression in the somatic mesoderm described above for stage 13 embryos (Fig. 4E,F). One high-expressing line, 4.40, also showed some abdominal labelling (Fig. 6A) which correlated well with the positions of the ventral and lateral clusters of *connectin* expression in the abdominal somatic mesoderm. In marked contrast to the endogenous *connectin* RNA distribution, no staining is seen in the central nervous system (Fig. 6B). Staining in gnathal segments was first detectable in stage 10 embryos whilst the thoracic and A9 labelling was not observed until slightly later at stage 11. Thus the 4 kb construct gives a subset of the complete *connectin* pattern with predominant expression in gnathal and thoracic segments.

Homeotic regulation of the 4 kb region

The clear differences in expression between thoracic and abdominal segments observed with the 4 kb construct suggested that the 110 bp immunopurified sequence had led us to a regulatory element under homeotic gene control. The reduced levels of expression of the 4 kb construct in abdominal segments A1-A8 suggest a down regulation by *Ubx* and the abdominal homeotic genes. In order to test this, we examined the expression in embryos homozygous for *Ubx* or *abd-A* null mutations and also in embryos homozygous for the *Ubx abd-A* double mutant combination. In *Ubx* mutants, expression is derepressed in the abdominal segments, A1 and A2 (Fig. 6D). This is consistent with the requirement for *Ubx* function in both A1 and A2 somatic musculature (Hooper, 1986). In *abd-A* mutants, there were no consistent changes in the pattern of expression of the 4 kb construct (data not shown) but, in the *Ubx-abdA* double mutant, the derepression is more dramatic than that seen in embryos mutant for *Ubx* alone: high levels of expression are now seen in A3 and, in late embryos, intermediate levels of expression are detected further posteriorly (Fig. 6E). This indicates that, in the wild type, both *Ubx* and *abd-A* repress this construct. The explanation for the lack of effect of the *abd-A* single mutant on the expression pattern is likely to be transregulation between *abd-A* and *Ubx*. The *abd-A* gene normally represses *Ubx* expression in the posterior abdominal segments and, in *abd-A* mutants, *Ubx* expression in the abdomen is derepressed (Struhl and White, 1985). Thus, in the *abd-A* single mutant, high levels of *Ubx* expression are




Fig. 4. Expression of *connectin* RNA in wild-type embryos. (A) Stage 11; lateral view. Labelling is seen in segmentally repeated subsets of the splanchnopleura, which will form the visceral mesoderm of the midgut, and in the visceral mesoderm of the foregut (fgm) and hindgut (out of focus). (B) Later stage 11; lateral view. Strong labelling is seen in the visceral mesoderm of the foregut (fgm) and hindgut (hgm) and the arrowheads indicate two of the segmentally repeated subsets of labelling in the presumptive visceral mesoderm of the midgut. The predominant labelling in the three thoracic segments T1, T2 and T3 is in the somatic mesoderm. There is also weaker labelling in cells of the somatic mesoderm in the abdominal segments. The arrow indicates labelling in one of the longitudinal tracts of cells in the ventral nervous system. (C) Stage 12; lateral view focussing on the somatic mesoderm. Labelling is predominantly in specific cell clusters in the thoracic somatic

mesoderm. (D) Stage 12; ventral view. The two longitudinal tracts of cells in the ventral nervous system are stained with uniform intensity along the anteroposterior axis. In the somatic mesoderm, there is a clear difference in labelling intensity between the thoracic and abdominal segments. Also, T1 is weaker than T2/T3 and shows a different arrangement of labelled cells. The arrow indicates labelling associated with gnathal sense organ precursors. (E) Stage 14; dissected preparation. The labelling in the ventral nervous system (vns) shows the beginnings of segment-to-segment modulation with the thoracic neuromeres showing slightly higher intensity labelling than the abdominal. In the somatic mesoderm, the intensity differences between thoracic and abdominal segments are less pronounced than at earlier stages. There are clear differences in the arrangement of the labelled cell clusters between thoracic and abdominal segments. (F) Stage 14; lateral view focussing on the somatic mesoderm showing the positions of the labelled cell clusters. The labelling in the T1 somatic mesoderm is rather obscured by out-of-focus labelling in the foregut visceral mesoderm. (G) Stage 16; labelling is predominantly in the brain lobes and in the thoracic neuromeres of the ventral nervous system. The arrows indicate labelling associated with head and gnathal sense organs. DNA probes were used for the in situ hybridizations in A, E and G; RNA probes were used for the others.




Fig. 5. Structure of transformation constructs containing *connectin* regulatory regions. For each of the four constructs (4 kb, 4 kb-35, and 1 kb-35; see text), the *connectin* DNA is shown boxed whilst the minimal hsp70 promoter-*lacZ* sequences of the vector are shown (see Materials and methods) as a line. The arrow indicates the start and direction of *lacZ* transcription from these constructs. S, P and R indicate the location of *SalI*, *PstI* and *EcoRI* restriction sites, respectively (see Fig. 1). The position of clone 35 (in the 4 kb and 1 kb constructs) is indicated by a filled box, and its absence in the 4 kb-35 and 1 kb-35 is indicated by a line. The 'no. of lines' indicates the total number of independent transformant lines analysed for *lacZ* expression (denominator) and the number of these showing a consistent pattern of expression (numerator).

likely to repress the 4 kb construct in the posterior abdominal segments. In all the mutants described above, the posterior domain of expression in A9 appeared normal.

To investigate potential activating homeotic loci, we examined the expression of the 4 kb construct in embryos homozygous for an *Antp* null mutation. In these mutant embryos, expression is strongly reduced in the thoracic segments, T2 and T3, but unaffected in T1, the gnathal region, and in A9 (Fig. 6F). This indicates that *Antp* is required for the high levels of expression found in T2 and T3 of the wild-type embryo. Consistent with this, the segments T2 and T3 are known to be the principal segments affected by *Antp* loss-of-function mutations (Wakimoto and Kaufman, 1981) and correspond to the domain of *Antp* expression in the mesoderm (Boulet and Scott, 1988; Martinez Arias, 1986).

Transcription driven by a 1 kb region encompassing clone 35

In order to examine the regulatory role of sequences more immediately surrounding clone 35, we next tested a 1 kb *PstI-SalI* fragment in which the 110 bp clone 35 is centrally located (Fig. 5). Like the 4 kb fragment, this 1 kb fragment was placed upstream of the hsp70 minimal promoter in both orientations. Consistent staining, related to the *connectin* gene expression pattern, was obtained only when the 1 kb element was in the same orientation as the functioning 4 kb element. Expression is confined to a subset of the 4 kb pattern with most labelling in the gnathal region (Fig. 7A). Compared to the 4 kb pattern, the staining observed in T1-T3 and in A9 appears much reduced in intensity relative to the gnathal expression (and is com-

pletely absent in several transformant lines). Thus, the 1 kb element lacks some of the sequences required for the high levels of expression in T1-T3 and A9 that are characteristic of the 4 kb construct and presumably these sequences reside in the 3 kb remainder fragment.

We were surprised to observe an additional pattern of expression in the visceral mesoderm of the midgut with the 1 kb construct that is not observed with the 4 kb construct. As described above, endogenous *connectin* RNA is expressed in a segmentally repeated subset of the visceral mesoderm. However, whilst the expression of *connectin* RNA in the presumptive visceral mesoderm of the midgut is seen transiently from stage 11-13, -galactosidase activity from the 1 kb construct does not appear until stage 15. The visceral mesoderm expression from the 1 kb construct is restricted to the developing second midgut constriction (Fig. 7B,C) and corresponds to a subset of parasegment 7. This is particularly interesting as *Ubx* is expressed in PS7 in the visceral mesoderm and *Ubx* function is required for the formation of the second midgut constriction (Bienz et al., 1988; Bienz and Tremml, 1988). The visceral mesoderm expression of the 1 kb construct is also dependent on *Ubx* as it is abolished in *Ubx* mutant embryos, whilst the gnathal pattern remains unaffected (Fig. 7D).

Deletion of clone 35 sequences from 4 kb and 1 kb regulatory regions

To test directly the requirement for the clone 35 sequence in the regulation conferred by the 4 kb or the 1 kb elements, constructs containing a precise deletion of this 110 bp region but similar in every other respect were made (called 4 kb-35 and 1 kb-35 constructs).

With the 4 kb-35 construct, the gnathal expression is little affected, but the expression in T1-T3 is weak and variable (Fig. 8A) indicating that the 110 bp region is required for the normal high levels of expression in T1-T3. As described above, comparison between the expression from the 4 kb and 1 kb constructs led to the conclusion that sequences outside the 1 kb region are also required for high levels of thoracic expression. Taken together, these observations imply that multiple sites of control are involved.

The 1 kb-35 construct gave an expression pattern identical to the 1 kb construct within the gnathal segments. However, whereas the 1 kb construct gave expression in the visceral mesoderm, staining in this tissue was completely absent in the 1 kb-35 construct (Fig. 8B). As the visceral mesoderm expression from the intact 1 kb element is also absent in a *Ubx*⁻ genetic background, we conclude that clone 35 sequences are required in vivo to mediate *Ubx*-dependent activation in PS7 of the visceral mesoderm.

Discussion

In this paper, we have shown that the immunopurified clone 35, which was isolated as a potential in vivo *Ubx* protein binding site, appears to be an important component of a regulatory element under homeotic gene control. This regulatory element is part of the control sequences regulating the expression of *connectin*, a gene encoding a cell-surface member of the LRR family.




Fig. 6. Expression from the 4 kb construct in wild-type and homeotic mutant embryos. (A) 4 kb construct, line 4.20 in wild-type background; stage 13; ventrolateral view. The most prominent labelling is in the thoracic mesoderm. Labelling is also seen in the gnathal segments and in the abdominal mesoderm. The pattern of expression in the mesoderm closely mimics the endogenous *connectin* expression (B) 4 kb construct, line 4.40 in wild-type background; stage 13; ventral view. In this line, the labelling is more tightly restricted to gnathal and thoracic segments with little or no labelling of abdominal segments except A9. Note absence of expression in the central nervous system. C-F show comparable ventrolateral views of the 4 kb construct, line 4.40, in the various genetic backgrounds. (C) 4 kb construct, line 4.40 in wild-type background; stage 13. (D) *Ubx* mutant (*Df bxd*¹⁰⁰) stage 13 embryo showing staining anteriorly in gnathal and thoracic segments as in wild type, but with additional labelling in A1 and A2. (E) *Ubx abd-A* mutant, stage 13. Labelling extends strongly posteriorly to A3, and there is some staining in abdominal segments between A3 and A9. (F) *Antp* mutant, stage 13. Labelling is restricted to gnathal segments, T1 and A9.

Connectin

The presence of multiple tandem copies of the LRR motif in the connectin molecule does not immediately assign the molecule to a specific functional class. LRRs were first identified in the leucine-rich alpha₂-glycoprotein (Takahashi et al., 1985) and have since been found in a diverse set of intracellular, membrane and extracellular proteins including yeast adenylate cyclase (Kataoka et al., 1985), the von Willebrand factor receptor (Titani et al., 1987) and extracellular matrix proteoglycans (Fisher et al., 1989). There is evidence that the LRRs are involved in protein-protein interactions. LRR peptides appear to adopt an amphipathic β -sheet presenting polar and apolar potential interaction surfaces (Gay et al., 1991; Krantz et al., 1991).

Deletion analysis of the LRR region in adenylate cyclase implicates LRRs in complex formation with RAS proteins (Field et al., 1990). Also, several cell surface LRR molecules, including the *Drosophila* molecules chaoptin and *Toll* protein, share the common property of mediating cell-cell adhesion (Krantz and Zipursky, 1990; Keith and Gay, 1990). Chaoptin shows additional similarity to connectin as it is also linked to the membrane via a GPI-anchor (Krantz and Zipursky, 1990). Mutants in chaoptin disrupt the organization of photoreceptor membranes consistent with a role for chaoptin as a cell adhesion molecule required for normal photoreceptor morphogenesis. Expression of connectin in tissue culture cells reveals that it can also function as a homophilic cell-cell adhesion molecule (Nose et al., 1992;




Fig. 7. Expression from the 1 kb construct. (A) Wild-type stage 12 embryo, showing gnathal expression (G) in the labial segment, together with weak expression in the thorax, in T1 and T2. (B) Wild-type stage 15 embryo. Labelling is seen anteriorly in the gnathal segments and the arrow indicates the expression domain in the visceral mesoderm. (C) Wild-type stage 16 embryo. Expression occurs anteriorly in gnathal segments and posteriorly in A9. The arrow indicates labelling in the visceral mesoderm at the second midgut constriction. (D) *Ubx* mutant (Df *bx^{d100}*) stage 17 embryo. The gnathal labelling is present, but there is no labelling in the visceral mesoderm. The arrow indicates the approximate position corresponding to the second midgut constriction in the wild type.




Fig. 8. Expression from 1 kb and 4 kb constructs lacking the clone 35 sequences. (A) 4 kb-35 construct viewed from above. Stage 14 embryo showing strong gnathal labelling but weak and variable expression in the thoracic segments. (B) 1 kb-35 construct, stage 17 embryo. The gnathal expression is present anteriorly, but there is no staining in the visceral mesoderm.

L. Meadows, A. P. G. and R. A. H. W., unpublished results). This supports the connection between cell surface LRR-containing molecules and adhesion, and further implicates the LRR sequences themselves in protein-protein interactions.

Regulation of the connectin gene by multiple homeotic loci

Previously, we have demonstrated that *connectin* RNA expression is regulated in the central nervous system by *Ubx* and the abdominal homeotic genes (Gould et al., 1990). Here we have examined the control of a 4 kb *connectin* regulatory element that directs expression predominantly in the somatic mesoderm and not in the ventral nervous system. We have shown that this 4 kb region, located at least 45 kb 3' of the transcription start site, is sufficient to drive a subset of the *connectin* expression pattern that is under the control of multiple homeotic genes, including *Ubx*. Genetic experiments have shown that *Antp* is the homeotic gene required to activate expression of the 4 kb construct in T2 and T3. However, in *Antp*⁻ mutants,

expression remains in gnathal segments, in T1 and in A9. The expression in T1 is likely to be dependent on *Sex-combs reduced* (*Scr*), whilst *Deformed*, *labial* and *proboscipedia* may be required for the more anterior activation (Mahaffey and Kaufman, 1988).

The 4 kb construct is strongly expressed in the thoracic segments but shows very little expression in abdominal segments. We have shown that *Ubx* and *abd-A* are required for this repression in the abdomen. However, the possibility exists that this regulation could be mediated through the influence of *Ubx* and *abd-A* on the expression of *Antp*. Both *Ubx* and *abd-A* normally repress *Antp* expression in the abdominal segments (Hafen et al., 1984). Thus, in *Ubx* or *Ubx abd-A* mutants, *Antp* expression in the abdomen is derepressed and these higher levels of *Antp* protein might directly enhance expression from the 4 kb construct. However, studies of transformed cuticular phenotypes induced by ectopic *Antp* expression indicate that *Antp* is not able to influence cuticle phenotype in regions expressing *Ubx* or the abdominal homeotic genes (Gibson and Gehring, 1988). This may suggest that *Ubx* and *abd-A* exert a dominant direct repression of target genes, blocking their activation by *Antp*. This phenomenon has been called 'phenotypic suppression' (Gonzalez-Reyes and Morata, 1990) and it is likely to be important for our understanding of homeotic gene function. The analysis of the molecular basis of phenotypic suppression has been impeded by the lack of identified target genes. Now, with the identification of specific target gene regulatory elements, such as the *connectin* 4 kb element, we can hope to clarify this phenomenon.

Deletion of either the clone 35 sequence or of the 3 kb *PstI-EcoRI* fragment leads to a similar reduction in the intensity of staining in T1-T3. Therefore, it is probable that there are at least two separable elements required for high levels of expression in this region. Their requirement in T2 and T3 indicates that elements within the 3 kb fragment and also within the 110 bp clone 35 both respond to *Antp* regulation. Another homeotic gene, most probably *Scr* (Mahaffey and Kaufman, 1988), may activate expression from these elements in T1.

With the 1 kb construct, we observed expression in PS7 of the visceral mesoderm, although this is not seen with the 4 kb construct. There may be two possible explanations for this; either repressing elements are present in the deleted 3 kb region or the visceral mesoderm expression element in the 1 kb construct may not function correctly when placed too far upstream of the hsp 70 minimal promoter. The *connectin* RNA is expressed in repeating subsets of the developing visceral mesoderm including PS7, whereas the 1 kb construct appears to give only the PS7-specific subset of this pattern. However, we first detect expression from this construct only after *connectin* RNA levels in this region have declined (compare Figs 4B and 7B). Hence, in this particular instance, the 1 kb construct does not faithfully reproduce the timing of onset of *connectin* expression. Nevertheless, the PS7 domain does allow a specific connection to be made between the 110 bp immunopurified fragment and *Ubx* control as expression is dependent on the presence of both the clone 35 sequence and *Ubx* function. Given that clone 35 was originally immunopurified using an antibody directed against *Ubx* proteins and given that it interacts

directly with *Ubx* protein in vitro (Gould et al., 1990), it seems likely that this 110 bp region mediates direct regulation by *Ubx*.

For the 1 kb construct, deletion of the 110 bp fragment has the same effect on expression in the visceral mesoderm as the removal of *Ubx* gene function. This is what would be expected if the 110 bp sequence identifies a specific *Ubx*-dependent regulatory site. However, for the 4 kb construct, a *Ubx* mutation results in increased expression in the abdominal somatic mesoderm whereas deletion of the 110 bp results in reduced thoracic expression. Our interpretation of these different effects is that the deletion of the 110 bp fragment removes, not only sites required for *Ubx* and *abd-A* repression, but also the sites required for activation by *Antp* and *Scr*.

The DNA sequence of the 110 bp region contains six copies of TAAT, the core homeodomain binding sequence (Egger et al., 1991; Kissinger et al., 1990). Using mutagenised versions of the 1 kb construct, it should now be possible to assay each of the TAAT cores for a direct and functional interaction with *Ubx*, *abd-A*, *Antp* and *Scr*. As the 110 bp sequence appears to be recognised by multiple homeotic loci, it will be interesting to determine whether the different homeotic gene products act on separate binding sites or whether they compete for the same sites.

In summary, the immunopurification procedure has successfully led us to a regulatory element under homeotic gene control. We are currently investigating whether other binding sites from the immunopurified DNA library are also associated with elements regulated by homeotic genes. In principle, this method should be generally applicable to searches for in vivo targets of characterised transcription factors or DNA-binding proteins.

Properties of genes regulated by the homeotic loci

Garcia-Bellido proposed that the homeotic genes would exert their effects on morphogenesis by controlling the activities of a group of cytodifferentiation genes. These cytodifferentiation genes would specify a number of cellular characteristics including, mitotic rate, preferential mitotic orientations, cell affinities and final cellular differentiation (Garcia-Bellido, 1975). The isolation of *connectin* as a homeotic target gene fits well with this scheme. *Connectin* is a cell surface molecule capable of mediating cell-cell adhesion and the analysis of its protein distribution in detail reveals that it is expressed in a subset of motoneurons and also in their specific target muscle cells both prior to and after the formation of neuromuscular junctions (Nose et al., 1992). Thus it is tempting to speculate that *connectin* is a cell-adhesion or recognition molecule involved in the development of specific neuromuscular connections. Interestingly, recent studies in *C. elegans* have implicated an *Antp* homologue *mab-5*, in the control of cell surface molecules that affect region-specific migratory cell behaviours in the nervous system (Salser and Kenyon, 1992). It has also been suggested that the neural cell adhesion molecule (N-CAM) is likely to be regulated by mouse Hox-gene products (Jones et al., 1992).

Our observation that *connectin* is likely to be regulated directly by homeotic genes is interesting with regard to the overall command chain of developmental gene controls.

The complex pattern of homeotic gene expression is largely the result of a hierarchy of regulation by transcription factors extending from *bicoid*, through gap genes, pair-rule genes and segment polarity genes (reviewed Ingham, 1988). It had always been possible that several more layers of transcriptional control might necessarily lie between the homeotic selector genes and their cytodifferentiation targets. Here, with *connectin*, this appears not to be the case and suggests that at least some genes involved in the common processes of morphogenesis are directly under the transcriptional control of homeotic genes.

We thank Nick Brown and Sarah Bray for guidance with cDNA library screening, Shashidhara L. S. for the RNA-probe *connectin* in situ hybridizations and Peter Lawrence for help with the transformations. We are grateful to Sarah Bray, Lisa Meadows and David Strutt for their comments on the manuscript. This work was supported by the MRC and the Wellcome Trust.

References

- Akam, M. (1987). The molecular basis for metameric pattern in the *Drosophila* embryo. *Development* **101**, 1-22.
- Akam, M. and Martinez-Arias, A. (1985). The distribution of *Ultrabithorax* transcripts in *Drosophila* embryos. *EMBO J.* **4**, 1689-1700.
- Andrew, D. J. and Scott, M. P. (1992). Downstream of the homeotic genes. *New Biologist* **4**, 5-15.
- Ashburner, M. (1989a). *Drosophila. A Laboratory Handbook*. New York: Cold Spring Harbor Laboratory Press.
- Ashburner, M. (1989b). *Drosophila. A Laboratory Manual*. New York: Cold Spring Harbor Laboratory Press.
- Beachy, P. A., Krasnow, M. A., Gavis, E. R. and Hogness, D. M. (1988). An *Ultrabithorax* protein binds near its own and *Antennapedia* P1 promoters. *Cell* **55**, 1069-1081.
- Bienz, M., Saari, G., Tremml, G., Muller, J., Züst, B. and Lawrence, P. A. (1988). Differential regulation of *Ultrabithorax* in two germ layers of *Drosophila*. *Cell* **53**, 567-576.
- Bienz, M. and Tremml, G. (1988). Domain of *Ultrabithorax* expression in *Drosophila* visceral mesoderm from autoregulation and exclusion. *Nature* **333**, 576-578.
- Boulet, A. and Scott, M. (1988). Control elements of the P2 promoter of the *Antennapedia* gene. *Genes Dev.* **2**, 1600-1614.
- Brown, N. H. and Kafatos, F. C. (1988). Functional cDNA libraries from *Drosophila* embryos. *J. Mol. Biol.* **203**, 425-437.
- Campos-Ortega, J. A. and Hartenstein, V. (1985) *The Embryonic Development of Drosophila melanogaster*. New York: Springer-Verlag.
- Cavener, D. R. (1987). Comparison of the consensus sequence flanking translational start sites in *Drosophila* and vertebrates. *Nucl. Acids Res* **1987**, 1353-1361.
- Devereux, J., Haerberli, P. and Smithies, O. (1984). A comprehensive set of sequence analysis programs for the VAX. *Nucl. Acids. Res.* **12**, 387-395.
- Ekker, S. C., Young, K. E., Von Kessler, D. P. and Beachy, P. A. (1991). Optimal DNA sequence recognition by the *Ultrabithorax* homeodomain of *Drosophila*. *EMBO J.* **10**, 1179-1186.
- Ferguson, M. A. F. and Williams, A. F. (1988). Cell-surface anchoring of proteins via glycosyl-phosphatidylinositol structures. *Ann. Rev. Biochem.* **57**, 285-320.
- Field, J., Xu, H.-P., Michaeli, T., Ballester, R., Sass, P. and Wigler, M. (1990). Mutations of the adenyl cyclase gene that block RAS function in *Saccharomyces cerevisiae*. *Science* **247**, 464-467.
- Fisher, L. W., Termine, J. D. and Young, M. F. (1989). Deduced protein sequence of bone small proteoglycan I (Biglycan) shows homology with proteoglycan II (Decorin) and several nonconnective tissue proteins in a variety of species. *J. Biol. Chem* **264**, 4571-4576.
- Garcia-Bellido, A. G. (1975). Genetic control of wing disc development in *Drosophila*. In *Cell Patterning, Ciba Foundation Symposium* **29** (ed. S. Brenner), pp. 161-178. New York: Associated Scientific Publishers.
- Gay, N. J., Packman, L. C., Weldon, M. A. and Barna, J. C. J. (1991). A leucine rich repeat peptide derived from the *Drosophila* Toll receptor forms extended filaments with a β -sheet structure. *FEBS Lett.* **291**, 87-91.
- Gibson, G. and Gehring, W. J. (1988). Head and thoracic transformations caused by ectopic expression of *Antennapedia* during *Drosophila* development. *Development* **102**, 657-675.
- Gonzalez-Reyes, A. and Morata, G. (1990). The developmental effect of overexpressing a Ubx product in *Drosophila* embryos is dependent on its interactions with other homeotic products. *Cell* **61**, 515-522.
- Gould, A. P., Brookman, J. J., Strutt, D. I. and White, R. A. H. (1990). Targets of homeotic gene control in *Drosophila*. *Nature* **348**, 308-312.
- Hafen, E., Levine, M. and Gehring, W. J. (1984). Regulation of *Antennapedia* transcript distribution by the bithorax complex of *Drosophila*. *Nature* **307**, 287-289.
- Hashimoto, C., Hudson, K. L. and Anderson, K. V. (1988). The Toll gene of *Drosophila*, required for dorso-ventral polarity, appears to encode a transmembrane protein. *Cell* **52**, 269-279.
- Hickey, M. J., Williams, S. A. and Roth, G. J. (1989). Human platelet glycoprotein IX: An adhesive prototype of leucine-rich glycoproteins with flank-center-flank structures. *Proc. Natl. Acad. Sci. USA* **86**, 6773-6777.
- Hiromi, Y. and Gehring, W. J. (1987). Regulation and function of the *Drosophila* segmentation gene *fushi tarazu*. *Cell* **50**, 963-974.
- Hooper, J. E. (1986). Homeotic gene function in the muscles of *Drosophila* larvae. *EMBO J.* **5**, 2321-2329.
- Immergluck, K., Lawrence, P. A. and Bienz, M. (1990). Induction across germ layers in *Drosophila* mediated by a genetic cascade. *Cell* **62**, 261-268.
- Ingham, P. W. (1988). The molecular genetics of embryonic pattern formation in *Drosophila*. *Nature* **335**, 25-34.
- Jones, F. S., Prediger, E. A., Bittner, D. A., De Robertis, E. M. and Edelman, G. M. (1992). Cell adhesion molecules as targets for Hox genes: Neural cell adhesion molecule promoter activity is modulated by cotransfection with Hox-2.5 and -2.4. *Proc. Natl. Acad. Sci. USA* **89**, 2086-2090.
- Kataoka, T., Broek, D. and Wigler, M. (1985). DNA sequence and characterization of the *S. cerevisiae* gene encoding adenylate cyclase. *Cell* **43**, 493-505.
- Keith, F. J. and Gay, N. J. (1990). The *Drosophila* membrane receptor Toll can function to promote cellular adhesion. *EMBO J.* **9**, 4299-4306.
- Kennison, J. A. and Tamkun, J. W. (1988). Dosage-dependent modifiers of Polycomb and *Antennapedia* mutations in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **85**, 8136-8140.
- Kissinger, C. R., Liu, B., Martin-Blanco, E., Kornberg, T. and Pabo, C. O. (1990). Crystal structure of an engrailed homeodomain-DNA complex at 2.8 Å resolution: a framework for understanding homeodomain-DNA interactions. *Cell* **63**, 579-590.
- Krantz, D. D., Zidovetski, R., Kagan, B. L. and Zipursky, S. L. (1991). Amphipathic structure of a leucine-rich repeat peptide. *J. Biol. Chem.* **266**, 16801-16807.
- Krantz, D. E. and Zipursky, S. L. (1990). *Drosophila* chaoptin, a member of the leucine-rich repeat family, is a photoreceptor cell-specific adhesion molecule. *EMBO J.* **9**, 1969-1977.
- Krasnow, M. A., Saffman, E. E., Kornfeld, K. and Hogness, D. S. (1989). Transcriptional activation and repression by *Ultrabithorax* proteins in cultured *Drosophila* cells. *Cell* **57**, 1031-1043.
- Kyte, J. and Doolittle, R. (1982). A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* **157**, 105-132.
- Lewis, E. B. (1963). Genes and developmental pathways. *Am. Zool.* **3**, 33-56.
- Lewis, E. B. (1978). A gene complex controlling segmentation in *Drosophila*. *Nature* **276**, 565-570.
- Lindsley, D. and Zimm, G. (1985) The genome of *Drosophila melanogaster*. Part 1: Genes A-K. *Drosophila Inform. Service* **62**.
- Lindsley, D. and Zimm, G. (1990) The genome of *Drosophila melanogaster*. Part 4: Genes L-Z, balancers, transposable elements. *Drosophila Inform. Service* **68**.
- Lopez, J., Chung, D., Fugikawa, K., Hagen, F. and Papayannopoulou, T. (1987). Cloning of the α -chain of glycoprotein 1b: a transmembrane protein with homology to leucine-rich α -2-glycoprotein. *Proc. Natl. Acad. Sci. USA* **84**, 5615-5619.
- Mahaffey, J. W. and Kaufman, T. C. (1988). The homeotic genes of the *Antennapedia* complex and the Bithorax complex of *Drosophila*. In *Developmental Genetics of Higher Organisms: A Primer in*

- Developmental Biology* (ed. G. M. Malacinski), pp 329-360. New York: Macmillan.
- Martinez-Arias, A.** (1986). The Antennapedia gene is required and expressed in parasegments 4 and 5 of the *Drosophila* embryo. *EMBO J.* **5**, 135-141.
- Muller, M., Affolter, M., Leupin, W., Otting, G., Wuthrich, K. and Gehring, W.** (1988). Isolation and sequence specific DNA binding of the *Antennapedia* homeodomain. *EMBO J.* **7**, 4299-4304.
- Nose, A., Mahajan, V. B., and Goodman, C. S.** (1992) Connectin: A Homophilic cell adhesion molecule expressed on a subset of muscles and the motoneurons that innervate them in *Drosophila*. *Cell* **70**, 553-567.
- Panganiban, G. E. F., Reuter, R., Scott, M. P. and Hoffman, F. M.** (1990). A *Drosophila* growth factor homolog, decapentaplegic, regulates homeotic gene expression within and across germ layers during midgut morphogenesis. *Development* **110**, 1041-1050.
- Reinke, R., Krantz, D. E., Yen, D. and Zipursky, S. L.** (1988). Chaoptin, a cell surface glycoprotein required for *Drosophila* cell morphogenesis, contains a repeat motif found in yeast and human. *Cell* **52**, 291-301.
- Reuter, R., Panganiban, G. E. F., Hoffman, F. M. and Scott, M. P.** (1990). Homeotic genes regulate the spatial expression of putative growth factors in the visceral mesoderm of *Drosophila* embryos. *Development* **110**, 1031-1040.
- Rothberg, J. M., Jacobs, J. R., Goodman, C. S. and Artavanis-Tsakonas, S.** (1990). slit: an extracellular protein necessary for the development of midline glia and commissural axon pathways contains both EGF and LRR domains. *Genes Dev.* **4**, 2169-2187.
- Salser, S. J. and Kenyon, C.** (1992). Activation of a *C. elegans* Antennapedia homologue in migrating cells controls their direction of migration. *Nature* **355**, 255-258.
- Sambrook, J., Fritsch, E. F. and Maniatis, T.** (1989). *Molecular Cloning - A Laboratory Manual. Second edition.* New York: Cold Spring Harbor Laboratory Press.
- Sanchez-Herrero, E., Vernos, I., Marco, R. and Morata, G.** (1985). Genetic organization of *Drosophila* bithorax complex. *Nature* **313**, 108-113.
- Sanger, F., Nicklen, S. and Coulson, A. R.** (1977). DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467.
- Schneider, D. S., Hudson, K. L., Lin, T.-Y. and Anderson, K. V.** (1991). Dominant and recessive mutations define functional domains of Toll, a transmembrane protein required for dorso-ventral polarity in the *Drosophila* embryo. *Genes Dev.* **5**, 797-807.
- Schneider, R. and Schweiger, M.** (1991). A novel modular mosaic of cell adhesion motifs in the extracellular domains of the neurogenic trk and trkB tyrosine kinase receptors. *Oncogene* **6**, 1807-1811.
- Spradling, A. C.** (1986). P-element mediated transformation. In *Drosophila; A Practical Approach* (ed. D. B. Roberts), pp 175-197. Washington: IRL.
- St. Johnston, R. D. and Gelbart, W. M.** (1987). Decapentaplegic transcripts are localized along the dorso-ventral axis of the *Drosophila* embryo. *EMBO J.* **6**, 2785-2791.
- Staden, R.** (1982). An interactive graphics program for comparing and aligning nucleic acid and amino acid sequences. *Nucl. Acids Res.* **10**, 2951-2961.
- Struhl, G. and White, R. A. H.** (1985). Regulation of the *Ultrabithorax* gene of *Drosophila* by other bithorax complex genes. *Cell* **43**, 507-519.
- Takahashi, N., Takahashi, Y. and Putnam, F. W.** (1985). Periodicity of leucine and tandem repetition of a 24-amino acid segment in the primary structure of leucine-rich α_2 -glycoprotein of human serum. *Proc. Natl. Acad. Sci. USA* **82**, 1906-1910.
- Tautz, D. and Pfeifle, C.** (1989). A non-radioactive method for the localisation of specific RNAs in *Drosophila* embryos reveals translational control of the segmentation gene *hunchback*. *Chromosoma* **98**, 81-85.
- Thali, M., Muller, M., Delorenzi, M., Matthias, P. and Bienz, M.** (1988). *Drosophila* homeotic genes encode transcriptional activators similar to mammalian OTF-2. *Nature* **336**, 598-601.
- Titani, K., Takio, K., Handa, M. and Ruggeri, Z. M.** (1987). Amino-acid sequence of von-Willebrand factor-binding domain of platelet membrane glycoprotein 1b. *Proc. Natl. Acad. Sci. USA* **84**, 5610-5614.
- von Heijne, G.** (1986). A new method for predicting signal sequence cleavage sites. *Nucl. Acids Res.* **14**, 4683-4690.
- Wagner-Bernholz, J. T., Wilson, C., Gibson, G., Schuh, R. and Gehring, W. J.** (1991). Identification of target genes of the homeotic gene Antennapedia by enhancer detection. *Genes. Dev.* **5**, 2467-2480.
- Wakimoto, B. T. and Kaufman, T. C.** (1981). Analysis of larval segmentation in lethal genotypes associated with the Antennapedia gene complex in *Drosophila melanogaster*. *Dev. Biol.* **81**, 51-64.
- White, R. A. H. and Wilcox, M.** (1985). The distribution of Ultrabithorax proteins in *Drosophila*. *EMBO J.* **4**, 2035-2043.
- Winslow, G. M., Hayashi, S., Krasnow, M., Hogness, D. S. and Scott, M. P.** (1989). Transcriptional activation by the *Antennapedia* and *fushi tarazu* proteins in cultured *Drosophila* cells. *Cell* **57**, 1017-1030.

(Accepted 21 September 1992)