

Table S1. Phenotypes in mice lacking TFs that are selectively expressed in the intestine and closely related endoderm-derived organs

TF	Gene ID	References for phenotype in knockout mice*
Nr1i3	12355	Kodama et al. (2004); Wei et al. (2000)
Cdx2	12591	Beck et al. (1999); Chawengsaksophak et al. (1997)
Foxa2	15376	Hallonet et al. (2002); Lantz et al. (2004)
HNF4 α	15378	Gupta et al. (2005)
Mist1	17341	Johnson et al. (2004)
Nfe2l2	18024	Ramos-Gomez et al. (2001)
Nr1i2	18171	Xie et al. (2000)
Pdx1	18609	Ahlgren et al. (1996); Offield et al. (1996)
Nr1h4	20186	Sinal et al. (2000)
HNF1 β	21410	Coffinier et al. (2002)
VitDR	22337	McCarthy et al. (2005)
Lisch7	54135	Mesli et al. (2004)
Mlxip1	58805	Iizuka et al. (2004)

*Some knockout strains demonstrate early embryonic lethality and the phenotypes listed here were observed after generation of conditional alleles.

Additional references

- Ahlgren, U., Jonsson, J. and Edlund, H.** (1996). The morphogenesis of the pancreatic mesenchyme is uncoupled from that of the pancreatic epithelium in IPF1/PDX1-deficient mice. *Development* **122**, 1409-1416.
- Coffinier, C., Gresh, L., Fiette, L., Tronche, F., Schutz, G., Babinet, C., Pontoglio, M., Yaniv, M. and Barra, J.** (2002). Bile system morphogenesis defects and liver dysfunction upon targeted deletion of HNF1beta. *Development* **129**, 1829-1838.
- Gupta, R. K., Vatamaniuk, M. Z., Lee, C. S., Flaschen, R. C., Fulmer, J. T., Matschinsky, F. M., Duncan, S. A. and Kaestner, K. H.** (2005). The MODY1 gene HNF-4alpha regulates selected genes involved in insulin secretion. *J. Clin. Invest.* **115**, 1006-1015.
- Hallonet, M., Kaestner, K. H., Martin-Parras, L., Sasaki, H., Betz, U. A. and Ang, S. L.** (2002). Maintenance of the specification of the anterior definitive endoderm and forebrain depends on the axial mesendoderm: a study using HNF3beta/Foxa2 conditional mutants. *Dev. Biol.* **243**, 20-33.
- Iizuka, K., Bruick, R. K., Liang, G., Horton, J. D. and Uyeda, K.** (2004). Deficiency of carbohydrate response element-binding protein (ChREBP) reduces lipogenesis as well as glycolysis. *Proc. Natl. Acad. Sci. USA* **101**, 7281-7286.
- Johnson, C. L., Kowalik, A. S., Rajakumar, N. and Pin, C. L.** (2004). Mist1 is necessary for the establishment of granule organization in serous exocrine cells of the gastrointestinal tract. *Mech. Dev.* **121**, 261-272.
- Kodama, S., Koike, C., Negishi, M. and Yamamoto, Y.** (2004). Nuclear receptors CAR and PXR cross talk with FOXO1 to regulate genes that encode drug-metabolizing and gluconeogenic enzymes. *Mol. Cell Biol.* **24**, 7931-7940.
- Lantz, K. A., Vatamaniuk, M. Z., Brestelli, J. E., Friedman, J. R., Matschinsky, F. M. and Kaestner, K. H.** (2004). Foxa2 regulates multiple pathways of insulin secretion. *J. Clin. Invest.* **114**, 512-520.
- McCarthy, T. C., Li, X. and Sinal, C. J.** (2005). Vitamin D receptor-dependent regulation of colon multidrug resistance-associated protein 3 gene expression by bile acids. *J. Biol. Chem.* **280**, 23232-23242.
- Mesli, S., Javorschi, S., Berard, A. M., Landry, M., Priddle, H., Kivlichan, D., Smith, A. J., Yen, F. T., Bihain, B. E. and Darmon, M.** (2004). Distribution of the lipolysis stimulated receptor in adult and embryonic murine tissues and lethality of LSR-/- embryos at 12.5 to 14.5 days of gestation. *Eur. J. Biochem.* **271**, 3103-3114.
- Ramos-Gomez, M., Kwak, M. K., Dolan, P. M., Itoh, K., Yamamoto, M., Talalay, P. and Kensler, T. W.** (2001). Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers is lost in nrf2 transcription factor-deficient mice. *Proc. Natl. Acad. Sci. USA* **98**, 3410-3415.
- Sinal, C. J., Tohkin, M., Miyata, M., Ward, J. M., Lambert, G. and Gonzalez, F. J.** (2000). Targeted disruption of the nuclear receptor FXR/BAR impairs bile acid and lipid homeostasis. *Cell* **102**, 731-744.
- Wei, P., Zhang, J., Egan-Hafley, M., Liang, S. and Moore, D. D.** (2000). The nuclear receptor CAR mediates specific xenobiotic induction of drug metabolism. *Nature* **407**, 920-923.
- Xie, W., Barwick, J. L., Downes, M., Blumberg, B., Simon, C. M., Nelson, M. C., Neuschwander-Tetri, B. A., Brunt, E. M., Guzelian, P. S. and Evans, R. M.** (2000). Humanized xenobiotic response in mice expressing nuclear receptor SXR. *Nature* **406**, 435-439.