

Development 135, 3969 (2008) doi:10.1242/dev.022061

## Differential contributions of *Mesp1* and *Mesp2* to the epithelialization and rostro-caudal patterning of somites

**Yu Takahashi<sup>1</sup>, Shuichi Hiraoka<sup>2</sup>, Satoshi Kitajima<sup>1</sup>, Tohru Inoue<sup>1</sup>, Jun Kanno<sup>1</sup> and Yumiko Saga<sup>3</sup>**

<sup>1</sup>Cellular & Molecular Toxicology Division, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagayaku, Tokyo 158-8501, Japan. <sup>2</sup>Laboratory for Developmental Genetics, RIKEN Research Center for Allergy and Immunology, 1-7-22 Suehirocho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan. <sup>3</sup>Division of Mammalian Development, National Institute of Genetics, Yata 1111, Mishima 411-8540, Japan.

There was an error published in *Development* **132**, 787-796.

Author Shuichi Hiraoka was omitted by mistake. The contribution made by Hiraoka was equal to that of the first author during the crucial phase when the main conclusion of the paper was reached.

The correct author list and affiliations are written above.

The authors apologise to readers for this mistake.