

Table S1. Primers used for quantitative RT-PCR

Gene	Forward (5'-3')	Reverse (5'-3')
<i>Bmp4</i>	AGATTGGCTCCCAAGAATCA	GCCTCCTAGCAGGACTTGG
<i>Dkk1</i>	CCGTCTGCCTCCGATCAT	GGACCAGAAGTGTCTTGCACAA
<i>Eomes</i>	CCACTGGATGAGGCAGGAGATTTC	AGTCTTGGAAGGTTCAATCAAGTCC
<i>Fgf5</i>	CGAGGAGTTTTTCAGCAACAA	CGCGGACGCATAGGTATTAT
<i>Gapdh</i>	TCAAGAAGGTGGTGAAGCAG	ACCACCCTGTTGCTGTAGCC
<i>Il2rg</i>	AATCCCCCATCAAGAATC	GTAGTCTGGCTGCAGACTCTCA
<i>Lef1</i>	AGCCTGTTTATCCCATCACG	TGAGGCTTCACGTGCATTAG
<i>Nanog</i>	CCTCCATTCTGAACCTGAGC	GCAATGGATGCTGGGATACT
<i>Nefl</i>	CCACTCTGCAAGCAAACAGA	CCACTGGATGAGGCAGGAGATTTC
<i>Nlgn3</i>	CTGTCTTTGGCTCTGGCATC	AGAGCCACTTTGGATGATGG
<i>Oct4</i>	AGCTGCTGAAGCAGAAGAGG	AGATGGTGGTCTGGCTGAAC
<i>Rest</i>	GTGCGAACTCACACAGGAGA	AAGAGGTTTAGGCCCGTTGT
<i>Rex1</i>	CCTGCACACAGAAGAAAGCA	CACTGATCCGCAAACACCT
<i>Sox2</i>	AGACCGTTTTTCGTGGTCTTG	TATCAACCTGCATGGGCATT
<i>T</i>	CAGCTGTCTGGGAGCCTGG	TGCTGCCTGTGAGTCATAAC