

Table S2. Sequences of primers used in QPCR and ChIP experiments

Primers used for QPCR on mRNA extracted from WT and <i>COUP-TFI</i> KO embryos (Fig. 4A)		
mRNA	Forward (5'-3')	Reverse (5'-3')
<i>Ngn2</i>	GACATTCCCGACACACAC	CCAGCAGCATCAGTACCTCC
<i>NeuroD1</i>	CGCTCTCGCTGTATGATTTG	ACGCAGAAGGCAAGGTGTC
<i>NeuroD2</i>	CTGGAGAGCTGCAGAACCG	TTTTTCTTTCCTTTGTCCG
<i>Tbr2</i>	AAACACGGATATCACCCAGC	GGCAAAGTGTGACAAAGGG
<i>Rnd2</i>	GTGTTTGAGAACTACACTGC	GTGTTTCTGCCGGCTAATG
Primers used to query chromatin enrichment by QPCR (Fig. 4D)		
Primers	Forward (5'-3')	Reverse (5'-3')
Bs A-A1	CGTTTGACCTTCCACCTTAG	TCCCACTTGCTTGGCCAGC
Bs B	GAGATTGGACGAATAATTGGC	CTGCCAGGTTTTGAATCTGC
Bs C	TCACTGGGTGACTCTTTTCG	GCAATGCCAGACAATGTGAG
Bs D	AGTCCTGAGCTGGCTGGGTCC	TCTCCTCACTAGGGCAACAG
mFabp7	CTAACCCAGTGGCCTGAGCA	TGGGCAGATGGCTCCAAT
mCycloA	AGGTCCTGGCATCTTGTTCCA	GAACCGTTTGTGTTGGTCCA