

Development

Contents

Volume 129 (11) 2002

Latimer, A. J., Dong, X., Markov, Y. and Appel, B. Delta-Notch signaling induces hypochord development in zebrafish	2555-2563	Imai, K. S., Satou, Y. and Satoh, N. Multiple functions of a Zic-like gene in the differentiation of notochord, central nervous system and muscle in <i>Ciona savignyi</i> embryos	2723-2732
Parker, D. S., Jemison, J. and Cadigan, K. M. Pygopus, a nuclear PHD-finger protein required for Wingless signaling in <i>Drosophila</i>	2565-2576	Auman, H. J., Nottoli, T., Lakiza, O., Winger, Q., Donaldson, S. and Williams, T. Transcription factor AP-2 γ is essential in the extra-embryonic lineages for early postimplantation development	2733-2747
Shvartsman, S. Y., Muratov, C. B. and Lauffenburger, D. A. Modeling and computational analysis of EGF receptor-mediated cell communication in <i>Drosophila</i> oogenesis	2577-2589	Lawrence, P. A., Casal, J. and Struhl, G. Towards a model of the organisation of planar polarity and pattern in the <i>Drosophila</i> abdomen	2749-2760
Sedaghat, Y., Miranda, W. F. and Sonnenfeld, M. J. The <i>jing</i> Zn-finger transcription factor is a mediator of cellular differentiation in the <i>Drosophila</i> CNS midline and trachea	2591-2606	DEVELOPMENT AND DISEASE	
Nechiporuk, A. and Keating, M. T. A proliferation gradient between proximal and <i>msxb</i> -expressing distal blastema directs zebrafish fin regeneration	2607-2617	Corsi, A. K., Brodigan, T. M., Jorgensen, E. M. and Krause, M. Characterization of a dominant negative <i>C. elegans</i> Twist mutant protein with implications for human Saethre-Chotzen syndrome	2761-2772
Katz, J. P., Perreault, N., Goldstein, B. G., Lee, C. S., Labosky, P. A., Yang, V. W. and Kaestner, K. H. The zinc-finger transcription factor Klf4 is required for terminal differentiation of goblet cells in the colon	2619-2628	Minasi, M. G., Riminiucci, M., De Angelis, L., Borello, U., Berarducci, B., Innocenzi, A., Caprioli, A., Sirabella, D., Baiocchi, M., De Maria, R., Boratto, R., Jaffredo, T., Broccoli, V., Bianco, P. and Cossu, G. The meso-angioblast: a multipotent, self-renewing cell that originates from the dorsal aorta and differentiates into most mesodermal tissues	2773-2784
Laudencia-Chingcuanco, D. and Hake, S. The <i>indeterminate floral apex1</i> gene regulates meristem determinacy and identity in the maize inflorescence	2629-2638		
Cornell, R. A. and Eisen, J. S. Delta/Notch signaling promotes formation of zebrafish neural crest by repressing neurogenin 1 function	2639-2648		
Rathjen, J., Haines, B. P., Hudson, K. M., Nesci, A., Dunn, S. and Rathjen, P. D. Directed differentiation of pluripotent cells to neural lineages: homogeneous formation and differentiation of a neurectoderm population	2649-2661		
Scanlon, M. J., Henderson, D. C. and Bernstein, B. SEMAPHORE1 functions during the regulation of ancestrally duplicated <i>knox</i> genes and polar auxin transport in maize	2663-2673		
Kahane, N., Cinnamon, Y. and Kalcheim, C. The roles of cell migration and myofiber intercalation in patterning formation of the postmitotic myotome	2675-2687		
Miura, T., Miura, C., Konda, Y. and Yamauchi, K. Spermatogenesis-preventing substance in Japanese eel	2689-2697		
Gamberi, C., Peterson, D. S., He, L. and Gottlieb, E. An anterior function for the <i>Drosophila</i> posterior determinant Pumilio	2699-2710		
Willem, M., Miosge, N., Halfter, W., Smyth, N., Jannetti, I., Burghart, E., Timpl, R. and Mayer, U. Specific ablation of the nidogen-binding site in the laminin $\gamma 1$ chain interferes with kidney and lung development	2711-2722		