

Development

Contents

Volume 130 (I) 2003

EDITORIAL

Smith, J. C.

Ch-ch-ch-changes...

1

RESEARCH ARTICLES

Przemeck, G. K. H., Heinzmann, U., Beckers, J. and Hrabé de Angelis, M.

Node and midline defects are associated with left-right development in *Delta1* mutant embryos

3-13

Kenney, A. M., Cole, M. D. and Rowitch, D. H.

Nmyc upregulation by sonic hedgehog signaling promotes proliferation in developing cerebellar granule neuron precursors

15-28

Bizzoca, A., Virgintino, D., Lorusso, L., Buttiglione, M., Yoshida, L., Polizzi, A., Tattoli, M., Cagiano, R., Rossi, F., Kozlov, S., Furley, A. and Gennarini, G.

Transgenic mice expressing F3/contactin from the TAG-1 promoter exhibit developmentally regulated changes in the differentiation of cerebellar neurons

29-43

Marcil, A., Dumontier, É., Chamberland, M., Camper, S. A. and Drouin, J.

Pitx1 and *Pitx2* are required for development of hindlimb buds

45-55

Moghal, N. and Sternberg, P. W.

A component of the transcriptional mediator complex inhibits RAS-dependent vulval fate specification in *C. elegans*

57-69

Vernon, A. E. and Philpott, A.

A single cdk inhibitor, p27^{Xic1}, functions beyond cell cycle regulation to promote muscle differentiation in *Xenopus*

71-83

Vernon, A. E., Devine, C. and Philpott, A.

The cdk inhibitor p27^{Xic1} is required for differentiation of primary neurones in *Xenopus*

85-92

Pringle, N. P., Yu, W.-P., Howell, M., Colvin, J. S., Ornitz, D. M. and Richardson, W. D.

Fgf3 expression by astrocytes and their precursors: evidence that astrocytes and oligodendrocytes originate in distinct neuroepithelial domains

93-102

Behrens, A., Haigh, J., Mechta-Grigoriou, F., Nagy, A., Yaniv, M. and Wagner, E. F.

Impaired intervertebral disc formation in the absence of *Jun*

103-109

Trieu, M., Ma, A., Eng, S. R., Fedtsova, N. and Turner, E. E.

Direct autoregulation and gene dosage compensation by POU-domain transcription factor Brn3a

111-121

Fichelton, P. and Gho, M.

The glial cell undergoes apoptosis in the microchaete lineage of *Drosophila*

123-133

Johansen, K. A., Iwaki, D. D. and Lengyel, J. A.

Localized JAK/STAT signaling is required for oriented cell rearrangement in a tubular epithelium

135-145

Hudson, C., Darras, S., Caillol, D., Yasuo, H. and Lemaire, P.

A conserved role for the MEK signalling pathway in neural tissue specification and posteriorisation in the invertebrate chordate, the ascidian *Ciona intestinalis*

147-159

Ha, C. M., Kim, G.-T., Kim, B. C., Jun, J. H., Soh, M. S., Ueno, Y., Machida, Y., Tsukaya, H. and Nam, H. G.

The *BLADE-ON-PETIOLE 1* gene controls leaf pattern formation through the modulation of meristematic activity in *Arabidopsis*

161-172

Deng, W.-M., Schneider, M., Frock, R., Castillejo-Lopez, C., Gaman, E. A., Baumgartner, S. and Ruohola-Baker, H.

Dystroglycan is required for polarizing the epithelial cells and the oocyte in *Drosophila*

173-184

López-Ríos, J., Tessmar, K., Loosli, F., Wittbrodt, J. and Bovolenta, P.

Six3 and Six6 activity is modulated by members of the groucho family

185-195

Baena-López, L. A. and García-Bellido, A.

Genetic requirements of *vestigial* in the regulation of *Drosophila* wing development

197-208

DEVELOPMENT AND DISEASE

Délot, E. C., Bahamonde, M. E., Zhao, M. and Lyons, K. M.

BMP signaling is required for septation of the outflow tract of the mammalian heart

209-220

Wallis, D., Hamblen, M., Zhou, Y., Venken, K. J. T., Schumacher, A., Grimes, H. L., Zoghbi, H. Y., Orkin, S. H. and Bellen, H. J.

The zinc finger transcription factor *Gfi1*, implicated in lymphomagenesis, is required for inner ear hair cell differentiation and survival

221-232

Corrigendum

233



Supplemental data available on-line