



Cover: A forelimb of E12.5 homozygous *Ifi88^{hypo}* mouse embryo. In this mutant, there is a loss of both activator and repressor functions of Gli proteins, leading to the formation of extra digits. See article by Liu et al. on p. 3103.

Review

Deschamps, J. and van Nes, J.
Developmental regulation of the Hox genes during axial morphogenesis in the mouse 2931-2942

Research articles

Tschopp, O., Yang, Z.-Z., Brodbeck, D., Dummler, B. A., Hemmings-Mieszczak, M., Watanabe, T., Michaelis, T., Frahm, J. and Hemmings, B. A.
Essential role of protein kinase B γ (PKB γ /Akt3) in postnatal brain development but not in glucose homeostasis 2943-2954

Xiao, T., Roeser, T., Staub, W. and Baier, H.
A GFP-based genetic screen reveals mutations that disrupt the architecture of the zebrafish retinotectal projection 2955-2967

Collombat, P., Hecksher-Sørensen, J., Broccoli, V., Krull, J., Ponte, I., Mundiger, T., Smith, J., Gruss, P., Serup, P. and Mansouri, A.
The simultaneous loss of *Arx* and *Pax4* genes promotes a somatostatin-producing cell fate specification at the expense of the α - and β -cell lineages in the mouse endocrine pancreas 2969-2980

Schlake, T.
FGF signals specifically regulate the structure of hair shaft medulla via IGF-binding protein 5 2981-2990

Kriks, S., Lanuza, G. M., Mizuguchi, R., Nakafuku, M. and Goulding, M.
Gsh2 is required for the repression of *Ngn1* and specification of dorsal interneuron fate in the spinal cord 2991-3002

Lallemand, Y., Nicola, M.-A., Ramos, C., Bach, A., Saint Clément, C. and Robert, R.
Analysis of *Msx1*; *Msx2* double mutants reveals multiple roles for *Msx* genes in limb development 3003-3014

Close, J. L., Gumuscu, B. and Reh, T. A.
Retinal neurons regulate proliferation of postnatal progenitors and Müller glia in the rat retina via TGF β signaling 3015-3026

Yamaguchi, M., Tonou-Fujimori, N., Komori, A., Maeda, R., Nojima, Y., Li, H., Okamoto, H. and Masai, I.
Histone deacetylase 1 regulates retinal neurogenesis in zebrafish by suppressing Wnt and Notch signaling pathways 3027-3043

Bouma, G. J., Albrecht, K. H., Washburn, L. L., Recknagel, A. K., Churchill, G. A. and Eicher, E. M.
Gonadal sex reversal in mutant *Dax1* XY mice: a failure to upregulate *Sox9* in pre-Sertoli cells 3045-3054

Cobb, J. and Duboule, D.
Comparative analysis of genes downstream of the *Hoxd* cluster in developing digits and external genitalia 3055-3067

Medioni, C. and Noselli, S.
Dynamics of the basement membrane in invasive epithelial clusters in *Drosophila* 3069-3077

Dixon, S. J. and Roy, P. J.
Muscle arm development in *Caenorhabditis elegans* 3079-3092

Merabet, S., Castelli-Gair Hombria, J., Hu, N., Pradel, J. and Graba, Y.
Hox-controlled reorganisation of intrasegmental patterning cues underlies *Drosophila* posterior spiracle organogenesis 3093-3102

Liu, A., Wang, B. and Niswander, L. A.
Mouse intraflagellar transport proteins regulate both the activator and repressor functions of Gli transcription factors 3103-3111

Brendolan, A., Ferretti, E., Salsi, V., Moses, K., Quaggin, S., Blasi, F., Cleary, M. L. and Selleri, L.
A *Pbx1*-dependent genetic and transcriptional network regulates spleen ontogeny 3113-3126

Knight, R. D., Javidan, Y., Zhang, T., Nelson, S. and Schilling, T. F.
AP2-dependent signals from the ectoderm regulate craniofacial development in the zebrafish embryo 3127-3138

Research article: Development and disease

Henseleit, K. D., Nelson, S. B., Kuhlbrodt, K., Hennings, J. C., Ericson, J. and Sander, M.
NKX6 transcription factor activity is required for α - and β -cell development in the pancreas 3139-3149