



Cover: Developing olfactory axons in a zebrafish embryo double-labeled with anti-PCAM (green) and anti-calretinin (magenta) antibodies. PCAM is a cell adhesion molecule present on all olfactory axons, whereas calretinin is expressed in a small population of olfactory sensory neurons projecting mostly to glomeruli in the lateral region of the olfactory bulb. See article by Miyasaka et al. on p. 1283.

Research articles

Morris, J. Z., Hong, A., Lilly, M. A. and Lehmann, R. twin, a CCR4 homolog, regulates cyclin poly(A) tail length to permit <i>Drosophila</i> oogenesis	1165-1174	Vincent, S. D., Dunn, N. R., Sciammas, R., Shapiro-Shalef, M., Davis, M. M., Calame, K., Bikoff, E. K. and Robertson, E. J. The zinc finger transcriptional repressor Blimp1/Prdm1 is dispensable for early axis formation but is required for specification of primordial germ cells in the mouse	1315-1325
Rosset, M., Loulier, K., Feuillet, C., Alonso, S. and Carroll, P. Reelin signaling is necessary for a specific step in the migration of hindbrain efferent neurons	1175-1185	Wagner, N., Wagner, K.-D., Hammes, A., Kirschner, K. M., Vidal, V. P., Schedl, A. and Scholz, H. A splice variant of the Wilms' tumour suppressor <i>Wt1</i> is required for normal development of the olfactory system	1327-1336
Montero, J.-A., Carvalho, L., Wilsch-Bräuninger, M., Kilian, B., Mustafa, C. and Heisenberg, C.-P. Shield formation at the onset of zebrafish gastrulation	1187-1198	Weninger, W. J., Lopes Floro, K., Bennett, M. B., Withington, S. L., Preis, J. I., Martinez Barbera, J. P., Mohun, T. J. and Dunwoodie, S. L. <i>Cited2</i> is required both for heart morphogenesis and establishment of the left-right axis in mouse development	1337-1348
Hudson, C. and Yasuo, H. Patterning across the ascidian neural plate by lateral Nodal signalling sources	1199-1210	Hejnol, A. and Schnabel, R. The eutardigrade <i>Thulinia stephaniae</i> has an indeterminate development and the potential to regulate early blastomere ablations	1349-1361
Escudero, L. M., Caminero, E., Schulze, K. L., Bellen, H. J. and Modolell, J. Charlatan, a Zn-finger transcription factor, establishes a novel level of regulation of the proneural <i>achaete/scute</i> genes of <i>Drosophila</i>	1211-1222	Okubo, T., Knoepfler, P. S., Eisenman, R. N. and Hogan, B. L. M. <i>Nmyc</i> plays an essential role during lung development as a dosage-sensitive regulator of progenitor cell proliferation and differentiation	1363-1374
Steinberg, Z., Myers, C., Heim, V. M., Lathrop, C. A., Rebustini, I. T., Stewart, J. S., Larsen, M. and Hoffman, M. P. FGFR2b signaling regulates ex vivo submandibular gland epithelial cell proliferation and branching morphogenesis	1223-1234	Bae, Y.-K., Shimizu, T. and Hibi, M. Patterning of proneuronal and inter-proneuronal domains by hairy- and enhancer of split-related genes in zebrafish neuroectoderm	1375-1385
Bommert, P., Lunde, C., Nardmann, J., Vollbrecht, E., Running, M., Jackson, D., Hake, S. and Werr, W. <i>thick tassel dwarf1</i> encodes a putative maize ortholog of the <i>Arabidopsis CLAVATA1</i> leucine-rich repeat receptor-like kinase	1235-1245	Dalpé, G., Brown, L. and Culotti, J. G. Vulva morphogenesis involves attraction of plexin 1-expressing primordial vulva cells to semaphorin 1a sequentially expressed at the vulva midline	1387-1400
Essner, J. J., Amack, J. D., Nyholm, M. K., Harris, E. B. and Yost, H. J. Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut	1247-1260	Ho, K. S., Suyama, K., Fish, M. and Scott, M. P. Differential regulation of Hedgehog target gene transcription by Costal2 and Suppressor of Fused	1401-1412
Rhinn, M., Lun, K., Luz, M., Werner, M. and Brand, M. Positioning of the midbrain-hindbrain boundary organizer through global posteriorization of the neuroectoderm mediated by Wnt8 signaling	1261-1272	Yang, L., Sym, M. and Kenyon, C. The roles of two <i>C. elegans</i> HOX co-factor orthologs in cell migration and vulva development	1413-1428
James, D., Levine, A. J., Besser, D. and Hemmati-Brivanlou, A. TGF β /activin/nodal signaling is necessary for the maintenance of pluripotency in human embryonic stem cells	1273-1282	Lee, H.-H. and Frasch, M. Nuclear integration of positive Dpp signals, antagonistic Wg inputs and mesodermal competence factors during <i>Drosophila</i> visceral mesoderm induction	1429-1442
Miyasaka, N., Sato, Y., Yeo, S.-Y., Hutson, L. D., Chien, C.-B., Okamoto, H. and Yoshihara, Y. Robo2 is required for establishment of a precise glomerular map in the zebrafish olfactory system	1283-1293	Research articles: Development and disease	
Mizutani, K.-i. and Saito, T. Progenitors resume generating neurons after temporary inhibition of neurogenesis by Notch activation in the mammalian cerebral cortex	1295-1304	Andreu, P., Colnot, S., Godard, C., Gad, S., Chafey, P., Niwa-Kawakita, M., Laurent-Puig, P., Kahn, A., Robine, S., Perret, C. and Romagnolo, B. Crypt-restricted proliferation and commitment to the Paneth cell lineage following <i>Apc</i> loss in the mouse intestine	1443-1451
Smith, T. G., Sweetman, D., Patterson, M., Keyse, S. M. and Münterberg, A. Feedback interactions between MKP3 and ERK MAP kinase control scleraxis expression and the specification of rib progenitors in the developing chick somite	1305-1314	Liu, W., Sun, X., Braut, A., Mishina, Y., Behringer, R. R., Mina, M. and Martin, J. F. Distinct functions for Bmp signaling in lip and palate fusion in mice	1453-1461
		Li, B., Nair, M., Mackay, D. R., Bilanchone, V., Hu, M., Fallahi, M., Song, H., Dai, Q., Cohen, P. E. and Dai, X. <i>Ovo1</i> regulates meiotic pachytene progression during spermatogenesis by repressing Id2 expression	1463-1473
		Erratum	1475