Development 134 (15)

Development

CONTENTS



A close-up image of a lawn of bri1 Id Arabidopsis thaliana plants during early vegetative growth. This double mutant was used to establish the positive role of steroid signalling during the developmental transition towards reproductive growth. See research article by Domagalska et al. on p. 2841.

REVIEW

2709 Notch signaling in vascular development and physiology Gridley, T.


RESEARCH REPORT

Unique mechanisms of growth regulation and tumor suppression upon Apc inactivation in the pancreas
 Strom, A., Bonal, C., Ashery-Padan, R., Hashimoto, N., Campos, M. L., Trumpp, A., Noda, T., Kido, Y., Real, F. X., Thorel, F. and Herrera, P. L.

RESEARCH ARTICLES

- 2727 Chondroitin sulfate glycosaminoglycans control proliferation, radial glia cell differentiation and neurogenesis in neural stem/progenitor cells Sirko, S., von Holst, A., Wizenmann, A., Götz, M. and Faissner, A.
- 2739 The IL-4/IL-13/Stat6 signalling pathway promotes luminal mammary epithelial cell development Khaled, W. T., Read, E. K. C., Nicholson, S. E., Baxter, F. O., Brennan, A. J., Came, P. J., Sprigg, N., McKenzie, A. N. J. and Watson, C. J.
- 2751 Requirement for ERK MAP kinase in mouse preimplantation development Maekawa, M., Yamamoto, T., Kohno, M., Takeichi, M. and Nishida, E.
- 2761 Foxa1 and Foxa2 regulate multiple phases of midbrain dopaminergic neuron development in a dosage-dependent manner Ferri, A. L. M., Lin, W., Mavromatakis, Y. E., Wang, J. C., Sasaki, H., Whitsett, J. A. and Ang, S.-L.
- 2771 Zebrafish relatively relaxed mutants have a ryanodine receptor defect, show slow swimming and provide a model of multi-minicore disease Hirata, H., Watanabe, T., Hatakeyama, J., Sprague, S. M., Saint-Amant, L., Nagashima, A., Cui, W. W., Zhou, W. and Kuwada, J. Y.
- 2783 Helt determines GABAergic over glutamatergic neuronal fate by repressing Ngn genes in the developing mesencephalon
 Nakatani, T., Minaki, Y., Kumai, M. and Ono, Y.
- 2795 Bi-compartmental communication contributes to the opposite proliferative behavior of Notch1-deficient hair follicle and epidermal keratinocytes Lee, J., Basak, J. M., Demehri, S. and Kopan, R.
- 2807 The *Drosophila* HMG-domain proteins SoxNeuro and Dichaete direct trichome formation via the activation of *shavenbaby* and the restriction of Wingless pathway activity
 Overton, P. M., Chia, W. and Buescher, M.
- 2815 Conserved pattern of tangential neuronal migration during forebrain development Métin, C., Alvarez, C., Moudoux, D., Vitalis, T., Pieau, C. and Molnár, Z.
- 2829 Two distinct sources for a population of maturing axial progenitors Cambray, N. and Wilson, V.
- 2841 Attenuation of brassinosteroid signaling enhances *FLC* expression and delays flowering Domagalska, M. A., Schomburg, F. M., Amasino, R. M., Vierstra, R. D., Nagy, F. and Davis, S. J.
- 2851 A novel eIF4G homolog, Off-schedule, couples translational control to meiosis and differentiation in *Drosophila* spermatocytes
 Franklin-Dumont, T. M., Chatterjee, C., Wasserman, S. A. and DiNardo, S.

Development 134 (15)



Notch signalling plays a crucial role in numerous developmental processes and is defective in several inherited diseases and cancers. In this issue, Thomas Gridley discusses its importance in vascular development and physiology, reviewing recent work showing that it regulates artery-vein differentiation and blood vessel sprouting and branching during angiogenesis. See review on p. 2709.

2863 Translational control of meiotic cell cycle progression and spermatid differentiation in male germ cells by a novel eIF4G homolog
Baker, C. C. and Fuller, M. T.

DEVELOPMENT AND DISEASE

- 2871 Early developmental specification of the thyroid gland depends on han-expressing surrounding tissue and on FGF signals
 Wendl, T., Adzic, D., Schoenebeck, J. J., Scholpp, S., Brand, M., Yelon, D. and Rohr, K. B.
- 2881 Midbrain dopaminergic neurogenesis and behavioural recovery in a salamander lesion-induced regeneration model
 Parish, C. L., Beljajeva, A., Arenas, E. and Simon, A.