



**Cover:** Comparative upper jaw morphogenesis and growth showing relative prominence proportions (frontonasal, blue; maxillary, red) in mouse, human, alligator and chicken. Amniote facial shape and proportions converge at fusion as a common developmental 'solution' to successful integration of independent outgrowths into a functional primary palate. See Research article by Young et al. on p. 1059.

## IN THIS ISSUE

- 975** Time to update the mammalian CV; An epigenetic bag of TRX; Mammary stratification caught on camera; RAd migration in the cortex; Oct4: the plot thickens; Maternal undernutrition stimulates embryo endocytosis

## PRIMER

- 977** The Mediator complex: a master coordinator of transcription and cell lineage development  
**Yin, J. and Wang, G.**

## REVIEW

- 988** Cell competition: how to eliminate your neighbours  
**Amoyel, M. and Bach, E. A.**

## STEM CELLS AND REGENERATION

- 1001** Oct4 is required for lineage priming in the developing inner cell mass of the mouse blastocyst  
**Chia Le Bin, G., Muñoz-Descalzo, S., Kurowski, A., Leitch, H., Lou, X., Mansfield, W., Etienne-Dumeau, C., Grabole, N., Mulas, C., Niwa, H., Hadjantonakis, A.-K. and Nichols, J.**
- 1011** Transcription factor expression uniquely identifies most postembryonic neuronal lineages in the *Drosophila* thoracic central nervous system  
**Lacin, H., Zhu, Y., Wilson, B. A. and Skeath, J. B.**

- 1022** The H3K4 methyltransferase Setd1a is first required at the epiblast stage, whereas Setd1b becomes essential after gastrulation  
**Bledau, A. S., Schmidt, K., Neumann, K., Hill, U., Ciotta, G., Gupta, A., Torres, D. C., Fu, J., Kranz, A., Stewart, A. F. and Anastassiadis, K.**

- 1036** Earmuff restricts progenitor cell potential by attenuating the competence to respond to self-renewal factors  
**Janssens, D. H., Komori, H., Grbac, D., Chen, K., Koe, C. T., Wang, H. and Lee, C.-Y.**

- 1047** *Sall1* balances self-renewal and differentiation of renal progenitor cells  
**Basta, J. M., Robbins, L., Kiefer, S. M., Dorsett, D. and Rauchman, M.**

## RESEARCH REPORT

- 1059** Embryonic bauplans and the developmental origins of facial diversity and constraint  
**Young, N. M., Hu, D., Lainoff, A. J., Smith, F. J., Diaz, R., Tucker, A. S., Trainor, P. A., Schneider, R. A., Hallgrímsson, B. and Marcucio, R. S.**

## RESEARCH ARTICLES

- 1064** Microtubule-severing protein Katanin regulates neuromuscular junction development and dendritic elaboration in *Drosophila*  
**Mao, C.-X., Xiong, Y., Xiong, Z., Wang, Q., Zhang, Y. Q. and Jin, S.**

- 1075** Eph-Ephrin signaling and focal adhesion kinase regulate actomyosin-dependent apical constriction of ciliary band cells  
**Krupke, O. A. and Burke, R. D.**

- 1085** Developmental stratification of the mammary epithelium occurs through symmetry-breaking vertical divisions of apically positioned luminal cells  
**Huebner, R. J., Lechler, T. and Ewald, A. J.**

- 1095** The neural crest cell cycle is related to phases of migration in the head  
**Ridenour, D. A., McLennan, R., Teddy, J. M., Semerad, C. L., Haug, J. S. and Kulesa, P. M.**

- 1104** An anterior limit of FGF/Erk signal activity marks the earliest future somite boundary in zebrafish  
**Akiyama, R., Masuda, M., Tsuge, S., Bessho, Y. and Matsui, T.**

- 1110** Single continuous lumen formation in the zebrafish gut is mediated by *smoothened*-dependent tissue remodeling  
**Alvers, A. L., Ryan, S., Scherz, P. J., Huisken, J. and Bagnat, M.**

- 1120** Molecular identification of venous progenitors in the dorsal aorta reveals an aortic origin for the cardinal vein in mammals  
**Lindskog, H., Kim, Y. H., Jelin, E. B., Kong, Y., Guevara-Gallardo, S., Kim, T. N. and Wang, R. A.**

- 1129** Trithorax monomethylates histone H3K4 and interacts directly with CBP to promote H3K27 acetylation and antagonize Polycomb silencing  
**Tie, F., Banerjee, R., Saiakhova, A. R., Howard, B., Monteith, K. E., Scacheri, P. C., Cosgrove, M. S. and Harte, P. J.**

- 1140** Mouse early extra-embryonic lineages activate compensatory endocytosis in response to poor maternal nutrition  
**Sun, C., Velazquez, M. A., Marfy-Smith, S., Sheth, B., Cox, A., Johnston, D. A., Smyth, N. and Fleming, T. P.**

- 1151** Activated retinoid receptors are required for the migration and fate maintenance of subsets of cortical neurons  
**Choi, J., Park, S. and Sockanathan, S.**

## TECHNIQUES AND RESOURCES

- 1161** BLIND ordering of large-scale transcriptomic developmental timecourses  
**Anavy, L., Levin, M., Khair, S., Nakanishi, N., Fernandez-Valverde, S. L., Degnan, B. M. and Yanai, I.**
- 1167** A targeted gene expression system using the tryptophan repressor in zebrafish shows no silencing in subsequent generations  
**Suli, A., Guler, A. D., Raible, D. W. and Kimelman, D.**