



**Cover:** Embryonic day 8 chicken hindgut was cultured for 72 hours on a fibronectin-coated surface in the presence of supplemental glial-derived neurotrophic factor (Gdnf). Immunostaining with a neural crest cell marker (Hnk1; red) and a neuronal marker (Tuj1; green) demonstrates robust migration of the enteric neural crest cells. See Research article by Nagy et al. on p. 264.

## SPOTLIGHT

- 183 An interview with Peter Lawrence  
**Vicente, C.**

## PRIMER

- 186 Measuring forces and stresses *in situ* in living tissues  
**Sugimura, K., Lenne, P.-F. and Graner, F.**

## REVIEW

- 197 The formation and function of the cardiac conduction system  
**van Weerd, J. H. and Christoffels, V. M.**

## STEM CELLS AND REGENERATION

- 211 Genetic analysis of Runx2 function during intramembranous ossification  
**Takarada, T., Nakazato, R., Tsuchikane, A., Fujikawa, K., Iezaki, T., Yoneda, Y. and Hinoi, E.**
- 219 Genes implicated in stem cell identity and temporal programme are directly targeted by Notch in neuroblast tumours  
**Zacharioudaki, E., Housden, B. E., Garinis, G., Stojnic, R., Delidakis, C. and Bray, S. J.**

- 232 Single epicardial cell transcriptome sequencing identifies Caveolin 1 as an essential factor in zebrafish heart regeneration  
**Cao, J., Navis, A., Cox, B. D., Dickson, A. L., Gemberling, M., Karra, R., Bagnat, M. and Poss, K. D.**

- 244 The zebrafish tailbud contains two independent populations of midline progenitor cells that maintain long-term germ layer plasticity and differentiate in response to local signaling cues  
**Row, R. H., Tsotras, S. R., Goto, H. and Martin, B. L.**

## RESEARCH ARTICLES

- 255 The transcriptional repressor Blimp-1 acts downstream of BMP signaling to generate primordial germ cells in the cricket *Gryllus bimaculatus*  
**Nakamura, T. and Extavour, C. G.**
- 264 Sonic hedgehog controls enteric nervous system development by patterning the extracellular matrix  
**Nagy, N., Barad, C., Graham, H. K., Hotta, R., Cheng, L. S., Fejszak, N. and Goldstein, A. M.**

- 276 RING1 proteins contribute to early proximal-distal specification of the forelimb bud by restricting *Meis2* expression

**Yakushiji-Kaminatsui, N., Kondo, T., Endo, T. A., Koseki, Y., Kondo, K., Ohara, O., Vidal, M. and Koseki, H.**

- 286 Neurogenesis in sea urchin embryos and the diversity of deuterostome neurogenic mechanisms  
**Garner, S., Zysk, I., Byrne, G., Kramer, M., Moller, D., Taylor, V. and Burke, R. D.**

- 298 Neurogenic gene regulatory pathways in the sea urchin embryo  
**Wei, Z., Angerer, L. M. and Angerer, R. C.**

- 306 Two Forkhead transcription factors regulate cardiac progenitor specification by controlling the expression of receptors of the fibroblast growth factor and Wnt signaling pathways  
**Ahmad, S. M., Bhattacharyya, P., Jeffries, N., Gisselbrecht, S. S. and Michelson, A. M.**

- 318 Prox1 and fibroblast growth factor receptors form a novel regulatory loop controlling lens fiber differentiation and gene expression  
**Audette, D. S., Anand, D., So, T., Rubenstein, T. B., Lachke, S. A., Lovicu, F. J. and Duncan, M. K.**

- 329 A new level of plasticity: *Drosophila* smooth-like testes muscles compensate failure of myoblast fusion  
**Kuckwa, J., Fritzen, K., Buttgerit, D., Rothenbusch-Fender, S. and Renkawitz-Pohl, R.**

- 339 Dual function of Bmpr1a signaling in restricting preosteoblast proliferation and stimulating osteoblast activity in mouse  
**Lim, J., Shi, Y., Karner, C. M., Lee, S.-Y., Lee, W.-C., He, G. and Long, F.**

- 348 c-Raf promotes angiogenesis during normal growth plate maturation  
**Liu, E. S., Raimann, A., Chae, B. T., Martins, J. S., Baccarini, M. and Demay, M. B.**

- 356 Crim1 regulates integrin signaling in murine lens development  
**Zhang, Y., Fan, J., Ho, J. W. K., Hu, T., Kneeland, S. C., Fan, X., Xi, Q., Sellarole, M. A., de Vries, W. N., Lu, W., Lachke, S. A., Lang, R. A., John, S. W. M. and Maas, R. L.**