



Cover: Craniofacial skeleton preparation after Alcian Blue staining of a 4-day-old *Xenopus* tadpole. Craniofacial structures are mainly derived from migratory neural crest cells. The migration of neural crest cells is dependent on dermatan sulfate. A failure in proper neural crest cell migration might explain the craniofacial anomalies and other congenital malformations in musculocontractural Danlos–Ehlos syndrome, which results from a defect in dermatan sulfate biosynthesis. See article by Gouignard et al. on page 607. 'Dermatan sulfate in neural crest cell migration' by N. Gouignard is licenced under a Creative Commons Attribution 4.0 International licence.

EDITORIAL

- 597 Modeling human diseases: an education in interactions and interdisciplinary approaches
Zon, L.

SPECIAL ARTICLE

- 601 Show and tell: disclosure and data sharing in experimental pathology
Schofield, P. N., Ward, J. M. and Sundberg, J. P.

RESEARCH ARTICLES

- 607 Musculocontractural Ehlers–Danlos syndrome and neurocristopathies: dermatan sulfate is required for *Xenopus* neural crest cells to migrate and adhere to fibronectin
Gouignard, N., Maccarana, M., Strate, I., von Stedingk, K., Malmström, A. and Pera, E. M.

- 621 Identification of benzopyrone as a common structural feature in compounds with anti-inflammatory activity in a zebrafish phenotypic screen
Robertson, A. L., Ogryzko, N. V., Henry, K. M., Loynes, C. A., Foulkes, M. J., Meloni, M. M., Wang, X., Ford, C., Jackson, M., Ingham, P. W., Wilson, H. L., Farrow, S. N., Solari, R., Flower, R. J., Jones, S., Whyte, M. K. B. and Renshaw, S. A.

- 633 Histone lysine crotonylation during acute kidney injury in mice
Ruiz-Andres, O., Sanchez-Niño, M. D., Cannata-Ortíz, P., Ruiz-Ortega, M., Egido, J., Ortíz, A. and Sanz, A. B.

- 647 Two different pathogenic mechanisms, dying-back axonal neuropathy and pancreatic senescence, are present in the YG8R mouse model of Friedreich's ataxia
Mollá, B., Riveiro, F., Bolinches-Amorós, A., Muñoz-Lasso, D. C., Palau, F. and González-Cabo, P.

- 659 A novel *Drosophila* model of TDP-43 proteinopathies: N-terminal sequences combined with the Q/N domain induce protein functional loss and locomotion defects

Langellotti, S., Romano, V., Romano, G., Klima, R., Feiguin, F., Cagnaz, L., Romano, M. and Baralle, F. E.

- 671 Cellular dynamics of regeneration reveals role of two distinct Pax7 stem cell populations in larval zebrafish muscle repair
Pipalia, T. G., Koth, J., Roy, S. D., Hammond, C. L., Kawakami, K. and Hughes, S. M.

- 685 High-throughput screening for modulators of ACVR1 transcription: discovery of potential therapeutics for fibrodysplasia ossificans progressiva
Cappato, S., Tonachini, L., Giacopelli, F., Tirone, M., Galletta, L. J. V., Sormani, M., Giovenzana, A., Spinelli, A. E., Canciani, B., Brunelli, S., Ravazzolo, R. and Bocciardi, R.

- 697 Muscles provide protection during microbial infection by activating innate immune response pathways in *Drosophila* and zebrafish
Chatterjee, A., Roy, D., Patnaik, E. and Nongthomba, U.

- 707 TLR-mediated albuminuria needs TNF α -mediated cooperativity between TLRs present in hematopoietic tissues and CD80 present on non-hematopoietic tissues in mice
Jain, N., Khullar, B., Oswal, N., Banoth, B., Joshi, P., Ravindran, B., Panda, S., Basak, S., George, A., Rath, S., Bal, V. and Sopory, S.