



Cover: A model of the flagellum and flagellar pocket area of an African trypanosome. The model is derived from a series of three electron tomograms. The model illustrates the flagellum basal body and pro-basal body with associated rootlet of four microtubules (blue). The single Golgi complex (yellow) is closely associated with the neck region of the flagellar pocket. The endoplasmic reticulum intertwines between structures and around the single mitochondrion (orange). See article by S. Lacombe et al. (pp. 1081-1090).

Cell Science at a Glance

1055 **HIF at a glance.** Brahimi-Horn, M. C. and Pouyssegur, J.

Commentaries

1059 **Adhesion signaling – crosstalk between integrins, Src and Rho.** Huvencers, S. and Danen, E. H. J.

1071 **How to scaffold the contractile ring for a safe cytokinesis – lessons from Anillin-related proteins.** D’Avino, P. P.

Research Articles

1081 **Three-dimensional cellular architecture of the flagellar pocket and associated cytoskeleton in trypanosomes revealed by electron microscope tomography.** Lacombe, S., Vaughan, S., Gadelha, C., Morphew, M. K., Shaw, M. K., McIntosh, J. R. and Gull, K.

1091 **A Golgi-associated protein 4.1B variant is required for assimilation of proteins in the membrane.** Kang, Q., Wang, T., Zhang, H., Mohandas, N. and An, X.

1100 **The nuclear pore component Nup358 promotes transportin-dependent nuclear import.** Hutten, S., Wälde, S., Spillner, C., Hauber, J. and Kehlenbach, R. H.

1111 **Nucleocytoplasmic shuttling of soluble tubulin in mammalian cells.** Akoumianaki, T., Kardassis, D., Polioudaki, H., Georgatos, S. D. and Theodoropoulos, P. A.

1119 **Necdin is expressed in cachectic skeletal muscle to protect fibers from tumor-induced wasting.** Sciorati, C., Touvier, T., Buono, R., Pessina, P., François, S., Perrotta, C., Meneveri, R., Clementi, E. and Brunelli, S.

1126 **Bax inhibitor 1 regulates ER-stress-induced ROS accumulation through the regulation of cytochrome P450 2E1.** Kim, H.-R., Lee, G.-H., Cho, E. Y., Chae, S.-W., Ahn, T. and Chae, H.-J.

1134 **Stability of the small γ -tubulin complex requires HCA66, a protein of the centrosome and the nucleolus.** Fant, X., Gnadt, N., Haren, L. and Merdes, A.

1145 **NDST1-dependent heparan sulfate regulates BMP signaling and internalization in lung development.** Hu, Z., Wang, C., Xiao, Y., Sheng, N., Chen, Y., Xu, Y., Zhang, L., Mo, W., Jing, N. and Hu, G.

1155 **Low-density lipoprotein receptor-related protein 1 is an essential receptor for myelin phagocytosis.** Gaultier, A., Wu, X., Le Moan, N., Takimoto, S., Mukandala, G., Akassoglou, K., Campana, W. M. and Gonias, S. L.

1163 ***S. pombe btn1*, the orthologue of the Batten disease gene *CLN3*, is required for vacuole protein sorting of Cpy1p and Golgi exit of Vps10p.** Codlin, S. and Mole, S. E.

1174 **Targeting of p0071 to the midbody depends on KIF3.** Keil, R., Kiebling, C. and Hatzfeld, M.

1184 **Redundant and differential regulation of multiple licensing factors ensures prevention of re-replication in normal human cells.** Sugimoto, N., Yoshida, K., Tatsumi, Y., Yugawa, T., Narisawa-Saito, M., Waga, S., Kiyono, T. and Fujita, M.

1192 **Knockout of GARPs and the β -subunit of the rod cGMP-gated channel disrupts disk morphogenesis and rod outer segment structural integrity.** Zhang, Y., Molday, L. L., Molday, R. S., Sarfare, S. S., Woodruff, M. L., Fain, G. L., Kraft, T. W. and Pittler, S. J.

1201 **Cell-cycle regulation and dynamics of cytoplasmic compartments containing the promyelocytic leukemia protein and nucleoporins.** Jul-Larsen, Å., Grudic, A., Bjerkvig, R. and Bøe, S. O.

1211 **Activity of the AMPA receptor regulates drebrin stabilization in dendritic spine morphogenesis.** Takahashi, H., Yamazaki, H., Hanamura, K., Sekino, Y. and Shirao, T.

1220 **A Stim1-dependent, noncapacitative Ca^{2+} -entry pathway is activated by B-cell-receptor stimulation and depletion of Ca^{2+} .** Morita, T., Tanimura, A., Baba, Y., Kurosaki, T. and Tojyo, Y.

1229 **An essential role for CIC-4 in transferrin receptor function revealed in studies of fibroblasts derived from *Cln4*-null mice.** Mohammad-Panah, R., Wellhauser, L., Steinberg, B. E., Wang, Y., Huan, L. J., Liu, X.-D. and Bear, C. E.

1238 **Heterodimerization controls localization of Duox-DuoxA NADPH oxidases in airway cells.** Luxen, S., Noack, D., Frausto, M., Davanture, S., Torbett, B. E. and Knaus, U. G.

1248 **PP2A regulates BMP signalling by interacting with BMP receptor complexes and by dephosphorylating both the C-terminus and the linker region of Smad1.** Bengtsson, L., Schwappacher, R., Roth, M., Boergemann, J. H., Hassel, S. and Knaus, P.

1258 **p300-mediated acetylation of the Rothmund-Thomson-syndrome gene product RECQL4 regulates its subcellular localization.** Dietschy, T., Shevelev, I., Pena-Diaz, J., Hühn, D., Kuenzle, S., Mak, R., Miah, M. F., Hess, D., Fey, M., Hottiger, M. O., Janscak, P. and Stagljar, I.