



**Cover:** Division process of the parasite *Toxoplasma gondii*, which consists of the formation of two new daughter cells inside a mother cell. When daughter cells reach maturity, they emerge from the mother body and require the apicoplast fatty acid biosynthesis products to form a new plasma membrane between the new daughter cells. Parasites are labelled for IMC1 (red; inner membrane complex) and GAP45 (green; a marker for mature pellicle), and with DAPI (blue; nucleus). See article by Martins-Duarte et al. (pp. 3320–3331).

### EDITORIAL

- 3205** The good, the bad and the median  
**Way, M. (Editor-in-Chief)**

### STICKY WICKET

- 3207** Mole's Wow! So now you have your own lab!  
Part XI – Lightning strikes!

### CELL SCIENTISTS TO WATCH

- 3209** Cell scientist to watch – Emmanuel Boucrot

### CELL SCIENCE AT A GLANCE

- 3211** Cellular functions of the ADF/cofilin family at a glance  
**Kanellos, G. and Frame, M. C.**

### COMMENTARY

- 3219** The LINC and NPC relationship – it's complicated!  
**Jahed, Z., Soheilypour, M., Peyro, M. and Mofrad, M. R. K.**

### RESEARCH ARTICLES

- 3231** Sar1 localizes at the rims of COPII-coated membranes *in vivo*  
**Kurokawa, K., Suda, Y. and Nakano, A.**
- 3238** Microtubule-independent secretion requires functional maturation of Golgi elements  
**Fourriere, L., Divoux, S., Roceri, M., Perez, F. and Boncompain, G.**
- 3251** COPI is essential for Golgi cisternal maturation and dynamics  
**Ishii, M., Suda, Y., Kurokawa, K. and Nakano, A.**
- 3262** Keratin impact on PKC $\delta$ - and ASMase-mediated regulation of hepatocyte lipid raft size – implication for FasR-associated apoptosis  
**Gilbert, S., Loranger, A., Omary, M. B. and Marceau, N.**

- 3274** The microtubule-severing protein fidgetin acts after dendrite injury to promote their degeneration  
**Tao, J., Feng, C. and Rolls, M. M.**

- 3282** *Drosophila* Ringmaker regulates microtubule stabilization and axonal extension during embryonic development  
**Mino, R. E., Rogers, S. L., Risinger, A. L., Rohena, C., Banerjee, S. and Bhat, M. A.**

- 3295** The role of Sema3–Npn-1 signaling during diaphragm innervation and muscle development  
**Saller, M. M., Huettl, R.-E., Hanuschick, P., Amend, A.-L., Alberton, P., Aszodi, A. and Huber, A. B.**

- 3309** Divergence of structural strategies for homophilic E-cadherin binding among bilaterians  
**Nishiguchi, S., Yagi, A., Sakai, N. and Oda, H.**

- 3320** Apicoplast fatty acid synthesis is essential for pellicle formation at the end of cytokinesis in *Toxoplasma gondii*  
**Martins-Duarte, É. S., Carias, M., Vommaro, R., Suroliá, N. and de Souza, W.**

- 3332** A role for the yeast CLIP170 ortholog, the plus-end-tracking protein Bik1, and the Rho1 GTPase in Snc1 trafficking  
**Boscheron, C., Caudron, F., Loeillet, S., Peloso, C., Mugnier, M., Kurzawa, L., Nicolas, A., Denarier, E., Aubry, L. and Andrieux, A.**

- 3342** Trafficking of MHC molecules to the cell surface creates dynamic protein patches  
**Blumenthal, D., Edidin, M. and Gheber, L. A.**

### TOOLS AND TECHNIQUES

- 3351** Non-invasive single-cell biomechanical analysis using live-imaging datasets  
**Pearson, Y. E., Lund, A. W., Lin, A. W. H., Ng, C. P., Alsuwaidi, A., Azzeh, S., Gater, D. L. and Teo, J. C. M.**