

## Transepithelial potential differences and Na<sup>+</sup> flux in isolated perfused gills of the crab *Chasmagnathus granulatus* (Grapsidae) acclimated to hyper- and hypo-salinity

C. M. Luquet<sup>1,\*</sup>, U. Postel<sup>2</sup>, J. Halperin<sup>1,3</sup>, M. R. Urcola<sup>1</sup>, R. Marques<sup>4</sup> and D. Siebers<sup>2</sup>

<sup>1</sup>Department of Biological Sciences, FCEN University of Buenos Aires, Building II, Ciudad Universitaria, C1428EHA Buenos Aires, Argentina, <sup>2</sup>Alfred Wegener Institute for Polar and Marine Research, Columbusstrasse, D-27568 Bremerhaven, Germany, <sup>3</sup>CONICET-Argentina and <sup>4</sup>Department of Inorganic Chemistry, FCEN University of Buenos Aires, Argentina

\*e-mail: luquet@bg.fcen.uba.ar

Accepted 1 October 2001

### Summary

We studied the transepithelial potential difference (TEPD) and <sup>22</sup>Na flux across isolated perfused gills (anterior pair 5 and posterior pairs 6–8) of the crab *Chasmagnathus granulatus* acclimated to either hypo- or hyper-osmotic conditions.

The gills of crabs acclimated to low salinity, perfused and bathed with 10‰ saline solutions, produced the following TEPDs (hemolymph side with respect to bath side): 0.4±0.7, -10.2±1.6, -10.8±1.3 and -6.7±1.3 mV for gills 5, 6, 7 and 8, respectively. Gills 6, 7 and 8 did not differ significantly. Reducing the saline concentration of bath and perfusate from 30‰ to 20‰ or 10‰ increased significantly the TEPDs of these gills. TEPDs of gill 6 (representative of posterior gills) were reduced by 69±5%

and 60±5% after perfusion with ouabain or BaCl<sub>2</sub> (5 mmol l<sup>-1</sup> each), respectively. The same gill showed a net ouabain-sensitive Na<sup>+</sup> influx of 1150±290 µequiv g<sup>-1</sup> h<sup>-1</sup>.

Gill 6 of crabs acclimated to high salinity produced TEPDs of -1.5±0.1 and -1.3±0.09 mV after perfusion with 30‰ or 40‰ salines, respectively. Perfusion with ouabain or BaCl<sub>2</sub> reduced TEPDs by 76±7% and 86±4%, respectively. A net ouabain-sensitive Na<sup>+</sup> efflux of 2282±337 µequiv g<sup>-1</sup> h<sup>-1</sup> was recorded in gill 6 perfused with 38‰ saline.

Key words: isolated perfused gill, transepithelial potential difference, Na<sup>+</sup>K<sup>+</sup>-ATPase, ion flux, hypo-regulation, hyper-regulation, ouabain, crab, *Chasmagnathus granulatus*.

### Introduction

Semiterrestrial estuarine crabs are adapted to a wide range of conditions. Water salinity varies from full-strength seawater to almost freshwater in a geographical and tidal basis. Excursions to intertidal and supratidal areas include the chance of entering into rain pools or into tide pools concentrated by evaporation. Moreover, aerial ventilation may cause evaporative concentration of water retained within the branchial chambers, leading to hyper-saline exposure of the gills and the gill chamber epithelium (Schmitt and Santos, 1993). Terrestrial and semiterrestrial crabs of marine or estuarine origin possess hyper- and hypo-regulation capacities, showing little variation in their internal osmotic and ionic concentrations within a wide range of salinity (Gross, 1964; Mantel and Farmer, 1983; Schubart and Diesel, 1998, 1999).

The gills of euryhaline crabs are histologically and functionally differentiated. While anterior gills are lined with thin epithelium and have a mainly respiratory function, the posterior ones play a key role in compensatory active uptake of ions (Siebers et al., 1982; Gilles and Péqueux, 1986; Towle and Kays, 1986; Compère et al., 1989). The predominant cells

in posterior gills are thick and possess large basolateral membrane interdigitations together with mitochondria (Copeland and Fitzjarrell, 1968; Compère et al., 1989; Luquet et al., 1997). There is much evidence from electrophysiological and ion flux studies on the ion-uptake capacity of posterior gills of different crab species (Gilles and Péqueux, 1981, 1985; Péqueux et al., 1988; Siebers et al., 1985; Lucu and Siebers, 1986; Burnett and Towle, 1990) (for a review, see Péqueux, 1995).

Two models for the mechanisms involved in such active ion transport through the gills, have been reviewed recently (Onken and Riestenpatt, 1998). (1) In the gills of species such as *Carcinus maenas*, which possess limited capacity to invade low-salinity environments (weak hyper-regulators), it is proposed that the ions enter the cell by crossing the apical membrane through Na<sup>+</sup>/2Cl<sup>-</sup>/K<sup>+</sup> symports coupled to K<sup>+</sup> channels. (2) In strong hyper-regulators, such as *Eriocheir sinensis*, Onken and Riestenpatt suggest that Na<sup>+</sup> crosses the apical membrane through epithelial Na<sup>+</sup> channels while Cl<sup>-</sup> is exchanged with HCO<sub>3</sub><sup>-</sup>, driven by an apical H<sup>+</sup>-V-ATPase. At

the basolateral side, Na<sup>+</sup>K<sup>+</sup>-ATPase and K<sup>+</sup> and Cl<sup>-</sup> channels are thought to drive ions into the hemolymph in both weak and strong hyper-regulating species.

Early work on salt and water regulation by intact fiddler crabs of the genus *Uca* (Green et al., 1959; Baldwin and Kirschner, 1976a; Evans et al., 1976) suggests that active ion excretion follows an extra-renal route. There is some physiological and histological evidence to suggest that the gills of hypo-regulating crabs are the organs involved in this function. Martínez et al. (1998), working with isolated perfused gills of *Ucides cordatus*, reported that gill 6 is capable of active ion excretion, while gill 5 is specialized in ion uptake. Ultrastructural studies also suggested an ion excretion capacity of the gills of crabs acclimated to hypersaline media; Martelo and Zanders (1986) and Luquet et al. (1997) described a cell architecture characteristic of an ion-excreting epithelium in the gills of grapsids and ocpodids. Some features of this cell architecture, such as the apparently low-resistance cell junctions, resemble those of the extensively studied vertebrate salt-secreting organs such as the avian salt gland (Riddle and Ernst, 1979), the teleost opercular epithelium (Ernst et al., 1980) and the rectal gland of elasmobranchs (Ernst et al., 1981).

*Chasmagnathus granulatus* Dana 1851 is a strong ion hyper- and hypo-regulating crab species that inhabits intertidal estuarine coasts of Brazil, Uruguay and Argentina (Boschi, 1964; Mougabure Cueto, 1998). The posterior gills of this species are believed to be involved in both ion uptake and excretion, since their epithelium thickness is increased to the same extent after transfer from full seawater to either dilute or concentrated seawater (Genovese et al., 2000).

The aim of this work was to establish, by electrophysiological and ion tracer flux experiments, the possible role of the different gill pairs of *C. granulatus* in ion-transport functions at low and high salinity. The participation of Na<sup>+</sup>K<sup>+</sup>-ATPase and K<sup>+</sup> channels was also studied.

## Materials and methods

### Animals

Crabs were collected from Faro San Antonio beach (36°18'S 56°48'W), near the southern edge of the Rio de la Plata estuary, Argentina. Once in the laboratory, the animals were separated at random into two groups and acclimated in glass containers with aerated seawater of either 12‰ or 45‰ salinity for at least 2 weeks. Laboratory temperature was kept at 20±2°C. Stage C intermoult adult male crabs (Drach and Tchernigovtzeff, 1967) of 26±0.3 mm carapace widths were selected for the study.

### Gill perfusion

Crabs were killed by destroying the ventral nervous ganglion with a spike. After removing the dorsal carapace, the gills were gently excised and placed in a Petri dish with saline solution. The afferent and efferent vessels of gills 5–8 were connected by fine polyethylene tubing of 0.4 mm diameter to a peristaltic

Table 1. Composition of the saline solutions used in each experiment

Salinity (%)	Component (mmol l <sup>-1</sup> )					
	NaCl	KCl	MgCl <sub>2</sub>	CaCl <sub>2</sub>	Hepes	NaHCO <sub>3</sub>
10	156	3.15	2.50	4.17	5.00	2.50
20	312	6.30	4.66	8.35	5.00	2.50
30	468	9.46	7.50	12.53	5.00	2.50
32	500	10.08	8.00	13.36	5.00	2.50
34	531	10.70	8.50	14.20	5.00	2.50
36	562	11.27	9.00	15.03	5.00	2.50
38	593	11.96	9.50	15.87	5.00	2.50
40	625	12.59	10.00	16.71	5.00	2.50

Perfusate also included 2 mmol l<sup>-1</sup> glucose.

pump (afferent) and to a glass tube (efferent). Perfusion rate was kept at 0.1 ml min<sup>-1</sup>. The tubing was held in position by an acrylic clamp and the preparation put into a glass beaker with the appropriate saline solution and constant aeration.

Gills 5–8 from crabs acclimated to low (12‰) and high (45‰) salinity were perfused and bathed with identical solutions, except that the perfusate contained 2 mmol l<sup>-1</sup> glucose. Table 1 shows the composition of the saline solutions used in the different experiments. All solutions were adjusted with Tris-base to the physiological pH 7.75 for *C. granulatus* (Luquet and Ansaldo, 1997). The effects of the following drugs were tested in gills perfused with 10‰ saline for low-salinity crabs and 38‰ saline for high-salinity crabs: ouabain (a specific Na<sup>+</sup>K<sup>+</sup>-ATPase inhibitor) (Skou, 1965), 5 mmol l<sup>-1</sup> applied basolaterally, and BaCl<sub>2</sub> and CsCl (K<sup>+</sup> channel blockers) (Zeiske, 1990; Draber and Hansen, 1994), 5–10 mmol l<sup>-1</sup> applied at both apical and basolateral sides.

### Trans epithelial potential difference (TEPD)

Ag/AgCl electrodes were connected by agar bridges to the external bath and to the glass tube collecting the perfusate. Potential differences were measured in the internal perfusate with respect to the external bath by means of a millivoltmeter.

### Na<sup>+</sup> flux

Na<sup>+</sup> outward and inward movements were measured in the same gill by applying <sup>22</sup>Na first in the perfusate and afterwards in the bathing solution. Identical ionic concentrations (20‰ or 38‰, for low- and high-salinity crabs, respectively) were used in the bath and perfusate. For Na<sup>+</sup> efflux, <sup>22</sup>Na was included in the perfusate at a final concentration of 0.25 µCi ml<sup>-1</sup>. After stabilization for 15 min, three samples (1 ml each) were collected from the perfusate and the bath at intervals of 15 min. Na<sup>+</sup> efflux was calculated from the radioactivity that appeared in the bath. To avoid overestimated flux due to a possible leak, we also measured the radioactivity lost by the perfusate after passing through the gill. Gills showing large discrepancies between both methods of measurement were discarded. Results from measurements of radioactivity that appeared in the bath were chosen for the study.

After washing with non-radioactive solution, Na<sup>+</sup> influx was measured by applying <sup>22</sup>Na in the bath at a final concentration of 9 kBq (0.25 µCi) ml<sup>-1</sup>. After stabilization for 15 min, radioactivity was measured in the collected perfusate at 10 min intervals during the subsequent 30 min.

Radioactivity was measured with a Canberra Series 35 plus gamma scintillation counter. Na<sup>+</sup> efflux and influx were calculated according to the formula of Lucu and Siebers (1986):

$$J = {}^{22}\text{Na} \times S / \text{SRA} \times m, \quad (1)$$

where *J* is the calculated unidirectional flux of Na<sup>+</sup> in µequiv h<sup>-1</sup> g<sup>-1</sup>; <sup>22</sup>Na is the radioactivity (cts min<sup>-1</sup>) collected during each interval; *S* is the number of samples collected during 1 h (6 and 4 for influx and efflux, respectively); *SRA* is the specific radioactivity (cts min<sup>-1</sup> µequiv<sup>-1</sup>) and *m* is the fresh mass of the gill (g).

*Chemicals*

NaCl, KCl, MgCl<sub>2</sub>, BaCl<sub>2</sub>, KCN and glucose were obtained from Merck Argentina; NaHCO<sub>3</sub> was obtained from Mallinckrodt USA; CaCl<sub>2</sub> and 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (Hepes) were purchased from J. T. Baker, USA; ouabain was purchased from Sigma USA and <sup>22</sup>Na was obtained from Amersham Pharmacia Biotech.

*Statistics*

Data were analyzed by one- or two-way repeated measures analysis of variance (ANOVA) or paired *t*-test when appropriate (Sokal and Rohlf, 1981). All values are means ± S.E.M.

**Results**

The preparations of isolated perfused gills remained viable for at least 3 h, since stable values of transepithelial potential difference (TEPDs) could be recorded along this period. The rate of perfusion was also constant during each experiment.

*Crabs acclimated to low salinity*

*Transepithelial potential difference*

Posterior gills of crabs acclimated to low salinity, perfused and bathed with identical salines, showed hemolymph-side negative TEPDs, which increased significantly in absolute value as the concentration of saline solutions decreased; no significant TEPD was recorded in gill 5 (representative of anterior gills) (Fig. 1). Two-way repeated measures ANOVA comparing gills 6, 7 and 8 indicated that saline concentration was the only significant source of variation (*P*<0.001, *N*=6 for each gill). TEPDs for gill 6 ranged between -2.4±0.5 mV (*N*=6) in 30‰ saline and -10.2±1.6 mV (*N*=6) in 10‰ saline. Since the three posterior gills responded similarly to the different saline concentrations, gill 6 was chosen as representative for further experiments because it was bigger than gills 7 and 8 and therefore easier to handle.

Basolaterally applied ouabain (5 mmol l<sup>-1</sup>) significantly reduced the TEPD of gill 6 by 69±5% (*P*<0.001, *N*=7).




Fig. 1. Transepithelial potential differences (TEPD) of gills of crabs acclimated to low salinity, perfused with different saline concentrations. TEPD of posterior gills increased significantly in absolute value at lower concentrations (*P*<0.001, *N*=6 for each gill). TEPD of gill 5 did not differ significantly from 0 mV (*P*>0.05).

5 mmol l<sup>-1</sup> BaCl<sub>2</sub> applied basolaterally caused a similar inhibitory effect of 60±5% (*P*<0.001, *N*=5). The same inhibitor was totally ineffective when applied at the apical (bath) side, even when the concentration was raised to 10 mmol l<sup>-1</sup>. In preliminary experiments CsCl (10 mmol l<sup>-1</sup>) was also ineffective at the apical side (data not shown). The effects of ouabain and BaCl<sub>2</sub> were totally reversed after washing out with normal saline solution (Table 2). Results from single experiments are shown in Fig. 2.

*Na<sup>+</sup> influx*

A net Na<sup>+</sup> influx of 1150±290 µequiv g<sup>-1</sup> h<sup>-1</sup> (*N*=6) was calculated for isolated gills 6 of crabs acclimated to low salinity, perfused and bathed with 20‰ saline. Unidirectional flux of Na<sup>+</sup> was 719±163 µequiv g<sup>-1</sup> h<sup>-1</sup> (*N*=6) in the outward (*J*<sub>out</sub>) direction and 1824±410 µequiv g<sup>-1</sup> h<sup>-1</sup> (*N*=6) in the inward direction (*J*<sub>in</sub>). The *J*<sub>in</sub>/*J*<sub>out</sub> ratio calculated from these results (2.87±0.67) was significantly different (*P*<0.05) from the ratio predicted by the Ussing equation (Ussing, 1949), which was 1.33±0.05. Ouabain (5 mmol l<sup>-1</sup>) added to the perfusate caused a reduction of 35±5% in *J*<sub>in</sub> (*P*<0.05, *N*=5); inhibition of the net Na<sup>+</sup> influx calculated from this value was 70±20%, *N*=5 (Fig. 3).

Table 2. Effect of different ion-transport inhibitors on the transepithelial potential difference (TEPD) of gill 6 of crabs acclimated to low salinity

TEPD (mV)	Ouabain			BaCl <sub>2</sub>		
	Control	Inhibitor	Wash out	Control	Inhibitor	Wash out
Mean	-6.6	-2.1	-6.8	-9	-3.8	-9
S.E.M.	0.8	0.5	0.8	1	0.8	1
<i>N</i>		7			5	

Ouabain and BaCl<sub>2</sub> (5 mmol l<sup>-1</sup>) were each applied basolaterally.




Fig. 2. Effects of ouabain and BaCl<sub>2</sub> on the transepithelial potential difference (TEPD) of individual gills 6 from a crab acclimated to low salinity, perfused with 10‰ saline.

#### Crabs acclimated to high salinity

##### Transepithelial potential difference

In a preliminary series of experiments, gills 6, 7 and 8 of crabs acclimated to high salinity, perfused with either 30‰ or 40‰ saline, produced TEPDs of the same polarity but substantially smaller than those shown by posterior gills of crabs acclimated to low salinity, i.e.  $-1.5 \pm 0.1$  and  $-1.3 \pm 0.09$  mV ( $N=13$ ) for gill 6 perfused with 30‰ and 40‰ solutions, respectively. These values did not differ significantly. Gill 5 did not show any significant TEPD.

In a second series of experiments, gill 6 was used as representative of posterior gills and the concentration of the bath and perfusate was increased stepwise from 30‰ to 40‰, in an attempt to minimize the effects of a possible osmotic shock. A slight tendency to increased TEPD at higher saline concentrations was observed, with the highest values recorded at 36–38‰ ( $N=9$ ). Nevertheless, these differences were not statistically significant (Fig. 4).

Effects of inhibitors are summarized in Table 3. Basolateral application of  $5 \text{ mmol l}^{-1}$  ouabain significantly inhibited the




Fig. 4. Transepithelial potential difference (TEPD) of gill 6 from crabs acclimated to high salinity, perfused at different saline concentrations ( $N=9$ ).

TEPD by  $76 \pm 7\%$  ( $P < 0.005$ ,  $N=6$ ). This effect was hardly reversed after washing out with normal saline. Similarly,  $10 \text{ mmol l}^{-1}$  BaCl<sub>2</sub> applied at the basolateral side had an inhibitory effect of  $86 \pm 4\%$  ( $P < 0.001$ ,  $N=5$ ) but did not cause consistent effects when applied at the apical side ( $N=9$ ). Another series of experiments was performed to test whether the effects of BaCl<sub>2</sub> were due to diffusive potentials caused by the asymmetrical addition of Cl<sup>-</sup> ions or by actual inhibition of K<sup>+</sup> channels. Choline chloride ( $20 \text{ mmol l}^{-1}$ ) applied basolaterally caused a significant decrease in TEPD of  $36 \pm 6\%$  ( $P < 0.01$ ,  $N=5$ ), whereas BaCl<sub>2</sub> under the same conditions produced a significantly higher effect of  $84 \pm 5\%$  ( $P < 0.001$ ,  $N=5$ ). Fig. 5 shows the results of individual experiments with ouabain, BaCl<sub>2</sub> and choline chloride.

##### Na<sup>+</sup> efflux

A mean net Na<sup>+</sup> efflux of  $2282 \pm 337 \mu\text{equiv g}^{-1} \text{ h}^{-1}$  ( $N=9$ ) was calculated for gill 6 of crabs acclimated to high salinity, perfused and bathed with 38‰ saline. Unidirectional Na<sup>+</sup> efflux ( $J_{\text{out}}$ ) was  $2900 \pm 329 \mu\text{equiv g}^{-1} \text{ h}^{-1}$  ( $N=9$ ), while unidirectional influx ( $J_{\text{in}}$ ) measured in the same preparations




Fig. 3. Unidirectional Na<sup>+</sup> fluxes of gills 6 from crabs acclimated to low and high salinity, perfused with 20 and 38‰ saline, respectively. Ouabain ( $5 \text{ mmol l}^{-1}$ ) added to the perfusate significantly inhibited both  $J_{\text{in}}$  of crabs acclimated to low salinity ( $P < 0.05$ ,  $N=5$ ) and  $J_{\text{out}}$  of crabs acclimated to high salinity ( $P < 0.001$ ,  $N=4$ ).




Fig. 5. Effects of ouabain, BaCl<sub>2</sub> and choline chloride on the transepithelial potential difference (TEPD) of individual gills 6 from a crab acclimated to high salinity, perfused with 38‰ saline. Inhibition was almost irreversible.

Table 3. Effect of basolateral application of ion-transport inhibitors on the transepithelial potential difference (TEPD) of gill 6 of crabs acclimated to high salinity

TEPD (mV)	Ouabain (5 mmol l <sup>-1</sup> )			KCN (2 mmol l <sup>-1</sup> )			BaCl <sub>2</sub> (10 mmol l <sup>-1</sup> )		
	Control	Inhibitor	Wash out	Control	Inhibitor	Wash out	Control	Inhibitor	Wash out
Mean	-0.92	-0.2	-	-1.45	-0.53	-1.3	-1.64	-0.22	-1.47
S.E.M.	0.13	0.7	-	0.12	0.10	0.7	0.27	0.6	0.26
N		6			6			5	

was  $619 \pm 73 \mu\text{equiv g}^{-1} \text{h}^{-1}$  ( $N=9$ ). The observed  $J_{\text{out}}/J_{\text{in}}$  ratio ( $4.48 \pm 0.91$ ) was significantly different ( $P < 0.05$ ) from the ratio predicted by the Ussing equation ( $0.95 \pm 0.01$ ).  $J_{\text{out}}$  was  $58 \pm 9\%$  inhibited by basolaterally applied  $5 \text{ mmol l}^{-1}$  ouabain ( $P < 0.001$ ,  $N=4$ ); inhibition of net efflux calculated from this value was  $81 \pm 6\%$ ,  $N=4$  (Fig. 3).

### Discussion

Ion-uptake functions of crab gills have been extensively studied in the hyper-regulators *Carcinus maenas*, *Eriocheir sinensis*, *Callinectes sapidus*, (Burnett and Towle, 1990; Gilles and Péqueux, 1981; Onken and Siebers, 1992; Siebers et al., 1985) (for reviews, see Péqueux, 1995; Onken and Riestenpatt, 1998) and the hyper-hyporegulators *Uca tangeri* and *Pachygrapsus marmoratus* (Drews and Graszynski, 1987; Krippeit Drews et al., 1989; Pierrot et al., 1995a,b). Ion excretion has received much less attention. Early studies conducted on fiddler crabs acclimated to hyper-osmotic media (Green et al., 1959; Baldwin and Kirschner, 1976a,b; Evans et al., 1976) demonstrated active excretion of Na<sup>+</sup> and Cl<sup>-</sup> by the whole animal, but the organ responsible for this function is still to be determined. Martínez et al. (1998) recorded active Na<sup>+</sup> excretion through the isolated perfused gills of *Ucides cordatus* acclimated to iso-osmotic media. The present paper presents a comparative study of ion-transport functions in the gills of the hyper-hyporegulating crab *Chasmagnathus granulatus* acclimated to either hypo-osmotic or hyper-osmotic conditions.

#### Acclimation to low salinity

Posterior gills of *C. granulatus* acclimated to low salinity, perfused and bathed with identical solutions, produce hemolymph-side negative TEPDs similar to those reported for hyper-regulating crabs (Lucu and Siebers, 1986; Siebers et al., 1985) and also for the hyper-hyporegulating crabs *Uca tangeri* and *Pachygrapsus marmoratus* (Drews and Graszynski, 1987; Krippeit Drews et al., 1989; Pierrot et al., 1995a,b). These TEPDs are sensitive to the dilution of the perfusion and bathing media, being the lowest absolute values measured at 30‰, which is near the physiological hemolymphatic concentrations of Na<sup>+</sup> and Cl<sup>-</sup> in this species (Mougabure Cueto, 1998). This change in potential difference seems to reflect an autoregulatory mechanism of gill tissue, which involves enhanced ion transport activity and/or increased paracellular resistance in response to hypo-osmotic stress. Onken (1996)

and Onken and Riestenpatt (1998) reported that changes in short-circuit currents measured in split gill lamellae of *Eriocheir sinensis* respond to osmotic variations at the basolateral side. Our results indicate that these changes in TEPD take place in *C. granulatus* within a few minutes and are totally reversed when the gill is perfused again with iso-ionic saline. This rapid response should be important for a species that is often observed emerging from brackish or seawater and entering into rain pools for feeding on supratidal plants.

Radioactive tracer flux suggests that posterior gills of this species actively take up Na<sup>+</sup> at low salinity. Both influx and efflux rates as well as net influx are somewhat high compared with data reported for other species in similar experimental conditions (Lucu and Siebers, 1986; Pierrot et al., 1995a). This high rate of Na<sup>+</sup> uptake is possibly a response to a high rate of ion loss by the animal. Gill and whole animal ionic permeability should be studied in order to test this hypothesis.

As a first approach to understanding the mechanisms involved in gill ion uptake, it can be concluded that Na<sup>+</sup>K<sup>+</sup>-ATPase located at the basolateral membrane is the major driving force, since both TEPD and Na<sup>+</sup> influx are inhibited by ouabain in similar proportions. The reason why ouabain does not cause total inhibition could be incomplete access of the drug to the enzyme molecules, due to the complex basolateral membrane interdigitations of gill ionocytes, as suggested by Burnett and Towle (1990) to explain similar results obtained with *Callinectes sapidus*.

As our results imply, barium-sensitive K<sup>+</sup> channels located in the basolateral membrane are also involved in generating the observed transepithelial potentials. The lack of effect of apical BaCl<sub>2</sub> and CsCl (preliminary data) suggests the absence of barium-sensitive K<sup>+</sup> channels in this membrane. It has been reported that these channels are necessary for electrogenic uptake of Na<sup>+</sup> and Cl<sup>-</sup> across the apical membrane through Na<sup>+</sup>/2Cl<sup>-</sup>/K<sup>+</sup> cotransporters in *Uca tangeri* and *Carcinus maenas* (Drews and Graszynski, 1987; Riestenpatt et al., 1996).

Onken and Riestenpatt (1998) have proposed that in strong hyper-regulators, such as *Eriocheir sinensis*, Na<sup>+</sup> and Cl<sup>-</sup> cross the apical membrane through Na<sup>+</sup> channels and Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> antiports, driven by an H<sup>+</sup>-V-ATPase. Although we have no direct evidence for apical transporters, expression of both Na<sup>+</sup>/H<sup>+</sup> exchangers and H<sup>+</sup>-V-ATPase has been detected in gill 6 of *C. granulatus* in preliminary molecular biology experiments (D. Weihrauch and C. M. Luquet, unpublished

observations). In addition, Genovese et al. (2000) reported increased activity of cytosolic carbonic anhydrase in posterior gills of *C. granulatus* after acclimation to low salinity. This enzyme is believed to produce  $H^+$  and  $HCO_3^-$  as counterions for  $Na^+$  and  $Cl^-$  exchangers, respectively, and is also important for the function of an  $H^+$ -V-ATPase (Henry, 1988; Henry and Swenson, 2000).

#### Acclimation to high salinity

The involvement of crab gills in hypo-regulatory ion excretion is still a matter of controversy. In the past two decades histological evidence has accumulated. There are reports of ultrastructural changes and increased posterior gill epithelium thickness after acclimation to hypersaline media (Martelo and Zanders, 1986; Luquet et al., 1997; Rosa et al., 1999; Genovese et al., 2000). Martínez et al. (1998) perfused gills 5 and 6 of *Ucides cordatus* acclimated in isosmotic medium, reporting net  $Na^+$  uptake by gill 5, even when the external saline was more concentrated than the perfusate. In contrast, they found net  $Na^+$  excretion by gill 6 at all concentrations tested. This experiment, however, is not comparable with the present results, since it was performed with asymmetrical perfusion. In addition, the authors acclimated the crabs to isosmotic medium. A clear difference does seem to exist between both species, however; whereas in *U. cordatus* different gills are specialized for transporting ions in opposite directions, the TEPDs measured on the three posterior gills of *C. granulatus* suggested that they have similar ion transport capacities. Thus gills 6, 7 and 8 of *C. granulatus* seem to be equally involved in both transport directions, after chronic acclimation to either low or high salinity.

At least two basolateral membrane proteins involved in ion uptake are also involved in ion excretion. These are  $Na^+K^+$ -ATPase, since both TEPD and  $Na^+$  flux are reduced by ouabain, and  $K^+$  channels, which are inhibited by basolateral application of  $BaCl_2$ . By contrast, our results provide no evidence for any apical ion-transport proteins.

Current models for salt excretion in vertebrates (gills, and opercular epithelia of teleost fish, rectal glands of elasmobranchs and avian salt glands) consider  $Na^+$  pumping into the paracellular space and a transcellular flux of  $Cl^-$ , which in turn generates a positive transepithelial potential difference that drives  $Na^+$  efflux via low-resistance tight junctions (Ernst et al., 1980, 1981; Lowy et al., 1987). Our results on crabs acclimated to high salinity indicate that  $Na^+K^+$ -ATPase is the main driving force for ion extrusion. In addition, previous electron microscopic work shows shorter septate junctions in gills of *Uca uruguayensis* and *C. granulatus* acclimated to high salinity (Luquet et al., 1997; Rosa et al., 1999) compared with gills of the same species acclimated to low salinity. Thus, paracellular flux of ions through these junctions also seems possible. However, the vertebrate model predicts a positive transepithelial potential difference for driving paracellular  $Na^+$  efflux. This is not the case for the gills of *C. granulatus*, which produce a little negative potential difference. Therefore the routes followed by  $Na^+$  and  $Cl^-$  at the apical side seem to differ

from known models for salt excretion and deserve further investigation.

This study was supported by a grant UBACYT TX 07. We acknowledge material support by the Instituto Antártico Argentino. We also wish to thank Susanne Luck-Kopp, Analía Pérez, Martín Ansaldo and Carolina Luchetti for their kind help and two anonymous reviewers for their helpful suggestions.

#### References

- Baldwin, G. F. and Kirschner, L. B. (1976a). Sodium and chloride regulation in *Uca* adapted to 175% sea water. *J. Exp. Zool.* **49**, 158–171.
- Baldwin, G. F. and Kirschner, L. B. (1976b). Sodium and chloride regulation in *Uca* adapted to 175% sea water. *J. Exp. Zool.* **49**, 172–180.
- Boschi, E. E. (1964). Los crustáceos decápodos brachyura del litoral bonaerense. *Bol. Instit. Biol. Marina (Mar del Plata)* **164**, 1–34.
- Burnett, L. E. and Towle, D. W. (1990). Sodium ion uptake by perfused gills of the blue crab *Callinectes sapidus*: effects of ouabain and amiloride. *J. Exp. Biol.* **149**, 293–305.
- Compère, Ph., Wanson, S., Péqueux, A., Gilles, R. and Goffinet, G. (1989). Ultrastructural changes in the gill epithelium of the green crab *Carcinus maenas* in relation to the external salinity. *Tissue Cell* **21**, 299–318.
- Copeland, D. E. and Fitzjarrell, A. T. (1968). The salt absorbing cells in the gills of the blue crab (*Callinectes sapidus* Rathbun) with notes on modified mitochondria. *Zeits. Zellf.* **92**, 1–22.
- Draber, S. and Hansen, U.-P. (1994). Fast single-channel measurements resolve the blocking effect of  $Cs^+$  on the  $K^+$  channel. *Biophys. J.* **67**, 120–129.
- Drach, P. and Tchernigovtzeff, C. (1967). Sur la méthode de détermination des stades d'intermue et son application générale aux crustacés. *Vie Milieu* **18**, 597–607.
- Drews, G. and Graszynsky, K. (1987). The transepithelial potential differences in the gills of the fiddler crab, *Uca tangeri*: influence of some inhibitors. *J. Comp. Physiol.* **157B**, 345–353.
- Ernst, S. A., Dodson, W. C. and Karnaky, K. J., Jr (1980). Structural diversity of occluding junctions in the low-resistance of seawater-adapted killifish (*Fundulus heteroclitus*). *J. Cell Biol.* **87**, 488–497.
- Ernst, S. A., Hootman, S. R., Schreiber, J. H. and Riddle, C. V. (1981). Freeze-fracture and morphometric analysis of occluding junctions in rectal glands of elasmobranch fish. *J. Membr. Biol.* **58**, 101–114.
- Evans, D. H., Cooper, K. and Bogan, M. B. (1976). Sodium extrusion by the seawater-acclimated fiddler crab *Uca pugilator*: comparison with other marine crustacea and marine teleost fish. *J. Exp. Biol.* **64**, 203–219.
- Genovese, G., Luquet, C. M., Paz, D. A., Rosa, G. A. and Pellerano, G. N. (2000). The morphometric changes in the gills of the estuarine crab *Chasmagnathus granulatus* under hyper- and hypo-regulation conditions are not caused by proliferation of specialized cells. *J. Anat.* **197**, 239–246.
- Gilles, R. and Péqueux, A. (1981). Cell volume regulation in Crustaceans: relationship between mechanisms for controlling the osmolality of extracellular fluids. *J. Exp. Zool.* **215**, 351–362.
- Gilles, R. and Péqueux, A. (1985). Ion transport in Crustacean gills: physiological and ultrastructural approaches. In *Transport Processes, Iono- and Osmoregulation: Current Comparative Approaches* (ed. R. Gilles and M. Gilles-Baillien), p. 136. Berlin, New York: Springer-Verlag.
- Gilles, R. and Péqueux, A. (1986). Physiological and ultrastructural studies of NaCl transport in crustacean gills. *Boll. Zool.* **53**, 173–182.
- Green, J. W., Harsch, M., Barr, L. and Prosser, C. L. (1959). The regulation of water and salt by the fiddler crabs, *Uca pugnax* and *Uca pugilator*. *Biol. Bull.* **116**, 76–87.
- Gross, W. J. (1964). Trends in water and salt regulation among aquatic and amphibious crabs. *Biol. Bull.* **127**, 447–466.
- Henry, R. P. (1988). Multiple functions of carbonic anhydrase in the crustacean gill. *J. Exp. Zool.* **248**, 19–24.
- Henry, R. P. and Swenson, E. R. (2000). The distribution and physiological significance of carbonic anhydrase in vertebrate gas exchange organs. *Respir. Physiol.* **121**, 1–12.
- Krippel-Drews, P., Drews, G. and Graszynski, K. (1989). Effects of ion substitution on the transepithelial potential difference of the gills of the

- fiddler crab *Uca tangeri*: evidence for a H<sup>+</sup>-pump in the apical membrane. *J. Comp. Physiol.* **159**, 43–49.
- Lowy, R. J., Dawson, D. C. and Ernst, S. A.** (1987). Mechanism of ion transport by avian salt gland primary cell cultures. *Am. J. Physiol.* **256**, R1184–R1191.
- Lucu, C. and Siebers, D.** (1986). Amilorid-sensitive sodium flux and potentials in perfused *Carcinus* gill preparations. *J. Exp. Biol.* **122**, 25–35.
- Luquet, C. and Ansaldo, M.** (1997). Acid–base balance and ionic regulation during emersion in the estuarine intertidal crab *Chasmagnathus granulata* Dana (Decapoda Grapsidae). *Comp. Biochem. Physiol.* **117A**, 407–410.
- Luquet, C., Pellerano, G. and Rosa, G.** (1997). Salinity-induced changes in the fine structure of the gills of the semiterrestrial estuarine crab, *Uca uruguayensis* (Nobili, 1901) (Decapoda, Ocypodidae). *Tissue Cell* **29**, 495–501.
- Mantel, L. H. and Farmer, L. L.** (1983). Osmotic and ionic regulation. In *The Biology of Crustacea*, vol. 5 (ed. L. H. Mantel), pp. 53–161. London, New York: Academic Press.
- Martelo, M. J. and Zanders, I. P.** (1986). Modifications of gill ultrastructure and ionic composition in the crab *Goniopsis cruentata* acclimated to various salinities. *Comp. Biochem. Physiol.* **84A**, 383–389.
- Martinez, C. B. R., Harris, R. R. and Santos, M. C. F.** (1998). Transepithelial potential differences and sodium fluxes in isolated perfused gills of the mangrove crab *Ucides cordatus*. *Comp. Biochem. Physiol.* **120A**, 227–236.
- Mougabure Cueto, G. A.** (1998). Interacción entre la regulación iónica y el equilibrio ácido-base en el cangrejo *Chasmagnathus granulata* Dana, 1851 (Decapoda, Grapsidae). pp. 1–55. Universidad de Buenos Aires. Graduate Thesis.
- Onken, H.** (1996). Active and electrogenic absorption of Na<sup>+</sup> and Cl<sup>-</sup> across posterior gills of *Eriocheir sinensis*: influence of short-term osmotic variations. *J. Exp. Biol.* **199**, 901–910.
- Onken, H. and Riestenpatt, S.** (1998). NaCl absorption across split gill lamellae of hyperregulating crabs: Transport mechanisms and their regulation. *Comp. Biochem. Physiol.* **119A**, 883–893.
- Onken, H. and Siebers, D.** (1992). Voltage-clamp measurements on single split lamellae of posterior gills of the shore crab *Carcinus maenas*. *Mar. Biol.* **114**, 385–390.
- Péqueux, A.** (1995). Osmotic regulation in crustaceans. *J. Crust. Biol.* **15**, 1–60.
- Péqueux, A., Gilles, R. and Marshall, W. S.** (1988). NaCl transport in gills and related structures. In *Advances in Comparative and Environmental Physiology*, vol. 1 (ed. R. Greger), pp. 1–73. Berlin: Springer.
- Pierrot C., Péqueux, A. and Thuet, P.** (1995a). Perfusion of gills isolated from the hyper-hyporegulating crab *Pachygrapsus marmoratus* (Crustacea, Decapoda): adaptation of a method. *Arch. Physiol. Biochem.* **103**, 401–409.
- Pierrot, C., Péqueux, A. and Thuet, P.** (1995b). Effects of ion substitution and of inhibitors on transepithelial potential difference and sodium fluxes in perfused gills of the crab *Pachygrapsus marmoratus*. *Arch. Physiol. Biochem.* **103**, 466–475.
- Riddle, C. V. and Ernst, S. A.** (1979). Structural simplicity of the *zonula occludens* in the electrolyte secreting epithelium of the avian salt gland. *J. Membr. Biol.* **45**, 21–35.
- Riestenpatt, S., Onken, H. and Siebers, D.** (1996). Active absorption of Na<sup>+</sup> and Cl<sup>-</sup> across the gill epithelium of the shore crab *Carcinus maenas*: voltage-clamp and ion-flux studies. *J. Exp. Biol.* **199**, 1545–1554.
- Rosa, G. A., Genovese, G., Luquet, C. M., Ansaldo, M. and Pellerano, G. N.** (1999). Histophysiological changes in *Chasmagnathus granulatus* during acclimation from low to high salinity. *Comp. Biochem. Physiol.* **124A**, S142. (Abstract)
- Schmidt, A. S. C. and Santos, E. A.** (1993). Behavior and haemolymphatic ionic composition of the intertidal crab *Chasmagnathus granulata* Dana, 1851 (Crustacea: Decapoda) during emersion. *Comp. Biochem. Physiol.* **106A**, 337–342.
- Schubart, C. D. and Diesel, R.** (1998). Osmoregulatory capacities and penetration into terrestrial habitats: A comparative study of jamaican crabs of the genus *Armases* Abele, 1992 (Brachyura: Grapsidae: Sesarminae). *Bull. Mar. Sci.* **62**, 743–752.
- Schubart, C. D. and Diesel, R.** (1999). Osmoregulation and the transition from marine to freshwater and terrestrial life: A comparative study of jamaican crabs of the genus *Sesarma*. *Arch. Hydrobiol.* **145**, 331–347.
- Siebers, D., Leweck, K., Markus, H. and Winkler, A.** (1982). Sodium regulation in the shore crab *Carcinus maenas* as related to ambient salinity. *Mar. Biol.* **69**, 37–43.
- Siebers, D., Winkler, A., Lucu, C., Thedens, G. and Weichart, D.** (1985). Na-K-ATPase generates an active transport potential in the gills of the hyperregulating shore crab *Carcinus maenas*. *Mar. Biol.* **87**, 185–192.
- Skou, J. C.** (1965). Enzymatic basis for active transport of Na<sup>+</sup> and K<sup>+</sup> across the cell membrane. *Physiol. Rev.* **45**, 596–617.
- Sokal, R. R. and Rohlf, F. J.** (1981). *Biometry*. 859pp. New York: W. H. Freeman.
- Towle, D. W. and Kays, W. T.** (1986). Basolateral localization of Na<sup>+</sup> K<sup>+</sup>-ATPase in gill epithelium of two osmoregulating crabs, *Callinectes sapidus* and *Carcinus maenas*. *J. Exp. Zool.* **239**, 311–318.
- Ussing, H. H.** (1949). The distinction by means of tracers between active transport and diffusion. *Acta Physiol. Scand.* **19**, 43–56
- Zeiske, W.** (1990). Noise from apical potassium ion channels. *Curr. Top. Membr. Transp.* **37**, 157–190.