

Limited extracellular but complete intracellular acid–base regulation during short-term environmental hypercapnia in the armoured catfish, *Liposarcus pardalis*

C. J. Brauner^{1,*}, T. Wang², Y. Wang³, J. G. Richards¹, R. J. Gonzalez⁴, N. J. Bernier⁵, W. Xi⁴, M. Patrick³ and A. L. Val⁶

¹Department of Zoology, University of British Columbia, 6270 University Boulevard, Vancouver, B.C., Canada V6T 1Z4, ²Department of Zoophysiology, University of Aarhus, 8000 Aarhus C, Denmark, ³Department of Biology, Queens University, Kingston, Ontario, Canada K7L 3N6, ⁴Department of Biology, University of San Diego, San Diego, CA 92110, USA, ⁵Department of Zoology, University of Guelph, Guelph, ON, Canada N1G 2W1 and ⁶National Institute for Research in the Amazon (INPA), Laboratory of Ecophysiology and Molecular Evolution, Manaus, AM, Brazil, CEP 69083-000

*Author for correspondence (e-mail: brauner@zoology.ubc.ca)

Accepted 17 June 2004

Summary

Environmental hypercapnia induces a respiratory acidosis that is usually compensated within 24–96 h in freshwater fish. Water ionic composition has a large influence on both the rate and degree of pH recovery during hypercapnia. Waters of the Amazon are characteristically dilute in ions, which may have consequences for acid–base regulation during environmental hypercapnia in endemic fishes. The armoured catfish *Liposarcus pardalis*, from the Amazon, was exposed to a water P_{CO_2} of 7, 14 or 42 mmHg in soft water (in $\mu\text{mol l}^{-1}$: Na^+ , 15, Cl^- , 16, K^+ , 9, Ca^{2+} , 9, Mg^{2+} , 2). Blood pH fell within 2 h from a normocapnic value of 7.90 ± 0.03 to 7.56 ± 0.04 , 7.34 ± 0.05 and 6.99 ± 0.02 , respectively. Only minor extracellular pH (pH_e) recovery was observed in the subsequent 24–96 h. Despite the pronounced extracellular acidosis, intracellular pH (pH_i) of the heart, liver and white muscle was tightly regulated within 6 h (the earliest time at which these parameters were measured) via a rapid accumulation of intracellular

HCO_3^- . While most fish regulate pH_i during exposure to environmental hypercapnia, the time course for this is usually similar to that for pH_e regulation. The degree of extracellular acidosis tolerated by *L. pardalis*, and the ability to regulate pH_i in the face of an extracellular acidosis, are the greatest reported to date in a teleost fish. The preferential regulation of pH_i in the face of a largely uncompensated extracellular acidosis in *L. pardalis* is rare among vertebrates, and it is not known whether this is associated with the ability to air-breathe and tolerate aerial exposure, or living in water dilute in counter ions, or with other environmental or evolutionary selective pressures. The ubiquity of this strategy among Amazonian fishes and the mechanisms employed by *L. pardalis* are clearly worthy of further study.

Key words: acid–base regulation, hypercapnia, pH_e , pH_i , intracellular pH regulation, catfish, *Liposarcus pardalis*, Amazon, water hardness, air-breathing.

Introduction

Environmental hypercapnia is common in freshwater systems with the most extreme examples being tropical aquatic environments covered with dense mats of vegetation, where CO_2 tensions can rise to as high as 60 mmHg (1 mmHg = approx. 0.133 kPa; Heisler et al., 1982). Hypercapnia causes a respiratory acidosis that, in most fish, is corrected within 10–72 h through branchial transfer of acid–base relevant ions (Heisler, 1984), while the kidney plays a relatively minor role (Perry et al., 1987). As a result, plasma HCO_3^- levels are elevated and plasma Cl^- levels reduced, implicating gill $\text{Cl}^-/\text{HCO}_3^-$ exchange as the primary mechanism for extracellular pH (pH_e) compensation. In general, intracellular

pH (pH_i) compensation follows a qualitatively similar temporal pattern (Heisler, 1984; Cameron and Kormanik, 1982).

Complete or significant degrees of pH compensation during environmental hypercapnia have been demonstrated in numerous species of freshwater and marine fish (e.g. Toews et al., 1983; Claiborne and Heisler, 1984; Cameron, 1985; Goss et al., 1995; Larsen and Jensen, 1997; Larsen et al., 1997; reviewed by Heisler, 1984; Claiborne et al., 2002). Water composition clearly affects the rate of acid–base compensation where the rate and degree of compensation is more pronounced in seawater than in freshwater, and more pronounced in hard

compared to soft freshwater (Heisler, 1984; Larsen and Jensen, 1997).

Many natural systems are characterized by soft water. One example is the Amazon, where the ionic composition of many rivers and tributaries approaches that of distilled water (Val and Almeida-Val, 1995). Given that extensive vegetation covers many of the stagnant lakes and ponds in the Amazon during the dry season, it is likely that resident fishes experience periods of hypercapnia (Heisler et al., 1982). The ability of fishes endemic to these ion-poor waters to tolerate and compensate for environmental hypercapnia has not previously been investigated. The armoured catfish *Liposarcus pardalis* is common in the Amazon and can withstand relatively severe environmental hypercapnia and large disturbances in plasma pH (Randall et al., 1996). Here, we investigate the temporal regulation of extra- and intracellular acid–base status during environmental hypercapnia of *L. pardalis* in natural softwater. The first experimental series describes extracellular acid–base status during hypercapnia (P_{CO_2} of 7, 14 and 42 mmHg) in softwater, while the second series investigates intracellular acid–base parameters of heart, liver and muscle during hypercapnia (P_{CO_2} of 14 and 32 mmHg). The third experimental series investigates whole animal fluxes of Na^+ , Cl^- and ammonia during hypercapnia (P_{CO_2} of 14 and 32 mmHg) to elucidate the mechanisms for acid–base regulation.

Materials and methods

Animal acquisition and holding

Liposarcus pardalis (Castelnaud 1855) (308±15.5 g) were obtained from a commercial fish supplier and held at the National Institute for Research of the Amazon (INPA), Manaus, Brazil, for at least 2 weeks prior to experimentation. Fish were maintained in aerated well-water in outdoor tanks on a natural photoperiod. The ionic composition (in $\mu\text{mol l}^{-1}$) was Na^+ , 15; Cl^- , 16; K^+ , 9; Ca^{2+} , 9; Mg^{2+} , 2. Water temperature was 28°C, and water pH was 6.9±0.3. Fish were fed commercially purchased pellets on a maintenance ration of 1% body mass per day, but feeding was withheld 48–72 h prior to experimentation.

Experimental protocols

Series I: Effect of hypercapnia on extracellular acid–base status

Fish were anaesthetized in 0.1 g l^{-1} Tricaine Metanesulfonate (MS-222) buffered with NaHCO_3^- . Once anaesthetized, fish were transferred to a surgery table, and an indwelling catheter (PE-50) was placed in the dorsal aorta according to Soivio et al. (1975). Fish were left to recover for 24–48 h in individual chambers, and the cannulae were flushed several times with heparinized saline. Following recovery, the fish were placed in 2 l chambers without access to air and exposed to a P_{CO_2} of 7, 14 or 42 mmHg. The low levels were achieved using a flow meter that mixed CO_2 with air into a 500 l recirculating system, while the high level was achieved using flow-through well-water directly from its source without

prior aeration. Water CO_2 levels stabilized within 1 h. Blood samples were withdrawn from the cannulae at 0, 2, 6 and 24 h at each level of hypercapnia. Whole blood pH (pH_e), P_{O_2} , true plasma total CO_2 content, haemoglobin concentration ($[\text{Hb}]$), and haematocrit (Hct) were measured immediately, and plasma was frozen for later analysis of Na^+ , Cl^- and protein concentrations (see Analytical procedures below). Cannulated fish exposed to 42 mmHg did not survive for 24 h. Thus, non-cannulated fish were exposed to 42 mmHg for 96 h, and a blood sample was obtained from the caudal vein within 10 s of removal from the water.

Series II: Effect of hypercapnia on intracellular acid–base status

Non-cannulated fish were placed in individual 2 l chambers overnight before being exposed to a P_{CO_2} of 14 or 32 mmHg without access to air. Tissues were sampled from normocapnic control fish and at 6, 24 or 72 h into the hypercapnic period. For tissue sampling, water flow to the chamber was stopped and a concentrated solution of buffered MS-222 was slowly added to the water to achieve a final concentration of 0.1 g l^{-1} , according to Wang et al. (1994). Within 2–3 min, fish lost equilibrium and could be removed from the water without struggling. This procedure has been shown to minimize any metabolic and acid–base changes associated with handling and sampling. Blood (0.5 ml) was drawn from the caudal vein into a heparinized syringe for measurement of pH_e , red cell intracellular pH (pH_i), P_{CO_2} , $[\text{Hb}]$, Hct and plasma Na^+ , Cl^- , Ca^{2+} concentrations, and osmolarity. The fish were then killed, and 0.5 g samples from heart, liver and muscle were removed and frozen immediately in liquid nitrogen for later analysis of pH_i . Sampling was complete within 2–3 min after the fish had been removed from the water.

Series III: Effect of hypercapnia on whole animal ion fluxes

Non-cannulated fish were placed in individual chambers (500 ml) overnight and exposed to the same regime as fishes of Series II. Fish were subjected to a P_{CO_2} of 14 or 32 mmHg. At 0, 6 and 24 h (and at 72 h at P_{CO_2} of 32 mmHg), water flow was interrupted for 1 h for measurements of whole animal unidirectional Na^+ influx ($J_{\text{in}}^{\text{Na}}$), efflux ($J_{\text{out}}^{\text{Na}}$), net flux ($J_{\text{net}}^{\text{Na}}$), net Cl^- flux ($J_{\text{net}}^{\text{Cl}}$), and total ammonia excretion.

Analytical procedures

Blood P_{O_2} and P_{CO_2} values were measured using Radiometer (Copenhagen, Denmark) P_{O_2} (E5046) and P_{CO_2} (E5036) electrodes thermostatted at 28°C in a BMS Mk2 electrode assembly with the output displayed on a Radiometer PHM 73. Total CO_2 content of true plasma was measured according to Cameron (1971). Extra- and intracellular pH were measured using a Radiometer microcapillary electrode (G299A) held within the BMS Mk2 system. Red cell pH_i was measured using the freeze–thaw method of Zeidler and Kim (1977), and pH_i of heart, liver and white muscle was measured according to Pörtner et al. (1990). Water pH was measured using a Corning (Corning, USA) combination pH electrode.

Haematocrit was measured in duplicate after centrifuging blood in micro-haematocrit tubes at 12 000 r.p.m. for 3 min. Blood [Hb] was measured spectrophotometrically following conversion to cyanomethaemoglobin, applying a millimolar extinction coefficient of 11.0. Plasma Na^+ and Ca^{2+} levels were measured using atomic absorption flame photometry, and plasma Cl^- concentration was measured according to Zall et al. (1956). Plasma ammonia, urea, glucose and lactate were measured using Sigma (St Louis, USA) diagnostic kits, and gill homogenate Na^+, K^+ -ATPase activity was measured according to McCormick (1993) and expressed relative to total homogenate protein which was measured using Bradford reagent (Bio-Rad, Richmond, CA, USA) and bovine serum albumin as a standard.

$J_{\text{in}}^{\text{Na}}$ was measured by adding 1.5 μCi (56 kBq) ^{22}Na to each experimental chamber and taking a 5 ml water sample at the respective time 0 (i.e. 0, 2, 6, 24 or 72 h after initiating exposure to hypercapnia) and 2 h following addition of isotope. Water samples were measured for radioactivity using a Beckman (Fullerton, USA) Coulter LS 6500 (Multi-purpose scintillation counter) for calculation of J_{in} as indicated below. Water Na^+ (measured spectrophotometrically), Cl^- (measured according to Zall et al., 1956), and total ammonia (measured using a Sigma diagnostic kit) were determined for calculation of net fluxes based upon the difference of the respective parameter over the 2 h incubation duration, taking into account the weight of individual fish.

Calculations

Plasma $[\text{HCO}_3^-]$ was calculated from the measured blood pH and P_{CO_2} (Series II). In series I, plasma $[\text{HCO}_3^-]$ was calculated from pH and total CO_2 concentration of plasma, using the Henderson–Hasselbalch equation. The CO_2 solubility coefficient and pK' for plasma were taken from Boutilier et al. (1984). Intracellular $[\text{HCO}_3^-]$ for heart, liver and muscle were calculated from the pH_i of the respective tissue, and the CO_2 solubility coefficient and pK' for plasma as described by (Heisler, 1982; Heisler et al., 1982).

$J_{\text{in}}^{\text{Na}}$ was calculated from the disappearance of isotope from the water and the average Na^+ concentration of the water during the flux period using the equation from Gonzalez and Dunson (1987):

$$J_{\text{in}} = \frac{(\ln Q_{\text{out}0} - \ln Q_{\text{out}1})Q_{\text{out}}}{(Mt)}, \quad (1)$$

where $Q_{\text{out}0}$ and $Q_{\text{out}1}$ are the total counts per minute (c.p.m.) in the flux chambers at the beginning and end of the flux period, respectively. Q_{out} is the average amount of Na^+ in the flux bath during the flux period, M is the mass of the fish in g, and t is the time in h.

Statistics

Differences among mean values within a given P_{CO_2} treatment were determined using a one-way ANOVA, or a one-way repeated-measures ANOVA as appropriate, followed by

a Tukey's *post-hoc* test. Data are presented as mean \pm S.E.M., $N=6$, unless otherwise indicated.

Results

Series I: Effect of hypercapnia on extracellular acid–base status

Blood P_{CO_2} remained elevated throughout the exposures to hypercapnia (Fig. 1A). When compared to normocapnia, extracellular pH was significantly reduced throughout all hypercapnic exposures. However, at a P_{CO_2} of 14 mmHg, pH_e increased significantly at 24 h compared to the initial and maximal reduction at 2 h, while pH_e increased significantly at 96 h compared to 2 h during exposure to a P_{CO_2} of 42 mmHg (Fig. 1B). Hypercapnia was associated with a progressive elevation of plasma HCO_3^- that was more marked at the more severe exposure (Fig. 1C). When expressed in a $\text{pH}-\text{HCO}_3^-$




Fig. 1. The effect of exposure to a water P_{CO_2} of 7 mmHg (circles), 14 mmHg (triangles) and 42 mmHg (squares) on (A) blood P_{CO_2} , (B) blood pH (pH_e) and (C) plasma HCO_3^- in *L. pardalis*. Within each P_{CO_2} treatment, lower case letters indicate values that differ significantly from each other.




Fig. 2. pH-HCO₃⁻ plot of blood from *L. pardalis* exposed to a P_{CO}₂ of 7 (circles), 14 (triangles) and 42 (squares) mmHg for up to 96 h. The broken line represents the *in vitro* buffer line for rainbow trout whole blood (Wood et al., 1982), and curved lines indicate P_{CO}₂ isopleths.

diagram (Fig. 2), and assuming that the whole blood non-bicarbonate buffer line of *L. pardalis* is similar to trout, it is apparent that half of the HCO₃⁻ accumulation in plasma can be ascribed to non-bicarbonate buffering (Fig. 2). pH_e did not recover at any level of hypercapnia. When expressed relative to the difference between pH_e in normocapnia and that after 2 h, pH_e only recovered by 8% following 24 h exposure to 7 mmHg, 32% following 24 h at 14 mmHg, and 23% following 96 h exposure to 42 mmHg.

Hypercapnia did not affect arterial P_O₂ (pooled value=45.4±2.7 mmHg), mean cell haemoglobin concentration (MCHC) (pooled value=3.9±0.1), plasma [Na⁺] (pooled value=140.5±3.0 meq l⁻¹) or [Cl⁻] (pooled value=114.6±1.1 meq l⁻¹). There was, however, a statistically significant reduction in Hct (from 40.4±1.6 at time 0, to 30.9±3.4 at time 24 h) which, most

likely, relates to blood sampling. There were no significant changes in Hct in Series II.

Series II: Effect of hypercapnia on intracellular acid-base status

When non-cannulated fish were exposed to hypercapnia, blood was withdrawn from the caudal vein following terminal anaesthesia to prevent struggling and associated acidosis. Nevertheless, pH_e of normocapnic fish was lower than pH_e of cannulated fish in Series I (Figs 1 and 3). Exposure to 14 mmHg significantly reduced pH_e and red cell pH_i at 6 and 24 h with no sign of pH compensation over 24h (Fig. 3). There were however, no significant changes in pH_i of heart, liver or white muscle (Fig. 3). When exposed to 32 mmHg, *L. pardalis* also experienced a substantial, and largely uncompensated, reduction in pH_e and red cell pH_i that persisted for up to 72 h (Fig. 4). Again, however, there were no statistically significant reductions in pH_i of heart, liver or white muscle during the 72 h of hypercapnia. Intracellular pH regulation of heart, liver and white muscle during the extracellular acidosis is evident when pH_i is plotted against pH_e for all time periods at both levels of hypercapnia (Fig. 5). Clearly, red cell pH decreases with lowered pH_e, while pH_i of liver heart and skeletal muscle tend to increase.

As in series I, plasma HCO₃⁻ increased during hypercapnia




Fig. 3. The effect of exposure to a water P_{CO}₂ of 14 mmHg on whole blood pH (open circles) and intracellular pH of red blood cells (closed circles), heart (triangles), liver (diamonds) and muscle (squares) in *L. pardalis*. Letters that differ within a tissue differ significantly from one another. No significant differences were observed at any time in heart, liver and muscle.




Fig. 4. The effect of exposure to a water P_{CO}₂ of 32 mmHg on whole blood pH (open circles) and intracellular pH of red blood cells (closed circles), heart (triangles), liver (diamonds) and muscle (squares) in *L. pardalis*. Letters that differ within a tissue differ significantly from one another. No significant differences were observed at any time in heart, liver and muscle.




Fig. 5. The relationship between tissue intracellular pH (pHi) and whole blood pH (pHe) in *L. pardalis* exposed to a water P_{CO_2} of 14 and 32 mmHg for up to 24 h. Circles represent red blood cells ($y=3.516+0.4546x$, $r^2=0.964$), triangles represent heart ($y=7.692-0.132x$, $r^2=0.211$), diamonds represent liver ($y=7.618-0.111x$, $r^2=0.247$) and squares represent white muscle ($y=7.388-0.045x$, $r^2=0.045$).

(Table 1). There were no significant changes in plasma $[Na^+]$, $[Cl^-]$ or $[Ca^{2+}]$; however, there was a trend toward a reduction in plasma $[Cl^-]$, consistent with the changes in plasma HCO_3^- . There was no significant effect of hypercapnia on haematocrit (pooled value= 27.9 ± 1.5), mixed-venous P_{O_2} (pooled value= 19.4 ± 1.6 mmHg), plasma urea (pooled value= 513 ± 56 $\mu\text{mol l}^{-1}$ nitrogen), plasma lactate (pooled value= 0.41 ± 0.13 mmol l^{-1}), plasma glucose (pooled value= 3.0 ± 0.2 mmol l^{-1}) or gill Na^+,K^+ -ATPase activity (pooled value= 0.82 ± 0.32 mmol $\text{l}^{-1} \text{g}^{-1} \text{h}^{-1}$). There was a significant increase in plasma ammonia levels at both levels of hypercapnia (Table 1). Calculated intracellular $[HCO_3^-]$ for heart, liver and kidney were significantly elevated at 6 h in both levels of hypercapnia. There were no significant differences after 6 h, indicating complete intracellular acid–base regulation at this time (Table 1).

While most of the intracellular HCO_3^- accumulation during hypercapnia is due to non-bicarbonate buffering, additional HCO_3^- accumulation must have occurred to compensate for the acidosis. Assuming a non-bicarbonate buffer value of 45 slykes for white muscle, as determined for the facultative air-breather, *Synbranchus marmoratus* (Heisler, 1982), it can be estimated that intracellular $[HCO_3^-]$, in the absence of intracellular pH compensation, would have risen to approximately 5.5 mmol l^{-1} at a P_{aCO_2} of 14 mmHg (A in Fig. 6). When compared to the calculated intracellular value of 6.8 mmol l^{-1} (Table 1), 1.3 mmol l^{-1} HCO_3^- must have accumulated in the white muscle within 6 h to alleviate the acidosis. At a P_{aCO_2} of 32 mmHg, intracellular $[HCO_3^-]$ would be expected to increase to approximately 9.2 mmol l^{-1} in the absence of pHi compensation (B in Fig. 6). When subtracted from the calculated intracellular value of 13.0 mmol l^{-1} (Table 1), it appears that 3.8 mmol l^{-1} HCO_3^- accumulated in white muscle to restore pHi.

Table 1. Series II plasma parameters and calculated intracellular $[HCO_3^-]$ in heart, liver and white muscle in fish exposed to a water P_{CO_2} of 14 or 32 mmHg for the indicated duration

Time (h)	Plasma parameters						Tissue $[HCO_3^-]$ (mmol l^{-1})			
	P_{CO_2} (mmHg)	Osmolarity (mOsm l^{-1})	$[Na^+]$ ($\mu\text{eq l}^{-1}$)	$[Cl^-]$ ($\mu\text{eq l}^{-1}$)	$[Ca^{2+}]$ ($\mu\text{eq l}^{-1}$)	HCO_3^- (mmol l^{-1})	[Ammonia] ($\mu\text{mol l}^{-1}$)	Heart	Liver	Muscle
14 mmHg										
0	5.5±0.3 ^a	238.0±3.5	135.4±5.3	111.8±3.4	3.6±0.2	9.8±0.6 ^a	209.7±25.3 ^a	0.7±0.1 ^a	1.0±0.1 ^a	2.2±0.3 ^a
6	16.9±0.8 ^b	239.2±2.8	130.0±5.0	107.1±3.2	3.6±0.1	15.3±0.6 ^b	455.4±29.7 ^b	3.2±0.4 ^b	4.0±0.7 ^b	6.8±0.8 ^b
24	14.9±0.3 ^c	230.2±5.0	131.4±2.4	106.5±2.3	3.6±0.1	15.3±0.6 ^b	412.8±44.9 ^b	3.1±0.3 ^b	2.7±0.2 ^b	5.2±0.3 ^b
32 mmHg										
0	4.5±0.2 ^a	247.7±6.6	135.9±6.0	111.1±4.7	3.4±0.2	8.8±0.3 ^a	422.2±28.1 ^{a,b}	0.7±0.1 ^a	0.9±0.1 ^a	1.9±0.2 ^a
6	33.4±1.2 ^b	246.8±5.2	134.8±2.8	108.4±2.5	4.0±0.5	20.6±1.7 ^b	693.2±83.4 ^{a,b}	5.4±0.7 ^b	7.0±0.8 ^b	13.0±1.0 ^b
24	31.9±1.3 ^b	242.3±2.3	127.7±2.9	104.2±1.9	3.4±0.3	25.9±3.1 ^b	768.7±127.1 ^{a,b}	5.6±0.4 ^b	7.3±0.3 ^b	13.3±1.0 ^b
72	31.9±1.3 ^b	238.3±6.8	135.1±5.4	102.7±6.8	3.8±0.1	22.7±1.9 ^b	636.2±49.3 ^b	6.0±0.7 ^b	6.7±0.9 ^b	15.9±0.9 ^b

Different superscript letters indicate values significantly different from each other for each parameter within each P_{CO_2} treatment.

Series III: Effect of hypercapnia on whole animal ion fluxes

Fish were exposed to hypercapnia simultaneously with those of Series II and thus experienced identical levels of hypercapnia. At a P_{CO_2} of 14 mmHg, fish exhibited about a 70% reduction in J_{in}^{Na} in conjunction with a large reduction in J_{out}^{Na} . As a result J_{net}^{Na} did not change at 6 and 24 h (Fig. 7). There was a significant decrease in J_{net}^{Cl} , but no significant effect of hypercapnia exposure on total ammonia excretion (pooled value of $396 \pm 28 \text{ nmol g}^{-1} \text{ h}^{-1}$). Urea excretion rates prior to hypercapnia were low ($11.5 \pm 1.0 \text{ nmol g}^{-1} \text{ h}^{-1}$) representing a total of $8.0 \pm 0.9\%$ of total nitrogenous waste excreted. Urea excretion was not measured during hypercapnia.

The effects of exposure to 32 mmHg were qualitatively similar with large reductions in both J_{in}^{Na} and J_{out}^{Na} , resulting in no significant change in J_{net}^{Na} over 72 h (Fig. 8). A significant reduction in J_{net}^{Cl} , relative to normocapnic control fish, was observed during exposure to hypercapnia. However, after 6 h, J_{net}^{Cl} was near 0.

Discussion

When *L. pardalis* were exposed to environmental hypercapnia, arterial pH was reduced in proportion to the rise in P_{aCO_2} , with little subsequent accumulation of plasma HCO_3^- and almost no compensation of pH_e over the following 24–96 h. In spite of the pronounced extracellular acidosis (reaching a level as low as 6.99), pH_i of the heart, liver and white muscle was tightly regulated within 6 h (the earliest time at which these parameters were measured). While most fish regulate pH_i during exposure to environmental hypercapnia, the time course for this is often similar to that for pH_e regulation (Heisler, 1984). The degree of extracellular acidosis tolerated by *L. pardalis*, and the ability to regulate pH_i in the face of an extracellular acidosis, are among the greatest reported to date in a teleost fish.

Extracellular acid–base regulation

The magnitude of the respiratory acidosis tolerated by *L. pardalis* is among the greatest reported in the literature. Blood pH fell from 7.98 to 6.99 within 2 h of exposure to a water P_{CO_2} of 42 mmHg. Eel (*Anguilla anguilla*) is also remarkably tolerant to hypercapnia, and blood pH fell from 7.92 to 7.16 as P_{CO_2} increased from 3 to 45 mmHg over a 3 h period




Fig. 6. pH- HCO_3^- plot of white muscle intracellular fluid of *L. pardalis* exposed to a P_{CO_2} of 14 and 32 mmHg for 6 h. The dotted line represents the *in vitro* buffer line for the white muscle of *Synbranchus marmoratus* (Heisler, 1982); curved lines indicate P_{CO_2} isopleths. A and B indicate the predicted HCO_3^- concentrations in white muscle in the absence of pH compensation at a water P_{CO_2} of 14 and 42 mmHg, respectively.




Fig. 7. Unidirectional Na^+ uptake (J_{in}) or loss (J_{out}) and net (J_{net}) Na^+ and Cl^- flux during exposure to a water P_{CO_2} of 14 mmHg for 24 h in *L. pardalis*. Letters that differ within a given parameter differ significantly from one another.

(McKenzie et al., 2002), resulting in no mortality with continued exposure to hypercapnia (McKenzie et al., 2003). Following 6 weeks exposure to a P_{CO_2} of 45 mmHg, there was considerable HCO_3^- accumulation (up to 72 mmol l^{-1}) associated with a large fall in plasma Cl^- in the eel (McKenzie et al., 2003); however, restoration of pH_e did not exceed 50%. White sturgeon *Acipenser transmontanus* also tolerate a




Fig. 8. Unidirectional Na^+ uptake (J_{in}) or loss (J_{out}) and net (J_{net}) Na^+ and Cl^- flux during exposure to a water P_{CO_2} of 32 mmHg for 72 h in *L. pardalis*. Letters that differ within a given parameter differ significantly from one another.

reduction in pH_e from 7.7 to 7.15 when exposed to severe hypercapnia where arterial P_{CO_2} reaches 30 mmHg (Crocker and Cech, 1998).

Most fish completely or partially restore pH_e , through net accumulation of plasma HCO_3^- in exchange for Cl^- , within 10 to 72 h of being exposed to hypercapnia. Sturgeon, however, show a very blunted recovery of pH during hypercapnia, where exposure to a P_{CO_2} of 30 mmHg resulted in no pH_e compensation at 24 h, and a mere 35% restoration of pH_e following 72 h (Crocker and Cech, 1998). The water used in this study was relatively hard, which in many species facilitates acid–base regulation during hypercapnia (Heisler, 1984; Larsen and Jensen, 1997). The degree of pH compensation observed in *L. pardalis* during hypercapnia was similar to sturgeon. 24 h after exposure to 7 mmHg, pH_e had only recovered by 8% and 96 h exposure to 42 mmHg merely led to a 23% recovery of pH_e . Hypercapnia was associated with a small reduction in J_{net}^{Cl} , indicating that branchial $\text{Cl}^-/\text{HCO}_3^-$ exchange was involved in the small rise of intra- and extracellular HCO_3^- . There was also a significant reduction of J_{in}^{Na} , which may be a direct effect of the low water pH (Wood, 1989). When *L. pardalis* were exposed to a P_{CO_2} of 10 mmHg in the presence of a 60- to 70-fold elevation in water NaCl concentration (1.0 mmol l⁻¹ NaCl), there was no change in the rate or magnitude of acid–base recovery within 24 h (data not shown), indicating that the blunted extracellular response to hypercapnia may not be solely due to limitations associated with ionic composition of the water.

L. pardalis can tolerate a water P_{CO_2} of 42 mmHg for weeks in water with low ionic levels, but it is not known whether they do compensate extracellular acid–base status over this duration. Interestingly, the ability of *L. pardalis* to tolerate extreme hypercapnia was size-dependent; in non-cannulated

fish, all fish greater than 150 g survived, while all those weighing between 50–80 g died. Cannulation also resulted in higher mortality. Fish reduced spontaneous activity and remained stationary for days during exposure to severe hypercapnia. Although *L. pardalis* is a facultative air-breather, they did not attempt to air-breathe even when they had access to air (C.J.B., unpublished observation). It has been proposed that the exceptional tolerance to environmental hypercapnia observed in the eel is associated with the need to tolerate air exposure (Heisler, 1982; McKenzie et al., 2002). This may also be the case for *L. pardalis*, which can survive aerial exposure for days (A. L. Val, unpublished data).

A complete lack of extracellular pH regulation during the respiratory acidosis that develops upon transition to air-breathing was observed in the facultative air-breather, *Synbranchus marmoratus* (Heisler, 1982). When exposed to aquatic hypoxia for 4 days, *S. marmoratus* relied exclusively upon air-breathing which, in combination with the reduction of water flow across the gills, led to a

fivefold increase of P_{aCO_2} and a reduction in pH_e from about 8.15 to 7.5. Plasma HCO_3^- did not increase to compensate for the acidosis, and there was very limited $[\text{HCO}_3^-]$ uptake from the water. This lack of pH_e compensation was concluded to result from the lack of gill–water contact and the associated impairment of branchial transfer of acid–base relevant ions (Heisler, 1982). It remains, however, to be established whether exposure to hypercapnic water, where gill–water interaction is maintained, would be associated with pH_e compensation in *S. marmoratus*. *Liposarcus pardalis* is also a facultative air-breather and it is possible that its limited compensation of pH_e during hypercapnia is associated with this trait rather than with the low ionic content of the water in which the fish lives. However, not all facultative air-breathers that experience a respiratory acidosis upon air-breathing exhibit a lack of pH compensation as seen in *S. marmoratus*. In another Amazonian catfish, *Hypostomus* sp., air-breathing was associated with a large reduction in gill ventilation (and presumably gill–water interaction) and blood P_{CO_2} increased from 3 to 20 mmHg; however, pH_e was fully compensated after 4–7 days due to a 15 mmol l⁻¹ rise of plasma $[\text{HCO}_3^-]$, in spite of the low ionic content of the water (Wood et al., 1979).

Intracellular acid–base regulation

Regulation of intracellular pH is important for enzyme function and metabolism and pH_i is tightly regulated in most animals. Regulation of pH_i is generally associated with partial to complete pH_e compensation during an acid–base disturbance in most animals (Heisler, 1984). A reduction in pH_i from 6.95 to 6.57 decreases glycolysis by 71% in mammalian cardiac muscle (see Somero and White, 1985). Severe impairment of aerobic metabolism is also likely with relatively small reductions of pH_i (0.1–0.3 pH units), due to the pH

sensitivity of enzymes such as citrate synthase and pyruvate dehydrogenase (see Hazel et al., 1978). The effects of intra- and extracellular acidosis can be partially alleviated by a high tissue buffer value, achieved through an elevation in histidyl imidazole or phosphate groups (Somero and White, 1985); but pH_i compensation requires active regulation.

Many teleost fishes regulate red cell pH during extracellular acidosis through release of catecholamines that stimulate red cell Na^+/H^+ exchange. This acts to safeguard oxygen transport, due to the presence of extremely pH-sensitive haemoglobins. This response protects blood oxygen binding that would have been reduced through the Root effect (see Nikinmaa, 1990). The armoured catfish do not exhibit a Root effect and do not possess adrenergic stimulation of red cell Na^+/H^+ exchange (Val et al., 1998). In contrast to the other tissues studied in *L. pardalis*, red blood cell pH decreased throughout the hypercapnic exposures. The ratio of the $\Delta\text{pH}_i/\Delta\text{pH}_e$ was 0.45, which is similar to that of other tissues in animals that experience a large extracellular acidosis over a 24–48 h period, and similar to that observed in red blood cells of rainbow trout *in vitro* in the absence of catecholamines (Heming et al., 1986). Turtle hearts, *in vitro*, exhibit a $\Delta\text{pH}_i/\Delta\text{pH}_e$ of approximately 0.55 (Jackson et al., 1991) and whole body pH_i of the catfish *Ictalurus punctatus* is reduced with a $\Delta\text{pH}_i/\Delta\text{pH}_e$ of 0.60 during hypercapnia (Cameron, 1980), following which changes in pH_e and pH_i occurred roughly in parallel. In the skate exposed to a P_{CO_2} of 7 mmHg, $\Delta\text{pH}_i/\Delta\text{pH}_e$ values for brain, heart and white muscle are approximately 0.39, 0.45 and 0.76, respectively, over a 24 h period, where recovery of pH_i is tightly correlated with that of pH_e . During continued exposure to hypercapnia (days to weeks), pH_i has been reported to be completely regulated in aquatic animals; however, this requires some degree of extracellular compensation, usually approaching 50% (Claiborne and Heisler, 1986; Pörtner et al., 1998; McKenzie et al., 2003).

The only two vertebrates reported to date that regulate pH_i (of heart and white muscle) in the face of a large, uncompensated, extracellular acidosis are *S. marmoratus* (Heisler, 1982) and the salamander *Siren lacertina* (Heisler et al., 1982). In the salamander, an increase in water pH or bicarbonate infusion did not alter the acid–base strategy, indicating that the animals do not attempt to regulate pH_e , and preferentially regulate pH_i . In both *S. marmoratus* and salamanders, pH_i of the heart and muscle was tightly regulated within 4 days, the earliest time measured. In the heart, liver and white muscle of *L. pardalis*, there was a trend towards an increase in pH_i during the pronounced extracellular acidosis, which occurred as early as 6 h, indicating that active pH regulation must be involved. It is not known whether *S. marmoratus* or *S. lacertina* are capable of regulating pH_i this rapidly.

The rapid, preferential regulation of pH_i in the face of a largely uncompensated extracellular acidosis in *L. pardalis* is rare among vertebrates. It is not known whether this trait is associated with the ability to air-breathe and tolerate aerial exposure, with the low ionic content of the water, or with other

environmental or evolutionary selective pressures. It is conceivable that this strategy could be selected in order to minimize problems associated with Cl^- homeostasis in an aquatic environment such as the Amazon, which is dilute in ions (particularly Cl^- and HCO_3^-), acidic, and prone to daily oscillations in hypercapnia. Complete pH_e compensation during environmental hypercapnia is associated with large reductions in plasma Cl^- levels that would have to be replaced rapidly when the fish moves from a hypercapnic to normocapnic environment. The ubiquity of this strategy among Amazonian fishes preferentially to regulate pH_i and not regulate pH_e is clearly worthy of further study.

Mechanisms of intracellular pH regulation

Intracellular pH regulation is associated with a net HCO_3^- uptake into the tissue, which is accomplished by active membrane transport of strong ions (Reeves, 1985). The HCO_3^- accumulated may ultimately come from the water in association with gill $\text{HCO}_3^-/\text{Cl}^-$ or Na^+/H^+ exchange. The $J_{\text{net}}^{\text{Cl}}$ required to account for the intracellular HCO_3^- accumulation during hypercapnia can be calculated, but is only reported for white muscle here. White muscle was the single largest tissue measured, and because white muscle pH_i is higher than heart and liver, it is likely to represent a large proportion of the total intracellular HCO_3^- accumulated by *L. pardalis* during hypercapnia. Assuming that white muscle comprises about 50% of body mass in *L. pardalis* (lower than other teleosts because of the dense bone of the skull), and that intracellular water represents about 85% of the white muscle mass, the active accumulation of $1.3 \text{ mmol l}^{-1} \text{ HCO}_3^-$ (see Results, Section II), requires that approximately $0.55 \mu\text{mol HCO}_3^- \text{ g}^{-1}$ must have been accumulated within the white muscle 6 h after exposure to a water P_{CO_2} of 14 mmHg. Assuming that this HCO_3^- ultimately came from the water in exchange for Cl^- , then a $J_{\text{net}}^{\text{Cl}}$ of $-92 \text{ nmol g}^{-1} \text{ h}^{-1}$ would be required to regulate white muscle pH_i , which is very similar to the measured $J_{\text{net}}^{\text{Cl}}$ value of $-102 \text{ nmol g}^{-1} \text{ h}^{-1}$ (Fig. 7). Thus, based upon the fluxes measured in this study, and consistent with studies on other species (Heisler, 1984; Larsen and Jensen, 1997), branchial $\text{HCO}_3^-/\text{Cl}^-$ exchange is most likely to account for intracellular acid–base regulation under hypercapnia in *L. pardalis*.

Net $\text{Cl}^-/\text{HCO}_3^-$ exchange across the gills, however, does not appear to be sufficient to account for muscle HCO_3^- accumulation at a higher P_{CO_2} . If $3.8 \text{ mmol l}^{-1} \text{ HCO}_3^-$ was actively accumulated in the white muscle at 32 mmHg (Table 1), and given the assumptions above, $1.6 \mu\text{mol HCO}_3^-$ per gram of fish would have to be accumulated in the white muscle over the 6 h exposure duration, which would require a $J_{\text{net}}^{\text{Cl}}$ of $-267 \text{ nmol g}^{-1} \text{ h}^{-1}$. This is far greater than the highest rate of $-80 \text{ nmol g}^{-1} \text{ h}^{-1}$ measured at 6 h during this exposure, indicating that other pathways must be involved. Hypercapnia did not affect net Na^+ uptake or ammonia excretion and *L. pardalis* does not appear to excrete acid *via* the kidney (Randall et al., 1996). Thus, the remaining HCO_3^- taken up by the muscle (and other tissues) may have been shuttled from the

extra to intracellular space. A reduction in plasma $[\text{HCO}_3^-]$ below the blood buffer line was observed at 2 h, indicating that this may have occurred. A reduction in plasma $[\text{HCO}_3^-]$ and an increase in plasma $[\text{Cl}^-]$ were observed in *S. marmoratus* during initial exposure to hypercapnia, illustrating the preference of pH_i over pH_e regulation (Heisler, 1982). *Liposarcus pardalis* is endowed with a high bone mass, predominantly skull, and given the role of the shell and bone in compensating for an acidosis in turtles (Jackson, 1997; Jackson et al., 2000), bone demineralization may be playing a role in intracellular acid–base regulation in *L. pardalis*. Bone does not appear to be a route for acid–base compensation in the channel catfish, *Ictalurus punctatus*, exposed to hypercapnia (Cameron, 1985), and there were no statistically significant changes in plasma $[\text{Ca}^{2+}]$ in *L. pardalis*; however, Ca^{2+} efflux from the fish during hypercapnia was not measured. Whether or not bone demineralization is involved in acid–base regulation in *L. pardalis* remains to be investigated. Clearly pH_i of the heart, liver, white muscle, and probably other tissues, is tightly regulated in the face of a large, predominantly uncompensated extracellular acidosis in *L. pardalis* during hypercapnia. Elucidating the precise mechanisms involved remains an exciting avenue for further studies.

This study was supported by National Sciences and Engineering Research Council (NSERC) of Canada Discovery Grants to C.J.B., Y.W. and N.J.B., NSERC and Killam Post-doctoral fellowship to J.R., University Faculty fellowship to R.G. and CNPq Brazil research grant to A.L.V. T.W. was supported by The Danish Research Council. We thank Aline Matsuo for excellent technical assistance and Nazare Paula da Silva for her tireless efforts in this and many other studies.

References

- Boutilier, R. G., Heming, T. A. and Iwama, G. K. (1984). Appendix: Physicochemical parameters for use in fish respiratory physiology. In *Fish Physiology*, vol. XA (ed. W. S. Hoar and D. J. Randall), pp. 403–430. New York: Academic Press.
- Cameron, J. N. (1971). A rapid method for determination of total carbon dioxide in small blood samples. *J. Appl. Physiol.* **31**, 632–634.
- Cameron, J. N. (1980). Body fluid pools, kidney function and acid–base regulation in the freshwater catfish, *Ictalurus punctatus*. *J. Exp. Biol.* **86**, 171–185.
- Cameron, J. N. (1985). The bone compartment in a teleost fish *Ictalurus punctatus*: size composition and acid–base response to hypercapnia. *J. Exp. Biol.* **117**, 307–318.
- Cameron, J. N. and Kormanik, G. A. (1982). Intracellular and extracellular acid–base status as a function of temperature in the fresh water channel catfish *Ictalurus punctatus*. *J. Exp. Biol.* **99**, 127–142.
- Claiborne, J., Edwards, S. and Morrison-Shetlar, A. (2002). Acid–base regulation in fishes: cellular and molecular mechanisms. *J. Exp. Zool.* **293**, 302–319.
- Claiborne, J. and Heisler, N. (1984). Acid–base regulation and ion transfers in the carp (*Cyprinus carpio*) during and after exposure to environmental hypercapnia. *J. Exp. Biol.* **108**, 25–43.
- Claiborne, J. and Heisler, N. (1986). Acid–base regulation and ion transfers in the carp (*Cyprinus carpio*): pH compensation during graded long- and short-term environmental hypercapnia, and the effect of bicarbonate infusion. *J. Exp. Biol.* **126**, 41–61.
- Crocker, C. E. and Cech, J. J., Jr (1998). Effects of hypercapnia on blood-gas and acid–base status in the white sturgeon, *Acipenser transmontanus*. *J. Comp. Physiol. B.* **168**, 50–60.
- Gonzalez, R. J. and Dunson, W. A. (1987). Adaptations of sodium balance to low pH in a sunfish (*Erneacanthus obesus*) from naturally acidic waters. *J. Comp. Physiol. B.* **157**, 555–566.
- Goss, G., Perry, S. and Laurent, P. (1995). Ultrastructural and morphometric studies on ion and acid–base transport processes in freshwater fish. In *Fish Physiology: Cellular and Molecular Approaches to Fish Ionic Regulation* (ed. C. M. Wood and T. J. Shuttleworth), pp. 257–284. New York: Academic Press.
- Hazel, J., Garlick, W. and Sellner, P. (1978). The effects of assay temperature upon the pH optima of enzymes from poikilotherms: a test of the imidazole alphastat hypothesis. *J. Comp. Physiol. B* **123**, 97–104.
- Heisler, N. (1982). Intracellular and extracellular acid–base regulation in the tropical fresh-water teleost fish *Synbranchus marmoratus* in response to the transition from water breathing to air breathing. *J. Exp. Biol.* **99**, 9–28.
- Heisler, N. (1984). Acid–base regulation in fishes. In *Fish Physiology*, vol. X (ed. W. S. Hoar and D. J. Randall), pp. 315–401. San Diego: Academic Press.
- Heisler, N., Forcht, G., Ultsch, G. R. and Anderson, J. F. (1982). Acid–base regulation in response to environmental hypercapnia in two aquatic salamanders, *Siren lacertina* and *Amphiuma means*. *Resp. Physiol.* **49**, 141–158.
- Heming, T. A., Randall, D. J., Boutilier, R. G., Iwama, G. K. and Primmitt, D. (1986). Ionic equilibria in red blood cells of rainbow trout (*Salmo gairdneri*): Cl^- , HCO_3^- and H^+ . *Resp. Physiol.* **65**, 223–234.
- Jackson, D. (1997). Lactate accumulation in the shell of the turtle *Chrysemys picta bellii* during anoxia at 3°C and 10°C. *J. Exp. Biol.* **200**, 2295–2300.
- Jackson, D. C., Arendt, E. A., Inman, K. C., Lawler, R. G., Panol, G. and Wasser, J. S. (1991). ^{31}P -NMR study of normoxic and anoxic perfused turtle heart during graded CO_2 and lactic acidosis. *Am. J. Physiol.* **260**, R1130–R1136.
- Jackson, D. C., Ramsey, A. L., Paulson, J. M., Crocker, C. E. and Ultsch, G. R. (2000). Lactic acid buffering by bone and shell in anoxic softshell and painted turtles. *Physiol. Biochem. Zool.* **73**, 290–297.
- Larsen, B. K. and Jensen, F. B. (1997). Influence of ionic composition on acid–base regulation in rainbow trout (*Oncorhynchus mykiss*) exposed to environmental hypercapnia. *Fish Physiol. Biochem.* **16**, 157–170.
- Larsen, B. K., Portner, H. O. and Jensen, F. B. (1997). Extra- and intracellular acid–base balance and ionic regulation in cod (*Gadus morhua*) during combined and isolated exposures to hypercapnia and copper. *Mar. Biol.* **128**, 337–346.
- McCormick, S. D. (1993). Methods for nonlethal gill biopsy and measurement of Na^+ , K^+ ATPase activity. *Can. J. Fish. Aquat. Sci.* **50**, 656–658.
- McKenzie, D. J., Piccolella, M., Valle, A. Z. D., Taylor, E. W., Bolis, C. L. and Steffensen, J. F. (2003). Tolerance of chronic hypercapnia by the European eel *Anguilla anguilla*. *J. Exp. Biol.* **206**, 1717–1726.
- McKenzie, D. J., Taylor, E. W., Dalla Valle, A. and Steffensen, J. F. (2002). Tolerance of acute hypercapnic acidosis by the European eel (*Anguilla anguilla*). *J. Comp. Physiol. B* **172**, 339–346.
- Nikinmaa, M. (1990). *Vertebrate Red Cells: Adaptations of Function to Respiratory Requirements*. Berlin: Springer Verlag, pp. 262.
- Perry, S. F., Malone, S. and Ewing, D. (1987). Hypercapnic acidosis in the rainbow trout (*Salmo gairdneri*). II. Renal ionic fluxes. *Can. J. Zool.* **65**, 896–902.
- Pörtner, H., Boutilier, R., Tang, Y. and Toews, D. (1990). Determination of intracellular pH and PCO_2 after metabolic inhibition by fluoride and nitrilotriacetic acid. *Resp. Physiol.* **81**, 255–274.
- Pörtner, H. O., Reipschlag, A. and Heisler, N. (1998). Acid–base regulation, metabolism and energetics in *Sipunculus nudus* as a function of ambient carbon dioxide level. *J. Exp. Biol.* **201**, 43–55.
- Randall, D., Brauner, C. and Wilson, J. (1996). Acid excretion in Amazonian fish. In *Physiology and Biochemistry of the Fishes of the Amazon* (ed. A. L. Val, V. M. F. Almeida-Val and D. J. Randall), pp. 91–99. Manaus, Brazil: INPA.
- Reeves, R. B. (1985). What are normal acid–base conditions in man when body temperature changes. In *Acid–base Regulation and Body Temperature* (ed. H. Rahn and O. Prakash), pp. 13–32. Boston: Martinus Nijhoff Publishers.
- Soivio, A., Nyholm, K. and Westman, K. (1975). A technique for repeated sampling of the blood of individual resting fish. *J. Exp. Biol.* **63**, 207–217.
- Somero, G. N. and White, F. N. (1985). Enzymatic consequences under alphastat regulation. In *Acid–base regulation and body temperature* (ed. H. Rahn and O. Prakash), pp. 55–80. Boston: Martinus Nijhoff Publishers.

- Toews, D. P., Holeton, G. F. and Heisler, N.** (1983). Regulation of the acid-base status during environmental hypercapnia in the marine teleost fish *Conger conger*. *J. Exp. Biol.* **107**, 9-20.
- Val, A. L. and Almeida-Val, V. M. F.** (1995). *Fishes of the Amazon and Their Environment: Physiological and Biochemical Aspects*. Berlin: Springer-Verlag. pp. 224.
- Val, A. L., de Menezes, G. C. and Wood, C. M.** (1998). Red blood cell adrenergic responses in Amazonian teleosts. *J. Fish Biol.* **52**, 83-93.
- Wang, X., Heigenhauser, G. J. F. and Wood, C. M.** (1994). The analysis of metabolites in rainbow trout white muscle: a comparison of different sampling and processing methods. *J. Fish Biol.* **45**, 855-873.
- Wood, C. M.** (1989). The physiological problems of fish in acid waters. In *Acid Toxicity and Aquatic Animals* (ed. R. Morris, E. W. Taylor, D. J. A. Brown and J. A. Brown), pp. 125-152. Cambridge: Cambridge University Press.
- Wood, C. M., McDonald, D. G. and McMahon, B. R.** (1982). The influence of experimental anaemia on blood acid-base regulation *in vivo* and *in vitro* in the starry flounder (*Platichthys stellatus*) and the rainbow trout (*Salmo gairdneri*). *J. Exp. Biol.* **96**, 221-237.
- Wood, C. M., Turner, J. D., Munger, R. S. and Graham, M. S.** (1990). Control of ventilation in the hypercapnic skate *Raja ocellata*: II. Cerebrospinal fluid and intracellular pH in the brain and other tissues. *Resp. Physiol.* **80**, 279-298.
- Wood, S. C., Weber, R. E. and Davis, B. J.** (1979). Effects of air-breathing on acid-base balance in the catfish, *Hypostomus* sp. *Comp. Biochem. Physiol.* **62A**, 185-187.
- Zall, D., Fisher, D. and Garner, M.** (1956). Photometric determination of chlorides in water. *Anal. Chem.* **28**, 1665-1678.
- Zeidler, R. and Kim, D. H.** (1977). Preferential hemolysis of postnatal calf red cells induced by internal alkalization. *J. Gen. Physiol.* **70**, 385-401.