

Table S1. Composition of flower-visiting insect community according to different flower species

Flowers	Insects					References
	Bees	Flies	Lepidoptera	Wasps	Others	
<i>Rosa carolina</i>	92	8	0	0	0	Morse, 1979
<i>Bidens</i> sp.	41	56	2	1	0	Schmalhofer, 2001
<i>Solidago</i> sp.	26	66	5	3	0	Schmalhofer, 2001
<i>Centaurea scabiosa</i>	93	4	0	0	3	Brechbühl et al., 2009a
<i>Anthemis tinctoria</i>	79	12	0	0	9	Brechbühl et al., 2009a
<i>Erigenia bulbosa</i>	58	42	0	0	0	Dailey and Scott, 2006
<i>Claytonia virginica</i>	78	22	0	0	0	Dailey and Scott, 2006
<i>Dentaria laciniata</i>	74	14	0	0	12	Dailey and Scott, 2006
<i>Stellaria media</i>	81	17	0	0	2	Dailey and Scott, 2006
<i>Barbarea vulgaris</i>	84	14	0	0	2	Dailey and Scott, 2006
<i>Chrysanthemum frutescens/Anthemis tinctoria/Knautia arvensis</i>	78	5	0	5	12	Brechbühl et al., 2009b

All values in table indicate percentages. Bold values indicate the highest percentage value for each flower species.