

RESEARCH ARTICLE

Dynamic changes in global and gene-specific DNA methylation during hibernation in adult thirteen-lined ground squirrels, *Ictidomys tridecemlineatus*

Sebastian Alvarado^{1,*}, Timothy Mak¹, Sara Liu¹, Kenneth B. Storey² and Moshe Szyf^{1,‡}

ABSTRACT

Hibernating mammals conserve energy in the winter by undergoing prolonged bouts of torpor, interspersed with brief arousals back to euthermia. These bouts are accompanied by a suite of reversible physiological and biochemical changes; however, much remains to be discovered about the molecular mechanisms involved. Given the seasonal nature of hibernation, it stands to reason that underlying plastic epigenetic mechanisms should exist. One such form of epigenomic regulation involves the reversible modification of cytosine bases in DNA by methylation. DNA methylation is well known to be a mechanism that confers upon DNA its cellular identity during differentiation in response to innate developmental cues. However, it has recently been hypothesized that DNA methylation also acts as a mechanism for adapting genome function to changing external environmental and experiential signals over different time scales, including during adulthood. Here, we tested the hypothesis that DNA methylation is altered during hibernation in adult wild animals. This study evaluated global changes in DNA methylation in response to hibernation in the liver and skeletal muscle of thirteen-lined ground squirrels along with changes in expression of DNA methyltransferases (DNMT1/3B) and methyl binding domain proteins (MBDs). A reduction in global DNA methylation occurred in muscle during torpor phases whereas significant changes in DNMTs and MBDs were seen in both tissues. We also report dynamic changes in DNA methylation in the promoter of the *myocyte enhancer factor 2C* (*mef2c*) gene, a candidate regulator of metabolism in skeletal muscle. Taken together, these data show that genomic DNA methylation is dynamic across torpor–arousal bouts during winter hibernation, consistent with a role for this regulatory mechanism in contributing to the hibernation phenotype.

KEY WORDS: Torpor, Arousal, Cytosine, Epigenetics

INTRODUCTION

DNA methylation is a mechanism that confers upon identical genomes different chemical identities by covalently modifying cytosine bases at different positions in the genome (Razin and Riggs, 1980). This mechanism has been implicated in cellular differentiation for the last three decades and a large body of data has linked DNA methylation with regulation of transcription (Razin, 1998). DNA methylation is known to be dynamically altered during

embryonic development (Lister et al., 2009; Razin and Szyf, 1984; Razin et al., 1984), resulting in tissue-specific patterns of methylation that are maintained throughout life. However, the idea that this mechanism is not completely ‘fixed’ during development and could also play a role in stably altering genome function in response to external environmental exposures after embryonic development and even in adults has recently been gaining support. Whereas there is an emerging body of evidence suggesting that exposure to social adversity early in life (Weaver et al., 2004), different diets (Bollati et al., 2007; Waterland and Jirtle, 2003) or toxins could alter DNA methylation, it is unclear whether these constitute stochastic responses to signals modulating DNA methylation enzymes or whether there is an organized programmed genomic response to distinct and predictable environmental signals. One of the most remarkable organized responses to predictable environmental change over seasons is mammalian hibernation. It provides a stark example of how the same adult genotype undergoes an organized, predictable and comprehensive but nevertheless dynamic phenotypic change. This provides a unique real-life model to test the hypothesis that there is a programmed DNA methylation response to the environment in adult animals.

Mammalian hibernation involves extended bouts of torpor during which core body temperature can drop to near 0°C and metabolic energy expenditure can be reduced to as low as 1–5% of euthermic rates (Carey et al., 2003). The hibernation phenotype includes alterations to multiple behavioral, physiological, biochemical and molecular parameters. In thirteen-lined ground squirrels, *Ictidomys tridecemlineatus* (Mitchill 1821), these changes include programmed down-regulation of most mRNA and protein expression accompanied by a suite of transcriptional (Morin and Storey, 2006; Srere et al., 1992), translational (Wu and Storey, 2012b) and post-translational (Abnous et al., 2012; Morin and Storey, 2006) signatures consistent with quiescent cellular function (for review, see Storey and Storey, 2010). In contrast, selected transcription factors have been shown to be activated during bouts of torpor and arousal, indicating a complex interplay between gene repression, activation and metabolic depression (Tessier and Storey, 2010, 2012). Transcriptional differences occurring during hibernation are also expressed in a tissue-specific manner (Fujii et al., 2006). However, the exact regulatory mechanisms that mediate transcriptional activation and repression in these altered metabolic states are largely unknown, particularly at the level of global transcriptional controls (Hampton et al., 2011; Yan et al., 2008).

DNA methylation controls gene function by the steric hindrance of transcriptional activators or recruitment of transcriptional repressors (Razin and Riggs, 1980; Siegfried et al., 1999). Additionally, recent evidence links DNA methylation with the regulation of seasonal behaviors (Alvarado et al., 2014) and

¹Department of Pharmacology, McGill University, 3655 Sir William Osler Suite 1309, Montreal, QC, Canada H3G 1Y6; Sackler program in Epigenetics and Psychobiology at McGill University. ²Institute of Biochemistry and Department of Biology, Carleton University, 1125 Colonel By Drive, Ottawa, ON, Canada K1S 5B6. *Present address: Department of Biology, Stanford University, 371 Serra Mall Rm 314, Palo Alto, CA 94305, USA.

[‡]Author for correspondence (moshe.szyf@mcgill.ca)

List of symbols and abbreviations

DNMT	DNA methyltransferase
LUMA	luminometric methylation assay
MBD	methyl binding domain
MeCP	methyl CpG binding protein
Mef2C	myocyte enhancer factor 2C
T_b	body temperature

responses to seasonal disruptions (Cao-Lei et al., 2014) in various species. However, dynamic changes in the methylome often reflect broader changes that cannot be solely accounted for by transcriptional activation or repression of protein-coding genes (Alvarado et al., 2013; Tajerian et al., 2013). DNA methylation is catalyzed by DNA methyltransferases that target either newly replicated hemi-methylated DNA (DNMT1) or unmethylated DNA to mediate *de novo* DNA methylation (DNMT3A/B). DNA methylation is often read by a family of methyl binding domain proteins (MBDs) to mediate multiple functions of transcriptional repression and activation (Angrisano et al., 2006; Brown and Szyf, 2007; Ng et al., 1999). These regulators define a ‘methylation toolkit’ that is largely conserved in the animal kingdom and is capable of regulating genomic function in response to a dynamic environment (Jurkowski and Jeltsch, 2011; Lee et al., 2010).

To date, the role of epigenetic modification of DNA in mammalian hibernation has been limited to two studies that showed altered post-translational modification of histones and transcriptional repression (Morin and Storey, 2006) and aberrant genomic methylation in brown adipose tissue (Biggar and Storey, 2014). In addition, Fujii et al. (2006) analyzed the role of DNA methylation of the promoter region in regulating the expression of a liver gene in chipmunks that is specifically down-regulated in hibernation. Hypomethylation of a CpG dinucleotide at –161 within the promoter region in liver was linked with tissue-specific expression of the *HP-27* gene (the site was hypermethylated in kidney and heart) but methylation status at –161 and three other CpG dinucleotides did not change in a comparison of liver from hibernating and non-hibernating chipmunks (Fujii et al., 2006). Hence, to advance the field, the present study tested the hypothesis that DNA methylation dynamically responds to cyclic changes occurring over torpor–arousal bouts in adult mammals. Our analysis focused on liver and skeletal muscle. Liver is of central importance to metabolic homeostasis of the whole animal including involvement in torpor-specific fuel metabolism (e.g. gluconeogenesis and ketone body synthesis to provide fuel for the brain), whereas muscle, although inactive during most of a torpor bout, plays a key role in arousal by generating heat via shivering thermogenesis. Both tissues exhibit a range of transcriptional, translational and post-transcriptional changes during hibernation and metabolic depression (Abnous et al., 2008; Green et al., 1984; Mamady and Storey, 2008; Storey, 1987). As the transcriptional and translational response in hibernation involves global changes, we first evaluated genomic changes in DNA methylation and its conserved machinery. We then used a candidate gene approach to measure DNA methylation in a gene that plays a role in muscle metabolic regulation during hibernation, the transcription factor *myocyte enhancer factor 2C* (*mef2c*) (Tessier and Storey, 2010). *mef2c* is a master transcription factor that plays roles in cellular differentiation including myogenesis, neurogenesis, and development of the heart and the neural crest (Potthoff and Olson, 2007). In mammals, *mef2c* is predominantly expressed in the brain and skeletal muscle with a suggested role for the maintenance of the myocyte differentiated state (Breitbart et al., 1993).

RESULTS**Global DNA methylation changes during hibernation**

In order to evaluate whether global DNA methylation states responded to changes related to hibernation (Fig. 1A), we assayed global DNA methylation in *I. tridecemlineatus* using LUMA (luminometric methylation assay) in two tissues: skeletal muscle and liver. In skeletal muscle, significant hypomethylation (an approximate 16% reduction, assuming 100% cutting by restriction enzymes) was observed in animals with cold body temperature (T_b) values that indicated they were in late torpor, just beginning to arouse or in interbout (Fig. 1B) as compared with animals that were euthermic in the cold (control). A query of the ground squirrel genome for *HpaII/MspI* sites (Ensemble release 66) indicated that the extent of changes in methylation that were observed in our study corresponded, at the very least, to 206,573 differentially methylated CpG sites in the muscle genome. If we consider 16% of CGs outside of the CCGG restriction site, this could represent the entire population of genomic CGs ranging close to ~1.5 million CGs. Significant changes in global methylation state were observed in the liver between animals that were euthermic in the cold and those in interbout (Fig. 1E).

Altered expression of DNA methylation regulators accompanies hibernation and seasonal change

Considering the changes in DNA methylation levels that occurred in muscle over torpor–arousal, we next investigated whether these changes were associated with overall changes in the expression of known DNA methylation regulators. We assessed changes in the mRNA transcript levels of DNA methyltransferases (*dnmt1*, *dnmt3b*) over the course of hibernation bouts. We observed that DNMT transcript levels in skeletal muscle did not change significantly over the torpor–arousal cycle (Fig. 1C,D). Thus, the observed reduction in global DNA methylation in muscle is not a consequence of a major change in DNMT transcription. However, in the liver where we saw no genomic changes in DNA methylation, we observed differences in *dnmt1* and *dnmt3b* expression between animals that were euthermic in the cold versus those at the early torpor stage (Fig. 1F,G). Thus, changes in DNMT were not correlated with global changes in DNA methylation. Knowing that the DNA methylation signal is in part transduced by several proteins, we measured mRNA levels of four MBD proteins [*mbd1*, *mbd2*, *mbd3*, *mecp2* (methyl CpG binding protein 2)] over the hibernation stages. Expression of these regulators was altered both between tissues and across the torpor–arousal cycle (Fig. 2). In muscle, a 14-fold increase was seen in *mecp2* mRNA levels between animals that were euthermic in the cold and those in the early arousal stage (Fig. 2, top). In liver, we observed significant increases in transcript levels between animals that were euthermic in the cold and those in early or late torpor for *mbd2* (Fig. 2, bottom), and between animals that were euthermic in the cold and those in the early arousal stage for *mbd3* and *mecp2* (Fig. 2, bottom). Although these changes in expression cannot directly explain the changes in global DNA methylation observed in skeletal muscle, they do support the hypothesis that the DNA methylation machinery in this candidate screen is altered during different stages of hibernation.

Dynamic expression and promoter methylation of the candidate regulator of muscle function, *mef2c*, across hibernation

We examined *mef2c* because its expression has been shown to be altered during hibernation in a tissue-specific manner (Bratincsak et al., 2007; Tessier and Storey, 2010, 2012). In skeletal muscle, the




Fig. 1. Levels of genomic DNA methylation over the torpor–arousal cycle of winter hibernation in thirteen-lined ground squirrels, *Ictidomys tridecemlineatus*. (A) A scheme of body temperature (T_b) variations over the course of torpor–arousal bouts showing the different sampling points used in this study. (B–D) Global DNA methylation levels across torpor–arousal in skeletal muscle (B) and expression levels of DNA methyltransferases *dnmt1* (C) and *dnmt3b* (D). (E–G) Global DNA methylation levels across torpor–arousal in liver (E) and expression levels of *dnmt1* (F) and *dnmt3b* (G). Data in histograms are means+s.e.m., $N=7$ independent samples from different animals. One-way ANOVA was carried out followed by a Dunnett's *post hoc* test. Asterisks indicate a significant difference from the control (euthermic in the cold) value, * $P<0.05$, ** $P<0.01$, **** $P<0.0001$.

expression of the gene was dramatically reduced during entry into torpor, when T_b was decreasing towards the stable low T_b value (Fig. 3A, left). Expression was slightly increased again in early torpor but remained low in late torpor and early arousal. In contrast, in the liver, *mef2c* gene expression remained low across all torpor stages and was up-regulated only during the early arousal phase (Fig. 3A, right).

We then used bisulfite pyrosequencing of the *mef2c* promoter (Fig. 3D) in skeletal muscle and liver to follow its state of methylation over the different hibernation stages. The promoter region of *mef2c* was predicted from an alignment to the human *mef2c* promoter (Fig. 3B,C) and was confirmed by transient expression using a luciferase expression vector in HEK293 cells (Fig. 3E). Several CpG sites upstream of the *mef2c* transcriptional start site (–297, –299, –321, –329, –331, –351 and –353) showed

dynamic changes between two hibernation stages in skeletal muscle but not in liver (Fig. 3D). A large increase in the methylation state of all CG sites across the promoter region of skeletal muscle *mef2c* occurred as the animals entered torpor (Fig. 3D, left), which correlates with the sharp reduction in gene expression (Fig. 3A, left). However, the CG sites were demethylated again during early torpor but showed a slight remethylation in late torpor before methylation was reduced again. A minor demethylation was again apparent upon early arousal but did not correlate with induced gene expression. The *mef2c* gene was silenced in the liver through most of the stages of hibernation (Fig. 3A, right); however, the promoter was methylated at the same level as in skeletal muscle (Fig. 3D, right) and the spike of *mef2c* expression during early arousal was not accompanied by a demethylation of the gene promoter. It is possible that a DNA methylation-independent process silences the




Fig. 2. Differential expression of mRNA transcript levels of methyl binding domain proteins (MBDs) across the torpor–arousal cycle of hibernation. Top, expression of *mbd1*, *mbd2*, *mbd3* and *mecp2* in skeletal muscle. Bottom, expression of *mbd1*, *mbd2*, *mbd3* and *mecp2* in liver. Other information as in Fig. 1. One-way ANOVA was carried out followed by a Dunnett's *post hoc* test. Asterisks indicate a significant difference from the control (euthermic in the cold) value, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

expression of this gene in the liver and is responsible for its induction during early arousal; for example, the formation of RNA stress granules (Tessier and Storey, 2014). Alternatively, this could indicate that this particular region is a muscle-specific promoter or enhancer that does not regulate liver-associated transcripts.

The above results suggested that DNA methylation might be involved in silencing *mef2c* promoter activity. Therefore, to investigate this further we analyzed whether methylation of the CG sites in the promoter would alter the ability of *mef2c* to direct transcription of a luciferase mRNA in a transient expression assay (Klug and Rehli, 2006). The *mef2c* promoter region was cloned into a CpG-less promoter construct in sense and antisense (as a negative control) directions. The plasmids were either *in vitro* methylated or mock methylated by the bacterial CG DNA methyltransferase *mSss1* (Nur et al., 1985). As the vector lacks CG sites, this will result in methylation only of the sites in the *mef2c* promoter and will thus recapitulate the situation *in vivo* where all of these sites were found to be methylated during entry into torpor (Fig. 3D, left). The plasmids were transfected into HEK293 cells and assayed after 48 h. The unmethylated promoter in the sense orientation expressed higher levels of luciferase compared with the background levels expressed from the antisense promoter (Fig. 3E). Furthermore, methylation of the sense promoter reduced the level of expression to the background levels expressed from the antisense promoter. These results show that DNA methylation is capable of reducing the transcriptional activity of the *mef2c* promoter *in vitro* and further supports evidence of muscle-specific regulation of transcription *in vivo*.

DISCUSSION

Epigenomic mechanisms such as DNA methylation are well suited to serve as an interface between the DNA of a cell and changing environments, but the main question is whether there exists an organized and predicted response of the methylome to natural environmental changes. Our current understanding of epigenetic mechanisms is largely limited to studies in human pathology or embryogenesis. As a result, very little is known about the role of epigenetic modification in adults facing naturally dynamic environments. Hibernation is one of the most dramatic phenotypic

responses to change in the climactic environment and could serve as an excellent paradigm for testing the response of the methylome to extreme changes in physiology. Hibernating mammals are capable of reducing their metabolic expenditures by 95–99% resulting in a very strong suppression of multiple core processes including respiration, heartbeat, transcription (Osborne et al., 2004), translation and the cell cycle (Frerichs et al., 1998; Wu and Storey, 2012a) over torpor bouts that can last for days or weeks. Considering the genetic similarity between seasonal hibernators and non-hibernators (Carey et al., 2003; Srere et al., 1992), this phenotype may not require novel genes but rather the reconfiguration of the transcriptome. In this respect, seasonal hibernators may be reiterating a neonatal program in preparation for torpor – suggesting that the genetic potential for heterothermy that exists in all mammalian neonates is invoked in adult hibernators by the continued or renewed expression of selected genes that are crucial for heterothermy (Harris et al., 2004).

In the present study, we show that DNA methylation, its effectors and its targets can be altered in hibernating mammals over winter torpor–arousal cycles. We further show tissue-specific genomic DNA methylation changes that accompany the torpor–arousal cycle and uncoupled transcription of methyltransferases and differential regulation of MBD proteins. We observed that the differentially expressed *mef2c* transcript was a tissue-specific target of differential DNA methylation *in vivo* during the hibernation cycle (Fig. 3), and that hypermethylation upon entrance into torpor is partly associated with reduced mRNA expression. The sites that are hypermethylated upon entrance into torpor could silence the *mef2c* promoter as determined by the promoter–luciferase assays (Fig. 3E). To our knowledge, this study is the first to demonstrate dynamic changes in genomic DNA methylation and its regulators in seasonal hibernating mammals. This study illustrates DNA methylation as a dynamic process in adult tissues that can be programmed to respond to distinct environmental cues. Although hibernation is a unique and extreme example of mammalian response to environment change, it provides a proof of principle that the genome of adult mammals can respond to environmental stress with tissue-specific changes in DNA methylation. Thus, this study provides evidence in support of the hypothesis that dynamic DNA methylation programs are not limited




Fig. 3. Dynamic methylation of the *mef2c* promoter at different hibernation stages. (A) Expression of *mef2c* mRNA measured in muscle (left) and liver (right). (B) Alignment of the *I. tridecemlineatus* *mef2c* promoter against the human *mef2c* promoter. (C) Physical map of the CpG dinucleotides in the *mef2c* region highlighted with balloons. (D) Bisulfite map of DNA methylation in the *mef2c* promoter in muscle (left) and liver (right). (E) Luciferase reporter constructs containing the *mef2c* regulatory region in pCpG in sense (S), antisense (AS) and methylated sense (mS) directions were transfected into HEK293 cells and their ability to direct expression of a downstream coding sequence was determined by assaying luciferase activity (RLuc, relative luciferase activity after normalization to total protein measurement) in extracts of transfected cells. Other information as in Fig. 1. One-way ANOVA was carried out followed by a Dunnett's *post hoc* test with euthermic in the cold as our control. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

to embryogenesis and that they can change in a predictable manner in response to external environmental and physiological signals.

Tissue-specific genomic methylation during hibernation

Our data show that the response of DNA methylation is tissue specific as liver and muscle altered their DNA methylation patterns differently during different phases of hibernation. In muscle, genomic DNA methylation decreased dramatically during late torpor, early arousal and interbout stages of hibernation (Fig. 1B) on a scale similar to that reported for global hypomethylation in cancer (Feinberg and Vogelstein, 1983; Lu et al., 1983) or autoimmune diseases (Cornacchia et al., 1988). In liver, genomic DNA methylation increased in animals during interbout (Fig. 1E). It should be noted that the use of LUMA in our study is limited in only being able to assay absolute changes in genomic methylation. As a result, we are unable to offer a comprehensive insight into which genes are differentially methylated, whether methylation is happening outside of a CpG context, or whether a reconfigured

methylome results in a zero sum change in genomic methylation. The exact role of the observed hypomethylation in muscle in controlling genome function is unknown as the extent of DNA demethylation is broader than what would be anticipated if only CpGs in regulatory regions and promoters were demethylated. This suggests that demethylation as a regulatory mechanism must involve wider genomic regions. At the very least, the genomic hypomethylation observed within the muscle represents a molecular rearrangement that is supported by the overall transcriptional activity observed in the skeletal muscle of hibernating mammals (Malatesta et al., 2009). This needs to be explored in the future with full genomic mapping of DNA methylation over different hibernation stages. Nevertheless, our study shows that global hypomethylation is a clear component of the altered physiological state of muscle associated with seasonal hibernation and it stands to reason that it has an important role in regulating genome function. Particularly interesting is the possibility that global DNA methylation may alter structural features of the genome (Espada

et al., 2007) considering that altered nucleolus structure and aggregation of RNA binding proteins into subnuclear particles has been reported during hibernation (Malatesta et al., 2011; Tessier et al., 2014).

Uncoupled transcription of epigenetic regulators from genomic DNA methylation

Given the differential genomic methylation over the course of torpor–arousal bouts, we further investigated whether altered transcription of DNA methyltransferases also occurred as a possible mechanism for the observed changes in methylation. We focused our interpretation of the data on the window of time that defined the greatest changes in genomic methylation of skeletal muscle, i.e. the stages from euthermic animals in the cold through to early arousal (Fig. 1A). Significantly altered expression of *dnmt* transcripts was only observed in the liver (Fig. 1F,G), but the changes in expression were not associated with changes in genomic DNA methylation at different stages of hibernation. Furthermore, no significant changes in *dnmt* expression were observed across the torpor–arousal cycle of hibernation in skeletal muscle (Fig. 1C,D) despite decreases in global DNA methylation (Fig. 1A). If a model of maintenance methylation is to be considered, this could further implicate trans-acting mechanisms known to regulate methylation, such as post-translational (Kang et al., 2001) or post-transcriptional (Garzon et al., 2009) suppression of DNA methyltransferases. However, given the scale and nature of changes observed in muscle, we propose mechanisms of hypomethylation that act independently of methyltransferase function through passive (Kagiwada et al., 2013) or active demethylation (Kohli and Zhang, 2013). In passive demethylation, muscle-specific protection from atrophy and wasting during hibernation (Lin et al., 2012) suggests the possibility of myosatellite rejuvenation of the cell pool in the absence of DNMTs, leading to genomic hypomethylation. We speculate this molecular regulation of the myocyte cell line based on previous reports where this tissue was shown to have the highest variability in genomic methylation (Horvath, 2013), suggesting a dynamic role in cell fate during self-renewal (Motohashi and Asakura, 2012). Similarly, active demethylation of the genome through a DNA demethylase could accompany hypomethylation. Currently, a bona fide DNA demethylase has yet to be fully characterized despite several lines of evidence in non-dividing cells and tissues within the brain (Bocker et al., 2011; Guo et al., 2011; Ma et al., 2009). In this respect, it has been suggested that DNA demethylation occurs through hydroxylation of methylcytosine mediated through complexes of the ten-eleven-translocation (TET) family of enzymes (Kohli and Zhang, 2013).

While changes in the transcription of DNMTs between euthermic and early torpor animals were not accompanied by global changes in DNA methylation in liver, their expression does not necessarily drive a genome-wide change in methylation. For example, many studies examining the localization of methyltransferases with respect to other repressive marks demonstrate the targeted hypermethylation of a small subset of the genome (Jin et al., 2012). Additionally, because of the limitations of LUMA, it would not be possible to discern small changes that affect a targeted area of the genome. These DNMTs could likely serve a function in *de novo* methylation independent of DNA replication due to cell cycle arrest seen in the liver during bouts of torpor (Wu and Storey, 2012a). Previous studies have shown that the expression of *dnmt3b* is mediated by the transcription factor Bmal1 with circadian rhythms in the liver known to underlie metabolic rate depression during hibernation in ground squirrels (Maekawa et al., 2012; Ruby et al., 2002).

The MBD family of proteins retains a conserved methyl binding domain motif that serves as a reader of DNA methylation patterns and mediates the activities of DNA methylation (Nan et al., 1993). These proteins can be present in varying amounts that do not necessarily rely on the abundance of genomic methylation; for example, MBD2 and MBD4 are overexpressed in CD4⁺ T-cells that exhibit genomic hypomethylation in erythematosus lupus (Balada et al., 2007). MBDs can act in both repressor (Klose and Bird, 2006; Ng et al., 1999) and activator (Baubec et al., 2013; Bhattacharya et al., 1999; Brown and Szyf, 2007) roles. In this study, we identified differential expression of MBD transcripts in both liver and muscle over torpor–arousal during hibernation. In muscle, *mecp2* expression changed between euthermia and arousal (Fig. 2, top). In the liver, *mbd3* and *mecp2* mRNA levels differed between animals that were euthermic in the cold room and those in early arousal. We speculate that *mbd3* transcription may support the activation of rRNA, as seen in previous studies (Brown and Szyf, 2007), consistent with the need for rapid reactivation of liver ribosomal protein synthesis when animals arouse from bouts of torpor (Epperson et al., 2004; Fedorov et al., 2009). Additionally, in muscle, MeCp2 has been reported to be involved in myogenesis as its role in forming pericentric heterochromatic centers is important for differentiation (Brero et al., 2005). However, these roles of MBDs in regulating changes in genomic function are intricate in many tissues (Gunther et al., 2013; Yasui et al., 2013) and will require additional studies to validate their actions within hibernation stages.

Tissue-specific methylation control of *mef2c*

In addition to our genome-wide screen, a candidate gene screen revealed a unique promoter-specific pattern of methylation that is dynamic during bouts of torpor. We focused on *mef2c* because previous studies showed that it is upregulated during torpor along with downstream targets GLUT4 and MyoD in skeletal muscle (Tessier and Storey, 2010, 2012). *mef2c* was particularly interesting to focus on because of its activity as a transcription factor known to regulate many downstream targets of muscle development (Potthoff et al., 2007a). For example, *mef2c* has been shown to directly regulate myomesin and M-protein, known mediators of skeletal muscle function (Potthoff et al., 2007a). *mef2c* deletion and overexpression was also shown to regulate activity-dependent fast to slow fiber transformation in skeletal muscle. Interestingly, the overexpression of *mef2c* can promote the slow fiber phenotype in skeletal muscle, enhancing endurance and enabling mice to run twice the distance of wild-type littermates (Potthoff et al., 2007b). Given a tentative role in maintaining the myocyte state (Breitbart et al., 1993) and endurance, we speculate that these roles support each other in a transcriptional program that sustains the functionality of skeletal muscle during hibernation bouts. This would support the crucial role of skeletal muscle in shivering thermogenesis to rewarm animals during arousal and potentially minimize/avoid atrophy during torpor so that the aroused animal is prepared to resume euthermic life.

Regarding DNA methylation, a previous study showed that pharmacologically induced hypomethylation in murine myocytes could enhance myogenesis (Montesano et al., 2013). Interestingly, induced transcripts in this study such as MyoD are known downstream targets of *mef2c* in *I. tridecemlineatus* during bouts of torpor (Tessier and Storey, 2010). Furthermore, methylation of Mef2 binding sites within gene promoters inhibited transcription factor binding (Palacios et al., 2010). We showed dynamic changes in DNA methylation of the *mef2c* gene during bouts of hibernation in muscle but not in the liver, while expression changes occurred in both tissues. It is important to note that methylation signatures and their

Table 1. Primer sequences used for bisulfite sequencing, *in vitro* promoter assays and expression analyses

Primer	Forward	Reverse
qpcr DNMT1	AGGCGGCTCAAAGACTTG	TCAATTTCTCCTTCACACTC
qpcr DNMT3B	CTGCAAGCCCAGCAGTCC	GGACTTGGAGCACTGTTGTTG
qpcr MBD1	GGAGACCCAAGAGGATGAG	AGCTGATTCCACAGTTCTCAC
qpcr MBD2	AAATAAGGGTAAACCAGACTTG	ACTTTGGTACTGGTTGTTTG
qpcr MBD3	AACCAGGTCAAAGGCAAG	TTAAAGATGGACGCTGTCTG
qpcr MEF2C	CGTCTCCAGCTCTCTGCTCT	CAGCCTTGAAGTGCTTCTCC
bis mef2c	AATAGTTGTTTTGTGATTTTTTTTT	TTTAAACCCCTCCACTTCTCAAAAC
seq mef2c	TTGTTTTGTGATTTTTTTTTTTTATT	
luc mef2c S	GGATCCTATAAAAGTCGGGATCTTTCCTCTG	AAGCTTTATATCTTTCGGTTCACTTTTAGCC
luc mef2c AS	AAGCTTTATAAAAGTCGGGATCTTTCCTCTG	GGATCCTATATCTTTCGGTTCACTTTTAGCC

function are tissue and cell type specific (Razin and Riggs, 1980). It is possible that *cis*-regulatory regions (enhancers and promoters) may constrain *mef2c* DNA demethylation to specific tissues (Andres et al., 1995). Although it is clear that in the muscle the region analyzed is not involved in silencing of the gene, our *in vitro* luciferase assay of the *mef2c* promoter at the very least validates that DNA methylation is capable of inhibiting its transcriptional activity, supporting our interpretation of muscle-specific *mef2c* regulation.

Taken together, our data demonstrate that DNA methylation is dynamically altered during hibernation and that there can be an organized response by DNA methylation to environmental cues in adult tissues. However, it is still unknown what the physiological role of these DNA methylation events is in orchestrating the molecular changes seen in hibernating mammals over the torpor–arousal cycle. Additionally, considering the tissue-specific expression of epigenetic regulators, this study begs for more comprehensive studies aimed at understanding how genome-wide changes in methylation affect specific gene functions. To our knowledge, our work is the first to report differential DNA methylation (and its machinery) within torpor–arousal cycles in a mammalian hibernator. We also characterized, for the first time (*in vivo* and *in vitro*), promoter methylation of a gene that is differentially regulated in hibernation. This work demonstrates that DNA methylation is dynamic and capable of changing significantly on both genome-wide and gene-specific levels long after mammals have developed and reached maturity. Considering the time scales over which these changes occur, the data also provide a new, but limited, understanding of the chronicity in which DNA methylation can change in cycles (in a matter of days). Whereas these mechanisms have been classically studied in cancer and disease states, few studies have examined the roles that dynamic DNA methylation in adult tissues could play in natural biological phenomena, thus emphasizing its importance to other fields of organismal and ecological biology.

MATERIALS AND METHODS

Animal collection and hibernation protocol

Thirteen-lined ground squirrels, *I. tridecemlineatus*, were captured by a trapper licensed by the United States Department of Agriculture (TLS Research, Bartlett, IL, USA) using humane traps. Male squirrels were used for hibernation studies as previously described; temperature transponders implanted under the skin allowed T_b to be monitored to identify sampling times (McMullen and Hallenbeck, 2010). Studies took place within the natural hibernation season (January and February) in the northern hemisphere. For sampling, animals were anesthetized with isoflurane and decapitated within 2 min of removal from the hibernation chamber. Tissues were rapidly excised, flash frozen in liquid nitrogen, transported to their destination on dry ice and then stored at -80°C until use. All animal procedures were approved by the Animal Care and Use Committee of the National Institute of Neurological Disorders and Stroke (National Institutes of Health, Bethesda, MD, USA; animal protocol no. ASP 1223-05). Animals were killed at the following seven stages as illustrated in Fig. 1A (McMullen

and Hallenbeck, 2010): (i) euthermic at room temperature, (ii) euthermic in the cold room for 3 days (T_b 34–37°C), (iii) entry into torpor with decreasing T_b ($18^\circ\text{C} \leq T_b \leq 31^\circ\text{C}$), (iv) early torpor with core T_b of 5–7°C for ~24 h, (v) late torpor with core T_b between 5 and 7°C for at least 3 days, (vi) early arousal from torpor with T_b rising to at least 12°C and (vii) interbout arousal with euthermic T_b re-established at 36–38°C.

DNA/RNA extraction

Thigh hindleg skeletal muscle and liver were collected in biological replicates of $N=7$ for each sampling stage. Tissues were ground to a powder under liquid nitrogen and split into 300 mg aliquots for separate RNA and DNA extractions. Tissues were homogenized with a Tissue Tearor (Biospec, Bartlesville, OK, USA) in Trizol (Invitrogen) for RNA extraction and extracts were prepared according to the manufacturer's protocols (Invitrogen). The quantity and quality of RNA were determined using spectrophotometric analysis. DNA extractions were carried out in an extraction buffer (500 μl) containing proteinase K (20 μl :20 mg μl^{-1} ; Roche, Basel, Switzerland) at 56°C for 12 h. Samples were then treated with RNase A (50 U mg^{-1} , 30 min; Roche) and phenol chloroform (1:1 v:v). DNA was pelleted with 70% ethanol and resuspended in 100 μl of buffer (10 mmol l^{-1} Tris-HCl, 1 mmol l^{-1} EDTA). The quantity and quality of DNA were determined using spectrophotometric analysis.

Bisulfite pyrosequencing

DNA was treated with sodium bisulfite as described previously (Clark et al., 1994) and primers were designed for a converted 136 bp fragment (scaffold name: JH393279.1; 7205500–7205635 according to SQUIRREL Ensembl Release 66) of the *mef2c* promoter. This promoter was predicted from an alignment of the fully annotated human *mef2c* gene with the *I. tridecemlineatus* homolog. Bisulfite-treated *mef2c* promoter region PCRs were amplified using two rounds of PCR with outer and nested primers (see Table 1). Cycling conditions involved an initial step of 5 min at 95°C followed by 35 cycles of 95°C for 1 min, 60°C for 2.5 min and 72°C for 1 min; the final step was a 5 min hold at 72°C. PCR products were sequenced using the Biotage Pyrosequencer according to the manufacturer's protocol (Biotage, Uppsala, Sweden).

Expression analysis

For all samples, 500 ng of RNA was subjected to RT-PCR according to the manufacturer's protocols (Roche) and quantified using quantitative PCR on the Lightcycler 480 (Roche, Indianapolis, IN, USA). Primers for all genes (see Table 1) were created across exon boundaries using Autoprime (Wrobel et al., 2004) and were validated with single peak melting curves following PCR and matching product sizes on agarose gels. *gapdh* was used as a reference gene for relative quantification as it was previously shown to have stable expression across hibernation bouts in *I. tridecemlineatus* (Wu and Storey, 2012a,b). Quantitative PCR was conducted with a pre-incubation at 95°C for 10 min followed by 45 cycles of 95°C for 10 s, 60°C for 10 s, 72°C for 10 s, with a final hold of 10 min at 72°C.

LUMA

LUMA was used to assay global DNA methylation across the genome. Our method was based on that described by Karimi et al. (2006b) with minor

modifications. LUMA involves the digestion of genomic DNA by methylation-sensitive (*HpaII*) or insensitive (*MspI*) restriction enzymes in combination with an internal control restriction enzyme (*EcoRI*) to normalize DNA input. *EcoRI* (R0101), *HpaII* (R0171) and *MspI* (R0106) were all purchased from New England Biolabs. Samples were incubated (37°C, 4 h) and then heat inactivated (80°C, 20 min). Digested genomic DNA (15 µl) was mixed with pyrosequencing annealing buffer (15 µl; Qiagen, Toronto, ON, Canada) and pyrosequenced according to the manufacturer's guidelines (PyroMark 24; Biotage). The nucleotide dispensation order used was based on Karimi et al. (2006a,b). All samples were run in technical triplicates with biological replicates of seven. Data were then presented as an index of overall genomic methylation calculated as (*HpaII/EcoRI*)/(*MspI/EcoRI*). This is a measure of CCGs that are demethylated divided by methylated CCGs.

In vitro luciferase assay

Luciferase construct PCR primers (Table 1) were made to amplify the 136 bp fragment of the *mef2c* promoter. *BamHI* and *HindIII* restriction sites were incorporated into primers in order to clone them into the CpG-less pCpGI (Klug and Rehli, 2006) and primers were cloned in 5' to 3' (sense) or 3' to 5' (antisense) orientation, respectively. The constructs were methylated *in vitro* with *SssI* CpG DNA methyltransferases (M0226L, New England Biolabs) as recommended by the manufacturer. Transfections were performed using calcium phosphate precipitation as described previously (Champagne et al., 2006) into HEK293 cells (CRL-1573, ATCC). Cells were harvested 48 h after transfection and luciferase activity was assayed using the Luciferase Assay System (Promega).

Statistical analysis

Data are expressed as means±s.e.m., *N*=7 independent samples from different animals. A one-way ANOVA was performed on each dataset followed by a Dunnett's test assigning euthermic in the cold room as control. Outliers were excluded based on Grubb's outlier tests on the data for a given stage of hibernation. Significance was set at *P*<0.05. Statistical analysis was undertaken using Prism (GraphPad Software Inc., San Diego, CA, USA).

Competing interests

The authors declare no competing or financial interests.

Author contributions

S.A., K.B.S. and M.S. designed the experiments. S.A., T.M. and S.L. performed experiments. S.A., K.B.S. and M.S. analyzed the data. S.A., K.B.S. and M.S. wrote the manuscript.

Funding

This study was funded by a grant from the Canadian Institute of Health awarded to M.S. (MOP-42411). We thank Dr J. M. Hallenbeck and Dr D. McMullen (National Institute of Neurological Disorders and Stroke, NIH, Bethesda, MD, USA) for providing the tissue samples for this study. S.A. was funded by the Sackler Foundation for Mental Health. K.B.S. was funded by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery grant and holds the Canada Research Chair in Molecular Physiology.

References

Abnous, K., Dieni, C. A. and Storey, K. B. (2008). Regulation of Akt during hibernation in Richardson's ground squirrels. *Biochim. Biophys. Acta* **1780**, 185–193.

Abnous, K., Dieni, C. A. and Storey, K. B. (2012). Suppression of MAPKAPK2 during mammalian hibernation. *Cryobiology* **65**, 235–241.

Alvarado, S., Tajerian, M., Millecamps, M., Suderman, M., Stone, L. S. and Szyf, M. (2013). Peripheral nerve injury is accompanied by chronic transcriptome-wide changes in the mouse prefrontal cortex. *Mol. Pain* **9**, 21.

Alvarado, S., Fernald, R. D., Storey, K. B. and Szyf, M. (2014). The dynamic nature of DNA methylation: a role in response to social and seasonal variation. *Integr. Comp. Biol.* **54**, 68–76.

Andres, V., Cervera, M. and Mahdavi, V. (1995). Determination of the consensus binding site for MEF2 expressed in muscle and brain reveals tissue-specific sequence constraints. *J. Biol. Chem.* **270**, 23246–23249.

Angrisano, T., Lembo, F., Pero, R., Natale, F., Fusco, A., Avvedimento, V. E., Bruni, C. B. and Chiariotti, L. (2006). TACC3 mediates the association of MBD2 with histone acetyltransferases and relieves transcriptional repression of methylated promoters. *Nucleic Acids Res.* **34**, 364–372.

Balada, E., Ordi-Ros, J., Serrano-Acedo, S., Martínez-Lostao, L. and Vilardell-Tarres, M. (2007). Transcript overexpression of the MBD2 and MBD4 genes in CD4+ T cells from systemic lupus erythematosus patients. *J. Leukoc. Biol.* **81**, 1609–1616.

Baubec, T., Ivánek, R., Lienert, F. and Schübeler, D. (2013). Methylation-dependent and -independent genomic targeting principles of the MBD protein family. *Cell* **153**, 480–492.

Bhattacharya, S. K., Ramchandani, S., Cervoni, N. and Szyf, M. (1999). A mammalian protein with specific demethylase activity for mCpG DNA. *Nature* **397**, 579–583.

Biggar, Y. and Storey, K. B. (2014). Global DNA modifications suppress transcription in brown adipose tissue during hibernation. *Cryobiology* **69**, 333–338.

Bocker, M. T., Hellwig, I., Breiling, A., Eckstein, V., Ho, A. D. and Lyko, F. (2011). Genome-wide promoter DNA methylation dynamics of human hematopoietic progenitor cells during differentiation and aging. *Blood* **117**, e182–e189.

Bollati, V., Baccarelli, A., Hou, L., Bonzini, M., Fustinoni, S., Cavallo, D., Byun, H.-M., Jiang, J., Marinelli, B., Pesatori, A. C. et al. (2007). Changes in DNA methylation patterns in subjects exposed to low-dose benzene. *Cancer Res.* **67**, 876–880.

Bratincsak, A., Brownstein, M. J., Cassiani-Ingoni, R., Pastorino, S., Szalayova, I., Toth, Z. E., Key, S., Németh, K., Pickel, J. and Mezey, E. (2007). CD45-positive blood cells give rise to uterine epithelial cells in mice. *Stem Cells* **25**, 2820–2826.

Breitbart, R. E., Liang, C. S., Smoot, L. B., Laheru, D. A., Mahdavi, V. and Nadal-Ginard, B. (1993). A fourth human MEF2 transcription factor, hMEF2D, is an early marker of the myogenic lineage. *Development* **118**, 1095–1106.

Brero, A., Easwaran, H. P., Nowak, D., Grunewald, I., Cremer, T., Leonhardt, H. and Cardoso, M. C. (2005). Methyl CpG-binding proteins induce large-scale chromatin reorganization during terminal differentiation. *J. Cell Biol.* **169**, 733–743.

Brown, S. E. and Szyf, M. (2007). Epigenetic programming of the rRNA promoter by MBD3. *Mol. Cell. Biol.* **27**, 4938–4952.

Cao-Lei, L., Massart, R., Suderman, M. J., Machnes, Z., Elgbeili, G., Laplante, D. P., Szyf, M. and King, S. (2014). DNA methylation signatures triggered by prenatal maternal stress exposure to a natural disaster: project ice storm. *PLoS ONE* **9**, e107653.

Carey, H. V., Andrews, M. T. and Martin, S. L. (2003). Mammalian hibernation: cellular and molecular responses to depressed metabolism and low temperature. *Physiol. Rev.* **83**, 1153–1181.

Champagne, F. A., Weaver, I. C. G., Diorio, J., Dymov, S., Szyf, M. and Meaney, M. J. (2006). Maternal care associated with methylation of the estrogen receptor- α 1b promoter and estrogen receptor- α expression in the medial preoptic area of female offspring. *Endocrinology* **147**, 2909–2915.

Clark, S. J., Harrison, J., Paul, C. L. and Frommer, M. (1994). High sensitivity mapping of methylated cytosines. *Nucleic Acids Res.* **22**, 2990–2997.

Cornacchia, E., Golbus, J., Maybaum, J., Strahler, J., Hanash, S. and Richardson, B. (1988). Hydralazine and procainamide inhibit T cell DNA methylation and induce autoreactivity. *J. Immunol.* **140**, 2197–2200.

Epperson, L. E., Dahl, T. A. and Martin, S. L. (2004). Quantitative analysis of liver protein expression during hibernation in the golden-mantled ground squirrel. *Mol. Cell. Proteomics* **3**, 920–933.

Espada, J., Ballestar, E., Santoro, R., Fraga, M. F., Villar-Garea, A., Nemeth, A., Lopez-Serra, L., Roperio, S., Aranda, A., Orozco, H. et al. (2007). Epigenetic disruption of ribosomal RNA genes and nucleolar architecture in DNA methyltransferase 1 (Dnmt1) deficient cells. *Nucleic Acids Res.* **35**, 2191–2198.

Fedorov, V. B., Goropashnaya, A. V., Toien, O., Stewart, N. C., Gracey, A. Y., Chang, C., Qin, S., Perte, G., Quackenbush, J., Showe, L. C. et al. (2009). Elevated expression of protein biosynthesis genes in liver and muscle of hibernating black bears (*Ursus americanus*). *Physiol. Genomics* **37**, 108–118.

Feinberg, A. P. and Vogelstein, B. (1983). Hypomethylation distinguishes genes of some human cancers from their normal counterparts. *Nature* **301**, 89–92.

Frerichs, K. U., Smith, C. B., Brenner, M., DeGracia, D. J., Krause, G. S., Marrone, L., Dever, T. E. and Hallenbeck, J. M. (1998). Suppression of protein synthesis in brain during hibernation involves inhibition of protein initiation and elongation. *Proc. Natl. Acad. Sci. USA* **95**, 14511–14516.

Fujii, G., Nakamura, Y., Tsukamoto, D., Ito, M., Shiba, T. and Takamatsu, N. (2006). CpG methylation at the USF-binding site is important for the liver-specific transcription of the chipmunk HP-27 gene. *Biochem. J.* **395**, 203–209.

Garzon, R., Liu, S., Fabbri, M., Liu, Z., Heaphy, C. E. A., Callegari, E., Schwind, S., Pang, J., Yu, J., Muthusamy, N. et al. (2009). MicroRNA-29b induces global DNA hypomethylation and tumor suppressor gene reexpression in acute myeloid leukemia by targeting directly DNMT3A and 3B and indirectly DNMT1. *Blood* **113**, 6411–6418.

Green, C. J., Brosnan, J. T., Fuller, B. J., Lowry, M., Stubbs, M. and Ross, B. D. (1984). Effect of hibernation on liver and kidney metabolism in 13-lined ground squirrels. *Comp. Biochem. Physiol. B* **79**, 167–171.

Gunther, K., Rust, M., Leers, J., Boettger, T., Scharfe, M., Jarek, M., Bartkuhn, M. and Renkawitz, R. (2013). Differential roles for MBD2 and MBD3 at methylated CpG islands, active promoters and binding to exon sequences. *Nucleic Acids Res.* **41**, 3010–3021.

Guo, J. U., Su, Y., Zhong, C., Ming, G.-L. and Song, H. (2011). Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain. *Cell* **145**, 423–434.

Hampton, M., Melvin, R. G., Kendall, A. H., Kirkpatrick, B. R., Peterson, N. and Andrews, M. T. (2011). Deep sequencing the transcriptome reveals seasonal adaptive mechanisms in a hibernating mammal. *PLoS ONE* **6**, e27021.

- Harris, M., Olson, L. and Milsom, W. (2004). The origin of mammalian heterothermy: a case for perpetual youth? In *Life in the Cold: Evolution, Mechanisms, Adaptation, and Application* Vol. 12, 44. Alaska, USA: Institute of Arctic Biology, University of Alaska Fairbanks.
- Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biol.* **14**, R115.
- Jin, B., Ernst, J., Tiedemann, R. L., Xu, H., Sureshchandra, S., Kellis, M., Dalton, S., Liu, C., Choi, J.-H. and Robertson, K. D. (2012). Linking DNA methyltransferases to epigenetic marks and nucleosome structure genome-wide in human tumor cells. *Cell Rep.* **2**, 1411-1424.
- Jurkowski, T. P. and Jeltsch, A. (2011). On the evolutionary origin of eukaryotic DNA methyltransferases and Dnmt2. *PLoS ONE* **6**, e28104.
- Kagiyama, S., Kurimoto, K., Hirota, T., Yamaji, M. and Saitou, M. (2013). Replication-coupled passive DNA demethylation for the erasure of genome imprints in mice. *EMBO J.* **32**, 340-353.
- Kang, E. S., Park, C. W. and Chung, J. H. (2001). Dnmt3b, de novo DNA methyltransferase, interacts with SUMO-1 and Ubc9 through its N-terminal region and is subject to modification by SUMO-1. *Biochem. Biophys. Res. Commun.* **289**, 862-868.
- Karimi, M., Johansson, S. and Ekstrom, T. J. (2006a). Using LUMA: a Luminometric-based assay for global DNA-methylation. *Epigenetics* **1**, 45-48.
- Karimi, M., Johansson, S., Stach, D., Corcoran, M., Grandér, D., Schalling, M., Bakalkin, G., Lyko, F., Larsson, C. and Ekstrom, T. J. (2006b). LUMA (Luminometric Methylation Assay)—a high throughput method to the analysis of genomic DNA methylation. *Exp. Cell Res.* **312**, 1989-1995.
- Klose, R. J. and Bird, A. P. (2006). Genomic DNA methylation: the mark and its mediators. *Trends Biochem. Sci.* **31**, 89-97.
- Klug, M. and Rehli, M. (2006). Functional analysis of promoter CpG-methylation using a CpG-free luciferase reporter vector. *Epigenetics* **1**, 127-130.
- Kohli, R. M. and Zhang, Y. (2013). TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* **502**, 472-479.
- Lee, T.-F., Zhai, J. and Meyers, B. C. (2010). Conservation and divergence in eukaryotic DNA methylation. *Proc. Natl. Acad. Sci. USA* **107**, 9027-9028.
- Lin, D. C., Hershey, J. D., Mattoon, J. S. and Robbins, C. T. (2012). Skeletal muscles of hibernating brown bears are unusually resistant to effects of denervation. *J. Exp. Biol.* **215**, 2081-2087.
- Lister, R., Pelizzola, M., Dowen, R. H., Hawkins, R. D., Hon, G., Tonti-Filippini, J., Nery, J. R., Lee, L., Ye, Z., Ngo, Q.-M. et al. (2009). Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* **462**, 315-322.
- Lu, L.-J. W., Randerath, E. and Randerath, K. (1983). DNA hypomethylation in Morris hepatomas. *Cancer Lett.* **19**, 231-239.
- Ma, D. K., Jang, M.-H., Guo, J. U., Kitabatake, Y., Chang, M.-L., Pow-Anpongkul, N., Flavell, R. A., Lu, B., Ming, G.-L. and Song, H. (2009). Neuronal activity-induced Gadd45b promotes epigenetic DNA demethylation and adult neurogenesis. *Science* **323**, 1074-1077.
- Maekawa, F., Shimba, S., Takumi, S., Sano, T., Suzuki, T., Bao, J., Ohwada, M., Ehara, T., Ogawa, Y. and Nohara, K. (2012). Diurnal expression of Dnmt3b mRNA in mouse liver is regulated by feeding and hepatic clockwork. *Epigenetics* **7**, 1046-1056.
- Malatesta, M., Perdoni, F., Battistelli, S., Muller, S. and Zancanaro, C. (2009). The cell nuclei of skeletal muscle cells are transcriptionally active in hibernating edible dormice. *BMC Cell Biol.* **10**, 19.
- Malatesta, M., Zancanaro, C. and Biggiogera, M. (2011). Immunoelectron microscopic characterization of nucleolus-associated domains during hibernation. *Microsc. Res. Tech.* **74**, 47-53.
- Mamady, H. and Storey, K. B. (2008). Coping with the stress: expression of ATF4, ATF6, and downstream targets in organs of hibernating ground squirrels. *Arch. Biochem. Biophys.* **477**, 77-85.
- McMullen, D. C. and Hallenbeck, J. M. (2010). Regulation of Akt during torpor in the hibernating ground squirrel, *Ictidomys tridecemlineatus*. *J. Comp. Physiol. B* **180**, 927-934.
- Montesano, A., Luzi, L., Senesi, P. and Terruzzi, I. (2013). Modulation of cell cycle progression by 5-azacytidine is associated with early myogenesis induction in murine myoblasts. *Int. J. Biol. Sci.* **9**, 391-402.
- Morin, P., Jr and Storey, K. B. (2006). Evidence for a reduced transcriptional state during hibernation in ground squirrels. *Cryobiology* **53**, 310-318.
- Motohashi, N. and Asakura, A. (2012). Molecular regulation of muscle satellite cell self-renewal. *J. Stem Cell Res. Ther. Suppl.* **11**, e002.
- Nan, X., Meehan, R. R. and Bird, A. (1993). Dissection of the methyl-CpG binding domain from the chromosomal protein MeCP2. *Nucleic Acids Res.* **21**, 4886-4892.
- Ng, H.-H., Zhang, Y., Hendrich, B., Johnson, C. A., Turner, B. M., Erdjument-Bromage, H., Tempst, P., Reinberg, D. and Bird, A. (1999). MBD2 is a transcriptional repressor belonging to the MeCP1 histone deacetylase complex. *Nat. Genet.* **23**, 58-61.
- Nur, I., Szyf, M., Razin, A., Glaser, G., Rottem, S. and Razin, S. (1985). Prokaryotic and eucaryotic traits of DNA methylation in spiroplasmas (mycoplasmas). *J. Bacteriol.* **164**, 19-24.
- Osborne, P. G., Gao, B. and Hashimoto, M. (2004). Determination in vivo of newly synthesized gene expression in hamsters during phases of the hibernation cycle. *Jpn. J. Physiol.* **54**, 295-305.
- Palacios, D., Summerbell, D., Rigby, P. W. J. and Boyes, J. (2010). Interplay between DNA methylation and transcription factor availability: implications for developmental activation of the mouse Myogenin gene. *Mol. Cell. Biol.* **30**, 3805-3815.
- Potthoff, M. J. and Olson, E. N. (2007). MEF2: a central regulator of diverse developmental programs. *Development* **134**, 4131-4140.
- Potthoff, M. J., Arnold, M. A., McAnally, J., Richardson, J. A., Bassel-Duby, R. and Olson, E. N. (2007a). Regulation of skeletal muscle sarcomere integrity and postnatal muscle function by Mef2c. *Mol. Cell. Biol.* **27**, 8143-8151.
- Potthoff, M. J., Wu, H., Arnold, M. A., Shelton, J. M., Backs, J., McAnally, J., Richardson, J. A., Bassel-Duby, R. and Olson, E. N. (2007b). Histone deacetylase degradation and MEF2 activation promote the formation of slow-twitch myofibers. *J. Clin. Invest.* **117**, 2459-2467.
- Razin, A. (1998). CpG methylation, chromatin structure and gene silencing—a three-way connection. *EMBO J.* **17**, 4905-4908.
- Razin, A. and Riggs, A. D. (1980). DNA methylation and gene function. *Science* **210**, 604-610.
- Razin, A. and Szyf, M. (1984). DNA methylation patterns formation and function. *Biochim. Biophys. Acta* **782**, 331-342.
- Razin, A., Webb, C., Szyf, M., Yisraeli, J., Rosenthal, A., Naveh-Many, T., Sciaky-Gallili, N. and Cedar, H. (1984). Variations in DNA methylation during mouse cell differentiation in vivo and in vitro. *Proc. Natl. Acad. Sci. USA* **81**, 2275-2279.
- Ruby, N. F., Dark, J., Burns, D. E., Heller, H. C. and Zucker, I. (2002). The suprachiasmatic nucleus is essential for circadian body temperature rhythms in hibernating ground squirrels. *J. Neurosci.* **22**, 357-364.
- Siegfried, Z., Eden, S., Mendelsohn, M., Feng, X., Tsuberi, B. Z. and Cedar, H. (1999). DNA methylation represses transcription in vivo. *Nat. Genet.* **22**, 203-206.
- Srere, H. K., Wang, L. C. and Martin, S. L. (1992). Central role for differential gene expression in mammalian hibernation. *Proc. Natl. Acad. Sci. USA* **89**, 7119-7123.
- Storey, K. B. (1987). Regulation of liver metabolism by enzyme phosphorylation during mammalian hibernation. *J. Biol. Chem.* **262**, 1670-1673.
- Storey, K. B. and Storey, J. M. (2010). Metabolic rate depression: the biochemistry of mammalian hibernation. *Adv. Clin. Chem.* **52**, 77-108.
- Tajerian, M., Alvarado, S., Millicamps, M., Vachon, P., Crosby, C., Bushnell, M. C., Szyf, M. and Stone, L. S. (2013). Peripheral nerve injury is associated with chronic, reversible changes in global DNA methylation in the mouse prefrontal cortex. *PLoS ONE* **8**, e55259.
- Tessier, S. N. and Storey, K. B. (2010). Expression of myocyte enhancer factor-2 and downstream genes in ground squirrel skeletal muscle during hibernation. *Mol. Cell. Biochem.* **344**, 151-162.
- Tessier, S. N. and Storey, K. B. (2012). Myocyte enhancer factor-2 and cardiac muscle gene expression during hibernation in thirteen-lined ground squirrels. *Gene* **501**, 8-16.
- Tessier, S. N. and Storey, K. B. (2014). To be or not to be: the regulation of mRNA fate as a survival strategy during mammalian hibernation. *Cell Stress Chaperones* **19**, 763-776.
- Tessier, S. N., Audas, T. E., Wu, C.-W., Lee, S. and Storey, K. B. (2014). The involvement of mRNA processing factors TIA-1, TIAR, and PABP-1 during mammalian hibernation. *Cell Stress Chaperones* **19**, 813-825.
- Waterland, R. A. and Jirtle, R. L. (2003). Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol. Cell. Biol.* **23**, 5293-5300.
- Weaver, I. C. G., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., Dymov, S., Szyf, M. and Meaney, M. J. (2004). Epigenetic programming by maternal behavior. *Nat. Neurosci.* **7**, 847-854.
- Wrobel, G., Kokocinski, F. and Lichter, P. (2004). AutoPrime: selecting primers for expressed sequences. *Genome Biol.* **5**, 11.
- Wu, C.-W. and Storey, K. B. (2012a). Pattern of cellular quiescence over the hibernation cycle in liver of thirteen-lined ground squirrels. *Cell Cycle* **11**, 1714-1726.
- Wu, C.-W. and Storey, K. B. (2012b). Regulation of the mTOR signaling network in hibernating thirteen-lined ground squirrels. *J. Exp. Biol.* **215**, 1720-1727.
- Yan, J., Barnes, B. M., Kohl, F. and Marr, T. G. (2008). Modulation of gene expression in hibernating arctic ground squirrels. *Physiol. Genomics* **32**, 170-181.
- Yasui, D. H., Xu, H., Dunaway, K. W., LaSalle, J. M., Jin, L.-W. and Maezawa, I. (2013). MeCP2 modulates gene expression pathways in astrocytes. *Mol. Autism* **4**, 3.