

## RESEARCH ARTICLE

# Variation in early-life telomere dynamics in a long-lived bird: links to environmental conditions and survival

Hannah Watson<sup>1,\*‡</sup>, Mark Bolton<sup>2</sup> and Pat Monaghan<sup>1</sup>

## ABSTRACT

Conditions experienced during early life can have profound consequences for both short- and long-term fitness. Variation in the natal environment has been shown to influence survival and reproductive performance of entire cohorts in wild vertebrate populations. Telomere dynamics potentially provide a link between the early environment and long-term fitness outcomes, yet we know little about how the environment can influence telomere dynamics in early life. We found that environmental conditions during growth have an important influence on early-life telomere length (TL) and attrition in nestlings of a long-lived bird, the European storm petrel *Hydrobates pelagicus*. Nestlings reared under unfavourable environmental conditions experienced significantly greater telomere loss during postnatal development compared with nestlings reared under more favourable natal conditions, which displayed a negligible change in TL. There was, however, no significant difference in pre-fledging TL between cohorts. The results suggest that early-life telomere dynamics could contribute to the marked differences in life-history traits that can arise among cohorts reared under different environmental conditions. Early-life TL was also found to be a significant predictor of survival during the nestling phase, providing further evidence for a link between variation in TL and individual fitness. To what extent the relationship between early-life TL and mortality during the nestling phase is a consequence of genetic, parental and environmental factors is currently unknown, but it is an interesting area for future research. Accelerated telomere attrition under unfavourable conditions, as observed in this study, might play a role in mediating the effects of the early-life environment on later-life performance.

**KEY WORDS:** Telomere length, Life-history evolution, Senescence, Telomere dynamics, Cohort effects

## INTRODUCTION

One of the principal aims of evolutionary ecology is to understand the mechanisms underlying individual variation in longevity and fecundity. Telomere dynamics link cellular processes with organismal ageing and thus optimisation of telomere length (TL) and attrition may play a major role in life-history evolution

(Monaghan and Haussmann, 2006). Telomeres comprise highly conserved non-coding DNA sequences that form protective caps at the ends of eukaryotic chromosomes (Blackburn, 2005). By protecting coding sequences from attrition, telomeres play an important role in maintaining genome stability (reviewed by Verdun and Karlseder, 2007). In the absence of the enzyme telomerase, telomeres shorten with each round of somatic cell division; when a critical length is reached, telomeres become dysfunctional and cells enter a state of replicative senescence (Blackburn, 2005; Verdun and Karlseder, 2007). Senescent cells subsequently die or adopt an altered secretory profile, resulting in secretion of inflammatory cytokines, growth factors and degradative enzymes that contribute to age-related declines in tissue and organ function (Campisi, 2005). The accumulation of senescent cells thus appears to be important to the ageing phenotype, thereby influencing lifespan (Campisi, 2005).

Longitudinal studies have shown that TL declines progressively with age in many vertebrates (e.g. Zeichner et al., 1999; Brummendorf and Mak, 2002; Salomons et al., 2009; Aviv et al., 2009; Bize et al., 2009; Heidinger et al., 2012; Barrett et al., 2013). Large within-species variability in TL and the rate of telomere shortening is reported among individuals of the same age (Hall et al., 2004; Aviv et al., 2009; Bize et al., 2009). Inter-individual variation in TL has been found to predict fitness components in natural populations; individuals with the shortest telomeres or the highest loss rate have the poorest survival prospects (Cawthon et al., 2003; Haussmann et al., 2005; Bize et al., 2009; Olsson et al., 2011; Heidinger et al., 2012) and TL was found to be positively correlated with lifetime reproductive success (Pauliny et al., 2006). Although early-life TL is partly determined by genetic factors (Njajou et al., 2007; Olsson et al., 2011), much of the inter-individual variation in TL may relate to environmental influences. Exposure to repeated stress (Epel et al., 2004; Kotrschal et al., 2007; Herborn et al., 2014), large-scale climatic processes (Mizutani et al., 2013), low habitat quality (Angelier and Vleck, 2013) and reproduction (Kotrschal et al., 2007; Heidinger et al., 2012) have all been associated with accelerated telomere attrition, possibly mediated by increased oxidative damage (von Zglinicki, 2002). TL therefore potentially reflects variation in individual state and past experiences.

It is well known that early-life conditions can have profound influences on phenotypic development and long-term fitness consequences (Lindström, 1999; Metcalfe and Monaghan, 2001). Variation in the pre- and early post-natal environment has been shown to influence survival and reproductive performance in a number of vertebrate species (Albon et al., 1987; Haywood and Perrins, 1992; Sedinger et al., 1995; Rose et al., 1998; Reid et al., 2003). Environmental conditions during early life can affect an entire cohort simultaneously, giving rise to substantial differences in life histories between successive cohorts, which can persist throughout the cohort's lifespan (Albon et al., 1987; Rose et al.,

<sup>1</sup>Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow G12 8QQ, UK. <sup>2</sup>RSPB Centre for Conservation Science, UK Headquarters, The Lodge, Sandy, Bedfordshire SG19 2DL, UK.

\*Present address: Department of Biology, Ecology Building, Lund University, Lund 223 62, Sweden.

‡Author for correspondence (hannah.watson@biol.lu.se)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

**List of abbreviations**

AUC	area under the curve
$C_t$	the number of PCR cycles required for accumulation of sufficient products to exceed a set threshold of fluorescent signal
GLM	generalised linear model
LM	linear model
LMM	linear mixed-effects model
LRT	likelihood ratio test
OCD	a non-variable copy control gene isolated from the European storm petrel encoding the protein ornithine decarboxylase
qPCR	quantitative PCR
REML	restricted maximum likelihood estimation
$T/S$	ratio of telomeric sequence to that of a single-copy gene, relative to the reference sample
TL	telomere length
$\Delta$ age	change in age (days) between first and second telomere length measurements

1998; Reid et al., 2003). Several studies on mammalian and avian species have found that telomere loss is greatest during early life, presumably as a result of the rapid growth and cell division that occurs during this period (Zeichner et al., 1999; Baerlocher et al., 2007; Salomons et al., 2009). Poor early nutrition and catch-up growth have been shown to result in accelerated telomere loss in a number of studies, including laboratory rats (*Rattus norvegicus*: Jennings et al., 1999; Tarry-Adkins et al., 2009) and wild birds (European shag *Phalacrocorax aristotelis*: Hall et al., 2004; king penguin *Aptenodytes patagonicus*: Geiger et al., 2012). Developmental stress has also been shown to accelerate telomere shortening through experimental brood enlargement (jackdaw *Corvus monedula*: Boonekamp et al., 2014) and increased exposure of nestlings to glucocorticoids (European shag: Herborn et al., 2014) in the wild. Early-life TL was found to be a better predictor of longevity than TL in adulthood in a longitudinal study of captive zebra finches *Taeniopygia guttata* (Heidinger et al., 2012). Early-life telomere dynamics may therefore mechanistically link developmental conditions with later-life senescence (Monaghan, 2010). Despite the potential significance of early-life telomere dynamics in influencing life-history traits, we still know relatively little about how variation in early-life conditions influences telomere loss during development in natural environments.

In this study, we examined the effects of inter-annual variation in the natal environment on TL and telomere dynamics during postnatal development in the European storm petrel *Hydrobates pelagicus* Linnaeus 1758 (hereafter, storm petrel). The storm petrel is a remarkably long-lived seabird, displaying low annual fecundity (obligate clutch of one) and low adult mortality. Its long lifespan may in part be attributed to superior cellular mechanisms for resistance to oxidative damage (Ogburn et al., 2001) and maintenance of telomere length (Hausmann et al., 2007). The oldest individuals in a population of a related species, the Leach's storm petrel *Oceanodroma leucorhoa*, show little or no accumulation of short telomeres over time (Hausmann and Mauck, 2008). We compared cohorts from two consecutive breeding seasons characterised by different environmental conditions, as indicated by overall reproductive performance of the colony. Productivity was relatively poor in 2011 and significantly lower than 2010 (Watson et al., 2014). This provided the opportunity to examine the effects of the natal environment on telomere length and dynamics during early life and relate these to nestling survival. The study site and design also enabled us to investigate whether exposure to an additional

source of developmental stress, arising from human recreational disturbance, has consequences for early-life telomere dynamics. Parent–offspring relationships in TL were examined within a single cohort.

**RESULTS**

The within-individual change in nestling TL was significantly different between the two cohorts raised under different environmental conditions: in 2010, there was little change in TL during the nestling period, whereas TL declined significantly with age in nestlings reared in 2011 (Table 1, model 1; Fig. 1A;  $\Delta$  age $\times$ cohort:  $F_{1,55,2}=12.17$ ,  $P\leq 0.001$ ). However, TL in early postnatal development was significantly longer in individuals from the 2011 cohort, compared with the 2010 cohort (Table 1, model 1;  $F_{1,94,3}=6.25$ ,  $P=0.014$ ). Removal of the main effect of cohort, while retaining it in the interaction with  $\Delta$  age (change in age in days between first and second telomere length measurements), showed that, even after adjusting for early differences in TL, the rate of change in TL was still significantly different between the two cohorts ( $\beta=-0.005\pm 0.002$ ,  $F_{1,95,5}=5.82$ ,  $P=0.018$ ) and therefore the cohort effect was above and beyond any effect due to regression to the mean. Furthermore, there was considerable variation between individuals in the degree of change in TL, with some individuals even exhibiting an apparent increase in TL during postnatal development (Fig. 1B). The between-individual differences in TL were not quite significant (Table 1, model 1; age at first measurement:  $F_{1,72,6}=3.81$ ,  $P=0.055$ ). Neither nestling TL nor the within-individual change in TL (tested by the respective interactions with  $\Delta$  age) was affected by any of the other variables considered, including sex, hatching date or visitor disturbance (all  $P>0.2$ ). Analysis of TL in late postnatal development demonstrated that, although there was a tendency for late TL to be shorter in 2011, the estimated cohort effect was not quite significant at this stage (Table 1, model 2;  $t_{47}=-1.73$ ,  $P=0.089$ ).

Of the 61 nestlings sampled, 6 died prior to fledging (3 individuals from each cohort). When examining whether early TL was a good predictor of survival of the nestling phase, we found that the probability of surviving to fledging increased significantly with TL (Table 1, model 3; Fig. 2A;  $z_{39}=2.12$ ,  $P=0.034$ ). The mean TL of chicks that did not survive to fledging was 22% shorter than that of chicks that successfully fledged (Fig. 2B). The model fit was significantly improved by the inclusion of the rate of mass gain, though this variable did not explain a significant portion of the variation in probability of fledging (Table 1, model 3;  $z_{39}=1.76$ ,  $P=0.078$ ). One of the chicks that did not fledge had very short telomeres ( $T/S$  ratio=0.68) and so the analysis was re-run excluding this nestling. The outlier had no influence on the relationship, its removal having no effect on the magnitude or significance of the effect of TL ( $\beta=0.99$ , 95% CI: 0.64–1.0,  $z_{38}=2.08$ ,  $P=0.038$ ). The probability of fledging was not affected by sex, hatching date, cohort or visitor disturbance (all  $P>0.1$ ). The relationship between nestling TL and probability of fledging was not affected by rate of mass growth (Table 1, model 3; TL $\times$ growth rate:  $z_{38}=-1.07$ ,  $P=0.286$ ). The AUC of the minimum adequate model was 0.90, indicating that the model performed very well in terms of accuracy.

Analysis of parent–offspring relationships in the 2011 cohort revealed that nestling TL was not significantly correlated with either maternal TL (Table 1, model 4;  $t_{19}=0.63$ ,  $P=0.538$ ) or paternal TL (Table 1, model 4;  $t_{19}=1.0$ ,  $P=0.330$ ). Parent–offspring TL relationships were unaffected by nestling sex, as shown by testing the respective two-way interactions (all  $P>0.5$ ).

**Table 1. Variation in telomere length (TL) and rate of telomere attrition during postnatal development in relation to natal environmental conditions, survival and parental TL**

Model and dependent variable	Fixed effects	Rejected terms	Parameter estimate (s.e. or 95% CI)	P-value
(1) Within-individual change in telomere length during postnatal development (LMM <sup>†</sup> )				
All TL (N=100)	$\Delta$ Age <sup>§</sup>		3.11e <sup>-4</sup> (1.90e <sup>-3</sup> )	<0.001***
	Cohort (2011)		0.147 (0.059)	0.014*
	$\Delta$ Age $\times$ cohort (2011)		-9.05e <sup>-3</sup> (2.59e <sup>-3</sup> )	<0.001***
		Age at first measurement	6.09e <sup>-3</sup> (3.12e <sup>-3</sup> )	0.055
		All 2-way interactions with $\Delta$ age		>0.2
(2) Telomere length in late postnatal development (LM)				
Late TL (N=49)		Cohort (2011)	-0.092 (0.053)	0.089
		Age	0.002 (0.004)	0.640
(3) Nestling survival and telomere length (binomial GLM)				
Survival (0/1) <sup>  </sup> (N=42)	Early TL		0.99 (0.68–1.0)	0.034*
	Growth rate		0.94 (0.42–1.0)	0.078
		Cohort (2011)	0.64 (0.15–0.95)	0.614
		TL $\times$ growth rate	5.2e <sup>-7</sup> (1.5e <sup>-18</sup> –1.0)	0.286
(4) Parent–offspring relationship in telomere length in 2011 cohort (LM)				
Early TL (N=21)		Maternal TL	0.148 (0.237)	0.538
		Paternal TL	0.194 (0.194)	0.330
		Maternal TL $\times$ nestling sex (male)	-0.301 (0.506)	0.560
		Paternal TL $\times$ nestling sex (male)	-0.139 (0.593)	0.818

TL of nestlings was measured in early and late development in two successive cohorts reared under favourable (2010) and unfavourable (2011) conditions. Parental TL was only measured in the 2011 cohort. Parameter estimates and P-values are shown for fixed effects retained in minimum adequate models and for key terms that were rejected during forward stepwise regression. Models 1 and 2 also tested for the fixed effects of sex, hatching date and visitor disturbance; the effects were not significant in any model and, for simplicity, are not shown in the table. Model type is specified as either linear mixed model (LMM), linear model (LM) or generalised linear model (GLM).

<sup>†</sup>Included random intercept for nestling identity; mean variance $\pm$ s.d., 0.005 $\pm$ 0.07.

<sup>§</sup> $\Delta$  age, change in age (days) between TL measurements.

<sup>||</sup>0=died prior to fledging, 1=survived to fledging.

\* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.001.


## DISCUSSION

Although it has been widely suggested that the most rapid telomere loss occurs early in life in a range of vertebrates (Zeichner et al., 1999; Baerlocher et al., 2007; Salomons et al., 2009), only a few studies have investigated telomere dynamics within the timeframe of postnatal development itself (see Foote et al., 2011; Geiger et al., 2012; Boonekamp et al., 2014; Herborn et al., 2014). We found that TL and dynamics in growing nestlings of a long-lived species were strongly influenced by the quality of the natal environment. Favourable environmental conditions during development, as reflected by moderate colony productivity in 2010, were associated with a negligible change in TL during early life, whereas conditions resulting in significantly lower chick survival among the 2011 cohort were associated with significant telomere attrition. Whereas human recreational disturbance was previously shown to reduce survival probabilities of nestlings (Watson et al., 2014), there was no effect of visitor activity on telomere length or dynamics.

While some of the cohort effect might be explained by regression to the mean, the difference in rate of change in TL remained, even after adjusting for the differences in early TL between the two cohorts. Accelerated attrition of longer telomeres, having controlled for regression to the mean, has previously been reported (Salomons et al., 2009; Aviv et al., 2009) and could be explained if longer telomeres offer a larger area for attack by free radicals (Aviv et al., 2009). The high early TL in the 2011 cohort could potentially be explained if the high early mortality observed within the 2011 cohort removed individuals with short telomeres from the population or if poor-quality and/or inexperienced birds did not breed. Early selective mortality of individuals with short telomeres was previously predicted to occur in storm petrels (Hausmann and

Mauck, 2008). Environmental conditions could be driving the accelerated telomere attrition observed in the 2011 cohort either directly, for example, via elevated oxidative stress, and/or indirectly, if selective mortality or breeding by only high-quality individuals were to result in a higher mean early TL, which is then subject to increased attrition. The functional links between the environment and telomere dynamics could be mediated by changes in parental care. Using a larger dataset from the same population, we found that, in 2011, parents spent less time brooding their young and maximal growth rate (in respect of mass) occurred later (H.W., unpublished results). An increased frequency of extreme weather events in 2011 (Watson, 2014) might have been a key factor influencing breeding success and parental care. Irrespective of the mechanism(s), this study presents good evidence for environmental conditions influencing the rate of telomere loss and shaping variation in early-life telomere dynamics between cohorts.

It is well known that cohort effects can arise in response to natal conditions. Differences in survival and recruitment rates among cohorts have previously been shown to be positively correlated with the natal environment in mammals (Albon et al., 1987; Rose et al., 1998) and birds (Sedinger et al., 1995; Reid et al., 2003). An increasing body of evidence links short telomeres and accelerated rate of telomere loss to a shorter lifespan (Cawthon et al., 2003; Bize et al., 2009; Olsson et al., 2011; Heidinger et al., 2012) and lower reproductive success (Pauliny et al., 2006). The results of this study suggest that early-life telomere dynamics may contribute to the marked differences in life-history traits that can arise among cohorts reared under different environmental conditions. Indeed, TL in early life has been shown to be a strong predictor of lifespan and a better predictor than TL in adulthood (Heidinger et al., 2012). In addition to the increase in chick mortality observed in the cohort reared under




**Fig. 1. Change in telomere length during growth in storm petrel nestlings from two consecutive cohorts reared under different natal conditions.**

(A) Mean within-individual change in TL in 2010 (black circles; solid line) and 2011 (open circles; dashed line). Lines represent model predictions from the minimum adequate LMM ( $\Delta \text{age} \times \text{cohort}$  (2011):  $F_{1,55.2}=12.17$ ,  $P \leq 0.001$ ,  $N=100$ ) fitted within the range of observed values.  $\Delta \text{age}$  is the change in age (days) between the first and second measurement. (B) Individual change in telomere length during postnatal development ( $N=98$ ) in 2010 (black circles; solid lines) and 2011 (open circles; dashed lines). Lines link TL measurements in early and late postnatal development for each individual.

unfavourable natal conditions, the population-level effects might be exacerbated by a reduction in the fitness of entire cohorts compared with those fledging under more favourable conditions. Such cohort-wide effects can subsequently destabilise population dynamics (Lindström, 1999). Although our understanding will benefit from examination of longer-term data of multiple cohorts, this study supports the idea that telomere dynamics may be a mechanism linking early-life conditions with later-life performance.

The maintenance of TL during postnatal development under favourable conditions in storm petrels contrasts with studies in other birds that suggest rapid telomere loss invariably occurs early in life (Hall et al., 2004; Baerlocher et al., 2007; Salomons et al., 2009; Heidinger et al., 2012). A negligible change in early-life TL was also demonstrated in chicks of the long-lived king penguin experiencing favourable growth conditions (Geiger et al., 2012). A slow rate of telomere shortening could be causally linked to a higher resistance to oxidative stress (Ogburn et al., 2001) and/or elevated telomerase activity (Verdun and Karlseder, 2007; Haussmann et al., 2007). Stressful conditions, however, have been shown to lead to accelerated telomere loss via downregulation



**Fig. 2. Relationship between early nestling telomere length and survival during postnatal development.** (A) Predicted probability of fledging (solid line) in relation to early TL ( $\leq 16$  days; GLM:  $z_{39}=2.12$ ,  $P=0.034$ ) with 95% confidence intervals (dashed lines). Line is fitted within the range of observed values (circles;  $N=42$ ) and based on a mean rate of mass gain of  $1.03 \text{ g day}^{-1}$  ( $z_{39}=1.76$ ,  $P=0.078$ ). Removal of the outlier at the lower end of the TL scale did not affect model predictions. (B) Mean  $\pm$  s.e. early TL of nestlings that died ( $N=6$ ) during the nestling phase and those that survived to fledging ( $N=36$ ).

of antioxidants in birds (Stier et al., 2009) and telomerase in humans (Epel et al., 2004). When faced with a costly reproductive event, adult Adélie penguins *Pygoscelis adeliae* are able to increase antioxidant defences and avoid accelerated telomere attrition (Beaulieu et al., 2011). The results of this study suggest that storm petrel nestlings, when faced with unfavourable natal conditions, are unable to modulate or activate such regulatory mechanisms. If poor early conditions were to lead to irreversible downregulation of antioxidants or telomerase, accelerated telomere loss may even persist beyond the nestling period, exacerbating the effects of early conditions on later-life senescence. An area for future research is to understand the links between telomere loss, oxidative stress and telomerase activity during early life.

While recent studies have investigated the relationship between TL in early life and adult survival in captive (Heidinger et al., 2012) and wild (Caprioli et al., 2013; Boonekamp et al., 2014) birds, we are aware of only one study that has directly examined the association between early-life TL and mortality within the development phase itself. Just as king penguin chicks that died prior to fledging displayed shorter telomeres (Geiger et al., 2012), we also found that post-hatching TL of storm petrel nestlings was directly related to imminent mortality. Despite a small sample size,



the model performed well (as measured by AUC) and the results were not influenced by an outlying observation. Although confidence intervals are wide at low TLs, they do not overlap with confidence intervals when the probability of fledging is equal to one. TL was a much better predictor of fledging success than the rate of mass gain. Although food load and rate of mass gain are likely to be of importance in determining survival within the first few days of hatching, storm petrel nestlings are well-buffered against periodic food shortages and mortality beyond the brooding stage is unlikely to be strongly influenced by environmental factors affecting rate of provisioning and subsequent mass gain. Additionally, because they nest in cavities, the risk of nest predation is very low. Consequently, it seems most likely that mortality was linked to intrinsic, rather than extrinsic, factors. Accelerated telomere shortening at this stage is likely to be a biomarker rather than a cause of poor survival prospects and may be indicative of exposure to high levels of glucocorticoids and/or oxidative stress. Early TL is partly determined by genetic factors (Njajou et al., 2007), but the rate of attrition is also known to be affected by environmental influences (Epel et al., 2004; Tarry-Adkins et al., 2009; Geiger et al., 2012). The results demonstrate that early-life TL is a good predictor of imminent mortality.

Recent evidence suggests that TL in birds is maternally inherited (an example of heterogametic inheritance) (Horn et al., 2011; Reichert et al., 2015; Asghar et al., 2015). However, Reichert et al. (2015) also showed that environmental factors exerted a strong influence on nestling TL and the maternal link disappeared with chick age. The absence of any observable heritable component to early TL in this study may therefore be a consequence of environmental effects obscuring any initial genetic effect on nestling TL. As we only have data on parent–offspring relationships for the cohort reared under unfavourable conditions, parent–offspring relationships in this study could have been obscured by parental age, which was not known; although, telomere length was found to be unrelated to age in two other long-lived avian species (Hall et al., 2004). Future research should seek to disentangle the relative contributions from genetic, parental and environmental effects on TL and dynamics. This will help to develop our understanding of the role of telomere dynamics in driving the evolution of life histories.

## MATERIALS AND METHODS

### Data collection

The study was conducted at the island of Mousa, located in the Shetland archipelago, UK (60°0'N, 1°10'W). Storm petrel nestlings are brooded for ~7 days and do not leave the underground nest cavity until fledging at ~65–70 days (Davis, 1957). Fledging success (of eggs hatched) was significantly lower in 2011 (0.60), compared with 2010 (0.78) (see Watson et al., 2014). This was probably linked, in part, to an increased frequency of extreme weather events in 2011 (Watson, 2014). It is not always clear what environmental correlates are relevant, however, and overall breeding performance of a population is widely used as a measure of the natal environment (e.g. Reid et al., 2003). Blood was collected from 32 nestlings in 2010 and 29 nestlings in 2011 and from both parents at 21 nests in 2011 only. Whole blood was obtained by venipuncture of the brachial vein under licence from the UK Home Office. Nestlings were sampled on two occasions (where possible) in the first (hereafter, 'early'; median age: 11 days, range: 4–33 days,  $N=59$ ) and second (hereafter, 'late'; median: 45 days, range: 34–63 days,  $N=52$ ) half of postnatal development. Fifty nestlings were sampled twice and the median number of days elapsed between repeated measurements was 31 days (range: 22–47 days). Only one sample was available for some nestlings, either because nestlings died ( $N=6$ ), were not accessible in the nest or samples were accidentally destroyed. Of the 50 nestlings sampled twice, 49 survived to fledging. The overall proportion of chicks that died was greatly underrepresented in the

data and the difference in mortality between the two cohorts was not reflected within the dataset; this is because the majority of chick mortality occurred within a few days of hatching and before initial blood sampling was carried out. Tarsus length (nestlings only), mass and wing length (maximum flattened chord; adults only) were recorded. Nestling mass was corrected to a standardised time (18:00 h) to account for the age-related rate of proportional weight loss that chicks undergo during diurnal fasting (Bolton, 1995).

### Determination of TL

Blood samples were stored at <5°C for up to 8 h prior to being separated by centrifugation. Plasma and red blood cells were stored at <5°C in the field for a maximum of 3 days before being transferred to –20°C for up to 3 months; following this, samples were stored at –80°C until laboratory analyses were performed. DNA was extracted from red blood cells using Machery–Nagel NucleoSpin Blood kits following the manufacturer's protocol. The quantity and purity of DNA was assessed using a NanoDrop 8000 spectrophotometer (Thermo Scientific); there were no differences in yield or purity between cohorts. Sex was determined following the molecular method described by Griffiths et al. (1998). TL was measured by quantitative PCR (qPCR) using an Mx3005P machine (Stratagene) as described by Cawthon (2002) and adapted by Criscuolo et al. (2009) for birds. The method gives a relative value for TL by determining the ratio of number of telomere repeats ( $T$ ) to that of a single-copy (or, more precisely, non-variable in copy number) control gene ( $S$ ) relative to a reference sample. Consequently, the measure will include any interstitial repeats of the telomeric sequence (Foote et al., 2013). However, good correlations have been found between TL measurements including and excluding the interstitial repeats using telomere restriction fragment (TRF) analysis (Foote et al., 2013) and between TRF and qPCR analyses (Criscuolo et al., 2009; Aviv et al., 2011). Since the extent of interstitial repeats is not expected to change with age, qPCR is well suited to detecting within-individual changes. Furthermore, measurement of telomeres from Leach's storm petrels, using TRF methods, suggests that interstitial repeats do not occur to the extent that they would influence telomere measurements (M. Haussmann, personal communication).

Amplification of telomere sequences was achieved using the forward and reverse primers: Tel1b (5'-CGGTTTGTTGGGTTTGGGTTTGGGTTTGGGTTTGGGTTTGGGTT-3') and Tel2b (5'-GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCTTACCCT-3'). The non-variable copy control gene used was ornithine decarboxylase (OCD), isolated from the European storm petrel (GenBank: DQ881744.1), which was amplified using the primers: OCD Fwd1 (5'-GACCTTGCCATCATTGGAGTTAG-3') and OCD Rev1 (5'-AAGGCATCCCTATTGTTAGGTAGA-3') sourced from Integrated DNA Technologies (Leuven, Belgium). qPCR was performed using 10 ng DNA per reaction. The concentrations of primers used were 500 nmol l<sup>-1</sup> for telomere and 70 nmol l<sup>-1</sup> for OCD reactions. Telomere qPCR reaction conditions started with 15 min at 95°C, followed by 27 cycles of 15 s at 95°C, 30 s annealing at 58°C and 30 s extending at 72°C. OCD reaction conditions started with 15 min at 95°C, followed by 40 cycles of 30 s at 95°C and 30 s at 60°C. Melting curves were assessed to confirm there was no non-specific amplification or primer–dimer formation. For both telomere and OCD reactions, the number of PCR cycles required for accumulation of sufficient products to exceed a threshold of fluorescent signal ( $C_t$ ) was determined. A standard curve, run on each plate, consisted of a serial dilution of a reference sample ranging from 40 ng to 2.5 ng. The  $C_t$  threshold for each reaction was determined from the reference sample. A negative control was also included on each plate. All samples, including the standard curve, were run in triplicate. Amplification efficiencies were within an acceptable range (mean±s.d.: telomere, 106.8±3.7%; OCD, 114.3±3.9%) and all samples fell within the bounds of the standard curve. Inter- and intra-assay variations were 3.6% and 1.4%, respectively, for telomere reactions and 0.82% and 0.33%, respectively, for OCD reactions. Inter-assay variation in  $T/S$  ratio was 10.4%. The  $T/S$  ratio for each sample was calculated from mean  $C_t$  values according to Pfaffl (2001). This method accounts for the variation in amplification efficiencies. All repeated samples from nestlings were run on the same plate along with parental samples.

### Statistical analysis

All statistical analyses were performed in R 3.0.0 (R Core Team, 2013). Linear mixed models (LMMs) with a normal error structure were fitted to data on nestling telomere length to examine change in TL in surviving nestlings ( $N=100$ ; number of individuals=55). Outlying observations were removed to normalise the data and improve model fits. Because we were primarily interested in the within-individual change in nestling TL, we used within-subject centring (van de Pol and Verhulst, 2006) to separate between-individual (cross-sectional) and within-individual (longitudinal) effects. The age variable was split into two covariates: age at first measurement (between-individual effects) and  $\Delta$  age (the change in age between measurements; within-individual effects). LMMs were fitted in the lme4 package (Bates et al., 2013). First, the optimal random effects structure was found by comparing nested models, fitted by restricted maximum likelihood (REML), using likelihood ratio tests (LRTs). A random intercept for nestling identity was included. We also considered the inclusion of a random intercept for nest identity, to account for potential non-independence of nestlings from different cohorts reared at the same nest-site. Thirty per cent of nests were sampled in both years, although we do not know whether nests were occupied by the same breeding pair in both years. However, the variance associated with nest identity was estimated to be zero and inclusion of the random effect did not significantly improve model fits. With nestling TL as the dependent variable, candidate explanatory variables included cohort (year: 2010 or 2011), nestling sex and human disturbance (high or low, defined as  $\leq 10$  m or  $> 150$  m from the visitor path, respectively) as two-level fixed factors, and age at first measurement,  $\Delta$  age and hatching date as covariates. We tested for effects on within-individual change in TL by considering all respective interactions with  $\Delta$  age. To examine variation in nestling TL in late development, linear models (LMs) with a normal error structure were fitted to log-transformed late nestling TL. Candidate explanatory variables included cohort, nestling sex and human disturbance as fixed factors and age and hatching date as covariates.

Generalised linear models (GLMs) with a binomial error structure and logit link function were fitted to data on survival (0/1;  $N=42$ ) to examine whether early TL (age:  $\leq 16$  days, median=11 days) was a good predictor of survival of the nestling phase. The threshold of 16 days was adopted because this was the upper limit of the 95% confidence interval of age at first measurement. These data did not include any repeated measures at the nest level. In addition to the fixed effect of nestling TL at  $\leq 16$  days, candidate explanatory variables included cohort, sex and visitor disturbance as fixed factors and body size, body mass, hatching date, growth rate (average daily incremental growth in mass) as covariates. We also tested for two-way interactions between TL and mass growth rate. The minimum adequate model was evaluated using receiver operating characteristic plots (Sing et al., 2005). The resulting area under the curve (AUC) offers a measure of predictive performance for a binomial model; a value of 1.0 indicates a perfect model, whereas a value of 0.5 indicates that a model performs no better than random.

Finally, parent–offspring relationships in the 2011 cohort were examined in a linear model (LM) analysis with early nestling TL as the dependent variable ( $N=21$ ). Candidate fixed effects included the covariates of maternal TL, paternal TL and nestling sex as a fixed factor. We also tested for potential effects of nestling sex on parent–offspring relationships with the respective two-way interactions between parental TL and nestling sex. We showed in previous analyses that age, hatching date and visitor disturbance did not affect early nestling TL and these variables were therefore not included in the parent–offspring analysis.

Taking into account the sample sizes and the number of variables to potentially control for, a conservative approach to model fitting was adopted; starting from a null model, fixed effects that significantly differed from zero ( $P < 0.05$ ) were added sequentially in a forward stepwise regression. Each time a new variable was added to the model, the significance of existing variables was re-examined. Nested models were compared using LRTs and the criterion for entry of a variable was a log-likelihood ratio  $P$ -value of  $< 0.05$ . For LMMs, model selection was performed using models fitted by maximum likelihood; parameter estimates with standard errors are quoted from the minimum adequate LMM fitted by REML. The significance of parameter estimates was

estimated using conditional  $F$ -tests based on Satterthwaite approximation for the denominator degrees of freedom. For LMs and GLMs, parameter estimates were estimated based on the  $t$ - and  $z$ -distributions, respectively; again estimates with standard errors are quoted from minimum adequate models.

### Acknowledgements

We thank Tom and Cynthia Jamieson for their overwhelming kindness and for running a superb ferry service; Julie Miller and Hannah Taylor for assistance with data collection in the field; Winnie Boner and Robert Gillespie for assistance with validation and optimisation of telomere assays; and, Aileen Adam and Kate Griffiths for assistance with molecular sexing. We also thank RSPB Shetland, Martin Heubeck, Alan Pottinger and Scottish Natural Heritage for logistical support in the field.

### Competing interests

The authors declare no competing or financial interests.

### Author contributions

Conception, study design and interpretation of the findings: H.W., M.B. and P.M. Data collection, laboratory analyses and data analysis: H.W. Preparation of the manuscript: H.W., M.B. and P.M.

### Funding

This study was supported by a DTA Organismal Biology studentship from the Biotechnology and Biological Sciences Research Council [BB/F016700/1 to H.W.] and a European Research Council Advanced Investigator Award [268926 to P.M.]. Additional financial support for fieldwork was provided by the Louise Hiom Trust [to H.W.]. M.B. did not receive any specific funding in respect of this study.

### References

- Albon, S. D., Clutton-Brock, T. H. and Guinness, F. E. (1987). Early development and population dynamics in red deer. II. Density-independent effects and cohort variation. *J. Anim. Ecol.* **56**, 69–81.
- Angelier, F. and Vleck, C. M. (2013). Telomere length, non-breeding habitat and return rate in male American redstarts. *Funct. Ecol.* **27**, 342–350.
- Asghar, M., Bensch, S., Tarka, M., Hansson, B. and Hasselquist, D. (2015). Maternal and genetic factors determine early life telomere length. *Proc. R. Soc. B* **282**, 20142263.
- Aviv, A., Chen, W., Gardner, J. P., Kimura, M., Brimacombe, M., Cao, X., Srinivasan, S. R. and Berenson, G. S. (2009). Leukocyte telomere dynamics: longitudinal findings among young adults in the Bogalusa Heart Study. *Am. J. Epidemiol.* **169**, 323–329.
- Aviv, A., Hunt, S. C., Lin, J., Cao, X., Kimura, M. and Blackburn, E. (2011). Impartial comparative analysis of measurement of leukocyte telomere length/DNA content by Southern blots and qPCR. *Nucleic Acids Res.* **39**, e134.
- Baerlocher, G. M., Rice, K., Vulto, I. and Lansdorp, P. M. (2007). Longitudinal data on telomere length in leukocytes from newborn baboons support a marked drop in stem cell turnover around 1 year of age. *Aging Cell* **6**, 121–123.
- Barrett, E. L. B., Burke, T. A., Hammers, M., Komdeur, J. and Richardson, D. S. (2013). Telomere length and dynamics predict mortality in a wild longitudinal study. *Mol. Ecol.* **22**, 249–259.
- Bates, D., Maechler, M., Bolker, B. and Walker, S. (2013). lme4: Linear Mixed-effects Models Using Eigen and S4. Available at: <http://lme4.r-forge.r-project.org/>
- Beaulieu, M., Reichert, S., Le Maho, Y., Ancel, A. and Criscuolo, F. (2011). Oxidative status and telomere length in a long-lived bird facing a costly reproductive event. *Funct. Ecol.* **25**, 577–585.
- Bize, P., Criscuolo, F., Metcalfe, N. B., Nasir, L. and Monaghan, P. (2009). Telomere dynamics rather than age predict life expectancy in the wild. *Proc. R. Soc. B* **276**, 1679–1683.
- Blackburn, E. H. (2005). Telomeres and telomerase: their mechanisms of action and the effects of altering their functions. *FEBS Lett.* **579**, 859–862.
- Bolton, M. (1995). Food delivery to nestling storm petrels: limitation or regulation? *Funct. Ecol.* **9**, 161–170.
- Boonekamp, J. J., Mulder, G. A., Salomons, H. M., Dijkstra, C. and Verhulst, S. (2014). Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proc. R. Soc. B* **281**, 20133287.
- Brümmendorf, T. H. and Mak, J. (2002). Longitudinal studies of telomere length in feline blood cells: implications for hematopoietic stem cell turnover in vivo. *Exp. Hematol.* **30**, 1147–1152.
- Campisi, J. (2005). Senescent cells, tumor suppression, and organismal aging: good citizens, bad neighbors. *Cell* **120**, 513–522.
- Caprioli, M., Romano, M., Romano, A., Rubolini, D., Motta, R., Folini, M. and Saino, N. (2013). Nestling telomere length does not predict longevity, but covaries with adult body size in wild barn swallows. *Biol. Lett.* **9**, 20130340.

- Cawthon, R. M.** (2002). Telomere measurement by quantitative PCR. *Nucleic Acids Res.* **30**, e47.
- Cawthon, R. M., Smith, K. R., O'Brien, E., Sivatchenko, A. and Kerber, R. A.** (2003). Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet* **361**, 393-395.
- Criscuolo, F., Bize, P., Nasir, L., Metcalfe, N. B., Foote, C. G., Griffiths, K., Gault, E. A. and Monaghan, P.** (2009). Real-time quantitative PCR assay for measurement of avian telomeres. *J. Avian Biol.* **40**, 342-347.
- Davis, P.** (1957). The breeding of the storm petrel. *Br. Birds* **50**, 371-383.
- Epel, E. S., Blackburn, E. H., Lin, J., Dhabhar, F. S., Adler, N. E., Morrow, J. D. and Cawthon, R. M.** (2004). Accelerated telomere shortening in response to life stress. *Proc. Natl. Acad. Sci. USA* **101**, 17312-17315.
- Foote, C. G., Gault, E. A., Nasir, L. and Monaghan, P.** (2011). Telomere dynamics in relation to early growth conditions in the wild in the lesser black-backed gull. *J. Zool.* **283**, 203-209.
- Foote, C. G., Vleck, D. and Vleck, C. M.** (2013). Extent and variability of interstitial telomeric sequences and their effects on estimates of telomere length. *Mol. Ecol. Resour.* **13**, 417-428.
- Geiger, S., Le Vaillant, M., Lebard, T., Reichert, S., Stier, A., Le Maho, Y. and Criscuolo, F.** (2012). Catching-up but telomere loss: half-opening the black box of growth and ageing trade-off in wild king penguin chicks. *Mol. Ecol.* **21**, 1500-1510.
- Griffiths, R., Double, M. C., Orr, K. and Dawson, R. J. G.** (1998). A DNA test to sex most birds. *Mol. Ecol.* **7**, 1071-1075.
- Hall, M. E., Nasir, L., Daunt, F., Gault, E. A., Croxall, J. P., Wanless, S. and Monaghan, P.** (2004). Telomere loss in relation to age and early environment in long-lived birds. *Proc. R. Soc. B* **271**, 1571-1576.
- Hausmann, M. F. and Mauck, R. A.** (2008). Telomeres and longevity: testing an evolutionary hypothesis. *Mol. Biol. Evol.* **25**, 220-228.
- Hausmann, M. F., Winkler, D. W. and Vleck, C. M.** (2005). Longer telomeres associated with higher survival in birds. *Biol. Lett.* **1**, 212-214.
- Hausmann, M. F., Winkler, D. W., Huntington, C. E., Nisbet, I. C. T. and Vleck, C. M.** (2007). Telomerase activity is maintained throughout the lifespan of long-lived birds. *Exp. Gerontol.* **42**, 610-618.
- Haywood, S. and Perrins, C. M.** (1992). Is clutch size in birds affected by environmental conditions during growth? *Proc. R. Soc. B* **249**, 195-197.
- Heidinger, B. J., Blount, J. D., Boner, W., Griffiths, K., Metcalfe, N. B. and Monaghan, P.** (2012). Telomere length in early life predicts lifespan. *Proc. Natl. Acad. Sci. USA* **109**, 1743-1748.
- Herborn, K. A., Heidinger, B. J., Boner, W., Noguera, J. C., Adam, A., Daunt, F. and Monaghan, P.** (2014). Stress exposure in early post-natal life reduces telomere length: an experimental demonstration in a long-lived seabird. *Proc. R. Soc. B* **281**, 20133151.
- Horn, T., Robertson, B. C., Will, M., Eason, D. K., Elliott, G. P. and Gemmill, N. J.** (2011). Inheritance of telomere length in a bird. *PLoS ONE* **6**, e17199.
- Jennings, B. J., Ozanne, S. E., Dorling, M. W. and Hales, C. N.** (1999). Early growth determines longevity in male rats and may be related to telomere shortening in the kidney. *FEBS Lett.* **448**, 4-8.
- Kotrschal, A., Ilmonen, P. and Penn, D. J.** (2007). Stress impacts telomere dynamics. *Biol. Lett.* **3**, 128-130.
- Lindström, J.** (1999). Early development and fitness in birds and mammals. *Trends Ecol. Evol.* **14**, 343-348.
- Metcalfe, N. B. and Monaghan, P.** (2001). Compensation for a bad start: grow now, pay later? *Trends Ecol. Evol.* **16**, 254-260.
- Mizutani, Y., Tomita, N., Niizuma, Y. and Yoda, K.** (2013). Environmental perturbations influence telomere dynamics in long-lived birds in their natural habitat. *Biol. Lett.* **9**, 20130511.
- Monaghan, P.** (2010). Telomeres and life histories: the long and the short of it. *Ann. NY Acad. Sci.* **1206**, 130-142.
- Monaghan, P. and Hausmann, M. F.** (2006). Do telomere dynamics link lifestyle and lifespan? *Trends Ecol. Evol.* **21**, 47-53.
- Njajou, O. T., Cawthon, R. M., Damcott, C. M., Wu, S.-H., Ott, S., Garant, M. J., Blackburn, E. H., Mitchell, B. D., Shuldiner, A. R. and Hsueh, W.-C.** (2007). Telomere length is paternally inherited and is associated with parental lifespan. *Proc. Natl. Acad. Sci. USA* **104**, 12135-12139.
- Ogburn, C. E., Carlberg, K., Ottinger, M. A., Holmes, D. J., Martin, G. M. and Austad, S. N.** (2001). Exceptional cellular resistance to oxidative damage in long-lived birds requires active gene expression. *J. Gerontol. A* **56**, B468-B474.
- Olsson, M., Pauliny, A., Wapstra, E., Uller, T., Schwartz, T. and Blomqvist, D.** (2011). Sex differences in sand lizard telomere inheritance: paternal epigenetic effects increases telomere heritability and offspring survival. *PLoS ONE* **6**, e17473.
- Pauliny, A., Wagner, R. H., Augustin, J., Szép, T. and Blomqvist, D.** (2006). Age-independent telomere length predicts fitness in two bird species. *Mol. Ecol.* **15**, 1681-1687.
- Pfaffl, M. W.** (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45.
- R Core Team** (2013). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing. <http://www.r-project.org>
- Reichert, S., Rojas, E. R., Zahn, S., Robin, J.-P., Criscuolo, F. and Massemin, S.** (2015). Maternal telomere length inheritance in the king penguin. *Heredity* **114**, 10-16.
- Reid, J. M., Bignal, E. M., Bignal, S., McCracken, D. I. and Monaghan, P.** (2003). Environmental variability, life-history covariation and cohort effects in the red-billed cormorant *Pyrrhocorax pyrrhocorax*. *J. Anim. Ecol.* **72**, 36-46.
- Rose, K. E., Clutton-Brock, T. H. and Guinness, F. E.** (1998). Cohort variation in male survival and lifetime breeding success in red deer. *J. Anim. Ecol.* **67**, 979-986.
- Salomons, H. M., Mulder, G. A., van de Zande, L., Hausmann, M. F., Linskens, M. H. K. and Verhulst, S.** (2009). Telomere shortening and survival in free-living corvids. *Proc. R. Soc. B* **276**, 3157-3165.
- Sedinger, J. S., Flint, P. L. and Lindberg, M. S.** (1995). Environmental influence on life-history traits: growth, survival, and fecundity in black brant (*Branta bernicla*). *Ecology* **76**, 2404-2414.
- Sing, T., Sander, O., Beerenwinkel, N. and Lengauer, T.** (2005). ROCr: visualizing classifier performance in R. *Bioinformatics* **21**, 3940-3941.
- Stier, K. S., Almasi, B., Gasparini, J., Piault, R., Roulin, A. and Jenni, L.** (2009). Effects of corticosterone on innate and humoral immune functions and oxidative stress in barn owl nestlings. *J. Exp. Biol.* **212**, 2085-2091.
- Tarry-Adkins, J. L., Chen, J. H., Smith, N. S., Jones, R. H., Cherif, H. and Ozanne, S. E.** (2009). Poor maternal nutrition followed by accelerated postnatal growth leads to telomere shortening and increased markers of cell senescence in rat islets. *FASEB J.* **23**, 1521-1528.
- van de Pol, M. and Verhulst, S.** (2006). Age-dependent traits: a new statistical model to separate within- and between-individual effects. *Am. Nat.* **167**, 766-773.
- Verdun, R. E. and Karlseder, J.** (2007). Replication and protection of telomeres. *Nature* **447**, 924-931.
- von Zglinicki, T.** (2002). Oxidative stress shortens telomeres. *Trends Biochem. Sci.* **27**, 339-344.
- Watson, H.** (2014). The occurrence of hypothermia in nestlings of the European Storm-petrel *Hydrobates pelagicus*. *Seabird* **26**, 106-109.
- Watson, H., Bolton, M. and Monaghan, P.** (2014). Out of sight but not out of harm's way: human disturbance reduces reproductive success of a cavity-nesting seabird. *Biol. Conserv.* **174**, 127-133.
- Zeichner, S. L., Palumbo, P., Feng, Y., Xiao, X., Gee, D., Sleasman, J., Goodenow, M., Biggar, R. and Dimitrov, D.** (1999). Rapid telomere shortening in children. *Blood* **93**, 2824-2830.