



Figure S1. Related to Figure 2. Comparison of hairiness of the tergites of the abdomen of uninfested (left) and stylopized females of *Andrena vaga* (right). Scale bar: 1 mm.




Figure S2. Distance between hairs of the abdominal tergites II–IV of the abdomen of stylopized and uninfested females of *Andrena vaga*.




Figure S3. Tarsus of *Stylops ovinae* (A) and hairiness of the abdominal tergites II–IV of a stylopized *Andrena vaga* (B–D), SEM-micrographs. (A) Prothoracic leg, ventral view (modified from Pohl & Beutel, 2004). (B) Abdominal tergite II. (C) Abdominal tergite III. (D) Abdominal tergite IV. Scale bar: 100 µm.




Figure S4. *Mengenilla moldrzyki* (Mengenillidae), male, tarsus of metathoracic leg, ventral view, SEM-micrograph (modified from Pohl & Beutel, 2004). Scale bar: 100 µm.

Table S1. Distance (μm) between hairs of abdominal tergites II–IV of stylopized and uninfested *Andrena vaga*.

	Stylopized				Uninfested				
	average	med.	max.	min.	average	med.	max.	min.	
Tergite II	55.6	54.8	91.3	27.7	Tergite II	63.1	62.3	155.2	28.3
Tergite III	43.9	41.7	126.5	13.4	Tergite III	98.5	80.7	286.9	34.3
Tergite IV	30.3	26.5	136.7	11.4	Tergite IV	111.1	94.3	350.5	34.3