



**Cover:** A clutch of first-stage stomatopod larvae (*Lysiosquillina maculata*). These larvae are illuminated in such a way to make both their transparent bodies as well as their green-reflective 'eyeshine' highly visible on a black background. In this issue, Feller and Cronin (pp. 3263–3273) examine properties of stomatopod larval eyeshine, concluding it serves as a novel pelagic camouflage reflective strategy spectrally matched to the natural light environment. Although eyeshine may appear highly visible on a black background, in nature, eye structures disguise the opaque retinas so that invisibility achieved by the larvae's transparent body tissue is minimally compromised. Photo credit: K. Feller.

## INSIDE JEB

- 3187** Transparent larvae hide eyes behind reflections; Lunar explorers will walk at higher speeds than thought; Parental lifestyle influenced eggshell evolution; Take-off with train is no drag for peacocks; Rats whisk whiskers to sense surroundings

## CLASSICS

- 3191** Hughes and Shelton: the fathers of fish respiration  
**Randall, D.**

## SHORT COMMUNICATIONS

- 3193** Patterns of fuel use during locomotion in mammals revisited: the importance of aerobic scope  
**Schippers, M.-P., LeMoine, C. M. R. and McClelland, G. B.**

- 3197** Host orientation using volatiles in the phoretic nematode *Caenorhabditis japonica*  
**Okumura, E. and Yoshiga, T.**

- 3200** The preferred walk to run transition speed in actual lunar gravity  
**De Witt, J. K., Edwards, W. B., Scott-Pandorf, M. M., Norcross, J. R. and Gernhardt, M. L.**

## METHODS & TECHNIQUES

- 3204** Use of a gyroscope/accelerometer data logger to identify alternative feeding behaviours in fish  
**Kawabata, Y., Noda, T., Nakashima, Y., Nanami, A., Sato, T., Takebe, T., Mitamura, H., Arai, N., Yamaguchi, T. and Soyano, K.**

## RESEARCH ARTICLES

- 3209** Gaze characteristics of freely walking blowflies *Calliphora vicina* in a goal-directed task  
**Kress, D. and Egelhaaf, M.**

- 3221** Social regulation of cortisol receptor gene expression  
**Korzan, W. J., Grone, B. P. and Fernald, R. D.**

- 3229** A context-dependent alarm signal in the ant *Temnothorax rugatulus*  
**Sasaki, T., Hölldobler, B., Millar, J. G. and Pratt, S. C.**

- 3237** The elaborate plumage in peacocks is not such a drag  
**Askew, G. N.**

- 3242** Spatial co-ordination of foot contacts in unrestrained climbing insects  
**Theunissen, L. M., Vikram, S. and Dürr, V.**

- 3254** Female gray treefrogs, *Hyla versicolor*, are responsive to visual stimuli but unselective of stimulus characteristics  
**Reichert, M. S., Galante, H. and Höbel, G.**

- 3263** Hiding opaque eyes in transparent organisms: a potential role for larval eyeshine in stomatopod crustaceans  
**Feller, K. D. and Cronin, T. W.**

- 3274** Exercising for food: bringing the laboratory closer to nature  
**Fonseca, I. A. T., Passos, R. L. F., Araújo, F. A., Lima, M. R. M., Lacerda, D. R., Pires, W., Soares, D. D., Young, R. J. and Rodrigues, L. O. C.**

- 3283** Physiological mechanisms underlying individual variation in tolerance of food deprivation in juvenile European sea bass, *Dicentrarchus labrax*  
**McKenzie, D. J., Vergnet, A., Chatain, B., Vandeputte, M., Desmarais, E., Steffensen, J. F. and Guinand, B.**

- 3293** Niche-specific cognitive strategies: object memory interferes with spatial memory in the predatory bat *Myotis nattereri*  
**Hulgard, K. and Ratcliffe, J. M.**

- 3301** Is warmer better? Decreased oxidative damage in notothenioid fish after long-term acclimation to multiple stressors  
**Enzor, L. A. and Place, S. P.**

- 3311** High paracellular nutrient absorption in intact bats is associated with high paracellular permeability in perfused intestinal segments  
**Brun, A., Price, E. R., Gontero-Fourcade, M. N., Fernandez-Marinone, G., Cruz-Neto, A. P., Karasov, W. H. and Caviedes-Vidal, E.**

- 3318** Fine-tuned echolocation and capture-flight of *Myotis capaccinii* when facing different-sized insect and fish prey  
**Aizpurua, O., Aihartza, J., Alberdi, A., Baagøe, H. J. and Garin, I.**

- 3326** Nesting behaviour influences species-specific gas exchange across avian eggshells  
**Portugal, S. J., Maurer, G., Thomas, G. H., Hauber, M. E., Grim, T. and Cassey, P.**

- 3333** Mechanical processing via passive dynamic properties of the cockroach antenna can facilitate control during rapid running  
**Mongeau, J.-M., Demir, A., Dallmann, C. J., Jayaram, K., Cowan, N. J. and Full, R. J.**

- 3346** Scavenging ROS dramatically increase NMDA receptor whole-cell currents in painted turtle cortical neurons  
**Dukoff, D. J., Hogg, D. W., Hawrysh, P. J. and Buck, L. T.**

- 3356** Physiological vagility and its relationship to dispersal and neutral genetic heterogeneity in vertebrates  
**Hillman, S. S., Drewes, R. C., Hedrick, M. S. and Hancock, T. V.**

- 3365** The search space of the rat during whisking behavior  
**Huet, L. A. and Hartmann, M. J. Z.**

- 3377** K<sup>+</sup> absorption by locust gut and inhibition of ileal K<sup>+</sup> and water transport by FGLamide allatostatins  
**Robertson, L., Donini, A. and Lange, A. B.**